

Discovery of a Novel Series of Tankyrase Inhibitors by a Hybridization Approach

Upendra Rao Anumala, Jo Waaler, Yves Nkizinkiko, Alexander Ignatev, Katina Lazarow, Peter Lindemann, Petter Angell Olsen, Sudarshan Murthy, Ezeogo Obaji, Alexander G. Majouga, Sergey V. Leonov, Jens Peter von Kries, Lari Lehtiö, Stefan Krauss, and Marc Nazaré

J. Med. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.7b00883 • Publication Date (Web): 20 Nov 2017

Downloaded from <http://pubs.acs.org> on November 20, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Discovery of a Novel Series of Tankyrase Inhibitors by a Hybridization Approach

Upendra Rao Anumala,^{†,‡} Jo Waaler,[§] Yves Nkizinkiko,[¥] Alexander Ignatev[¥] Katina Lazarow,[†] Peter Lindemann[†], Petter Angell Olsen,[§] Sudarshan Murthy,[¥] Ezeogo Obaji,[¥] Alexander G. Majouga,[∇] Sergey Leonov,^{⊥,◇} Jens Peter von Kries,^{†,‡} Lari Lehtiö,[¥] Stefan Krauss,^{§} and Marc Nazaré^{*†,‡}*

[†]Leibniz-Forschungsinstitut für Molekulare Pharmakologie (FMP), Campus Berlin-Buch,
13125 Berlin, Germany.

[‡]Berlin Institute of Health (BIH), Anna-Louisa-Karsch-Str. 2, 10178 Berlin, Germany.

[§]Unit for Cell Signaling, Institute of Medical Microbiology, Oslo University Hospital,
Gaustadalleen 34, 0372 Oslo, Norway and Hybrid Technology Hub - Centre of Excellence,
Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway.

[¥]Faculty of Biochemistry and Molecular Medicine, Biocenter Oulu, University of Oulu, PO
Box 5400, 90014 Oulu, Finland.

[∇]Moscow State University, Chemistry Department, Leninskie Gory 1/3, Moscow 119991,
Russia.

[⊥]National University of Science and Technology MISiS, Leninsky Avenue 4, Moscow
119049, Russia.

[◇]Moscow Institute of Physics and Technology (State University), Institutskiy Lane 9, 141700
Dolgoprudny, Russia.

ABSTRACT

A structure-guided hybridization approach using two privileged substructures gave instant access to a new series of tankyrase inhibitors. The identified inhibitor **16** displays high target affinity on tankyrase 1 and 2 with a biochemical and cellular IC₅₀ values of 29 nM, 6.3 nM and 19 nM, respectively, and high selectivity towards other Poly(ADP-ribose) polymerase enzymes. The identified inhibitor shows a favorable in-vitro ADME profile as well as good oral bioavailability in mice, rats and dogs. Critical for the approach was the utilization of an appropriate linker between 1,2,4-triazole and benzimidazolone moieties, whereby a cyclobutyl linker displayed superior affinity compared to a cyclohexane and phenyl linker.

Introduction

Catalytic modification of proteins using the redox metabolite nicotinamide adenine dinucleotide (NAD⁺) as a substrate to successively add ADP-ribose moieties onto the target protein is known as PARsylation.¹ In 1960s, this posttranslational protein modification was reported with the identification of PARP1 and its role in DNA repair.²⁻⁴ There are two classes of PARP enzymes: mono (ADP-ribosyl)ating and (oligo-)poly(ADP-ribosyl)ating proteins.⁵⁻⁶ The poly-ADP-ribose polymerases tankyrase 1 and tankyrase 2 (TNKS1, PARP5a and TNKS2, PARP5b) share almost 82% sequence identity and show predominantly cytoplasmic differential expression in a variety of tissues.⁷ Both tankyrase paralogs have three functional segments: the carboxy terminal catalytic ARTD domain (ADP-ribosyltransferase with Diphtheria toxin homology), five ankyrin domains involved in protein/protein interactions with target proteins such as AXIN, NuMA, TRF1, GRB and IRAP⁸ and a sterile alpha motif (SAM) domain that is involved in tankyrase polymerization and supports the catalytic activity of the PARP domain.⁹ The catalytic ARTD domains of both tankyrases show high similarity with 89% sequence identity.

1
2
3 Tankyrases can control in a context dependent manner several cellular pathways including,
4
5 the WNT/ β -catenin signaling pathway that executes key functions in embryonic
6
7 development, stem cell biology, cell fate specification, energy metabolism and cell migration.
8
9 Other roles of tankyrases comprise the involvement in e.g. mitosis, and cherubism.^{6, 10-14}
10
11 Tankyrase inhibition may have therapeutic potential in several diseases like selected cancers
12
13 such as colorectal carcinoma and non-small cell lung cancer as well as in fibrotic diseases,
14
15 and herpes simplex virus (HSV) infections.¹⁵⁻¹⁷
16
17

18 Several tankyrase inhibitors emerging from high throughput screening or chemical
19
20 optimization efforts have recently been reported and inhibitor binding modes in the active site
21
22 has been effectively elucidated by X-ray crystallography and complemented by in-silico
23
24 docking studies.^{13, 18-28} Two distinct binding sites, the nicotinamide and the adenosine sub-
25
26 pocket have been described to accommodate small molecule ligands. Compound **1**
27
28 (**XAV939**), binding to the nicotinamide sub-pocket, was the first reported tankyrase inhibitor
29
30 which showed an impact on WNT/ β -catenin signaling by stabilizing AXIN and decreasing β -
31
32 catenin levels.¹³ Due to the conservation of the nicotinamide sub-pocket throughout the entire
33
34 PARP family, **1**, and related inhibitors **2-4** show variable selectivity towards the other
35
36 members or the PARP family.
37
38
39

40 In contrast to the nicotinamide binding site, the architecture of the adenosine binding cavity
41
42 is more varied between PARPs. Consequently, inhibitors addressing the adenosine sub-
43
44 pocket of TNKS1/2 appear to be intrinsically more selective over the other members of the
45
46 PARP family. Notably only tankyrase has been reported to be effectively inhibited by a
47
48 compound which is solely binding to this sub-pocket. Recently, dual site tankyrase inhibitors
49
50 that span both, the nicotinamide and the adenosine binding site have shown to be potent and
51
52 tankyrase selective.^{22, 11, 26-28} Examples of reported tankyrase inhibitors are given in Figure 1.
53
54
55
56
57
58
59
60

1
2
3 We previously reported the discovery of, a 1,2,4-triazole based specific tankyrase inhibitor
4
5 **5** (JW74) which culminated in the development of **6** (G007-LK) which inhibits TNKS1/2 *in*
6
7 *vitro* and *in vivo* with high specificity.¹⁸ However, albeit being highly potent and selective,
8
9 and showing an excellent oral bioavailability in mice, **6** had poor PK in rats, hampering its
10
11 further preclinical development. Here we report the design of a new class of tankyrase
12
13 inhibitors based on a hybridization approach of privileged fragments from two distinct
14
15 inhibitor series¹⁸ **6** and compound **8** described by Bregman et al..²⁰ We show that the lead
16
17 compound of the series shows a further improved affinity and cellular activity, good
18
19 solubility, good target specificity within the PARP family, low adverse inhibition in a kinase
20
21 panel, good oral bioavailability in mouse, rat and dog as well as efficacy in mouse xenograft
22
23 models.
24
25




Figure 1. Examples of chemical structures of known tankyrase inhibitors; A) Inhibitors addressing the nicotinamide pocket (**1**¹³, **2**²³, **3**¹², **4**²¹), B) Inhibitors addressing the adenosine pocket (**5**¹⁸, **6**¹⁸, **7**²⁴, **8**²⁰, **9**²⁰, **10**¹⁹), C) Dual inhibitors addressing the adenosine and nicotinamide pocket (**11**²⁶, **12**²², **13**²⁷).

Results and Discussion

Design of the Hybrid Inhibitors.

Our previous studies of the diaryl 1,2,4-triazole series yielded potent and selective tankyrase inhibitors.^{18, 29} However, the lead compound **6** suffered from an extended, highly conjugated, aromatic system incorporating a vinylic bond in combination with the intrinsically high lipophilicity, a negligible low Fsp³ content and a critically high molecular weight for some derivatives. Collectively, this suggested that a further classical optimization with incremental structural changes may not solve these issues. We speculated that due to the resemblance of the N-disubstituted glycine pharmacophore in **8** with a diaryl 1,2,4-triazole the distal benzimidazolone could be grafted on this scaffold with the appropriate spacer. Thus, we resorted to a hybridization approach by joining the diaryl substituted 1,2,4-triazole and the benzimidazolone from the two inhibitors **6** and **8** (Figure 2). The available co-crystal structures of TNKS2-**6** and TNKS1-**8** analogs were used to guide the hybrid and linker design (Figure 3 a-c). In these crystal structures the hydrogen bonding pattern is conserved and importantly the atoms of the triazole and glycine moieties superpose well and the vinyl and cyclohexane linkers occupy the same space in the pocket (Figure 3c). An appropriate linker for the hybridization approach would thus enable us to preserve the overall binding mode and maintain distinct interaction anchor points of the merged fragments. Three different linkers were chosen to provide the appropriate distance and conformational adaptability within this new class of tankyrase inhibitors (Figure 2). Design was based on the parent compound crystal structures and docking was used to check compatibility with the binding pocket. A phenyl linker group present in **14** has been used in many tankyrase inhibitors at this position (Figure 1, compounds **7**, **9**, **11**), while a saturated cyclohexyl in **15** is equidistant but provides

more flexibility to the linker. A shorter and more rigid 1,3-trans substituted cyclobutyl (**16**) was designed as a novel linker in the hybrid compound (Figure 2).




Figure 2. Hybridization design strategy for the new tankyrase inhibitors **14-16**.





Figure 3. Co-crystal structures of TNKS1/2 and inhibitors. (a) Binding mode of **6** in TNKS2 catalytic domain (PDB: 4HYF). (b) Binding mode of **8** in TNKS1 catalytic domain (PDB: 4K4E). (c) superposition of **6** and **8** co-crystal structures. Only TNKS2 protein is shown for clarity. (d) Co-crystal structure of **14** with TNKS2 (PDB: 5NSP Sigma A weighted $2F_o - F_c$ electron density map around the ligand is contoured at 1.0σ). (e) Co-crystal structure of **16** with TNKS2. Sigma A weighted $2F_o - F_c$ electron density map around the ligand is contoured at 1.5σ . (PDB: 5NOB). (f) Superposition of **6** and **16** co-crystal structures showing the compounds and TNKS2 protein corresponding to TNKS2-**6** co-crystal. The black dash lines represent hydrogen bonds and the red spheres represent water molecules.

Synthesis of the Designed Tankyrase Inhibitors.

The phenyl derivative **14** was synthesized by coupling of pyrimidyl imidohydrazide derivative **20** with the benzimidazolone acid **22** in a key condensation step. The pyrimidyl imidohydrazide derivative **20** was obtained via activation of the amide as imidoyl chloride intermediate **19** and subsequent substitution with hydrazine hydrate. This building block **20**

was then coupled with benzimidazolone acid **22** using EDCI and cyclized by dehydration in refluxing toluene to afford the phenyl triazole **14** (Scheme 1).

Scheme 1. Synthesis of compound **14**




Reagents and conditions: (a) HATU, DIPEA, dichloromethane, rt, 14 h, 68%; (b) PCl_5 , POCl_3 , Toluene, reflux, 7h, not isolated; (c) $\text{N}_2\text{H}_4 \cdot \text{H}_2\text{O}$, THF, 3 h, 48%; (d) SnCl_2 , methanol, reflux, 45 min, 92%; (e) triphosgene, dichloromethane, rt, 24 h, then reflux, 24 h, 72%; (f) LiOH, THF:H₂O (4:1), rt, overnight, 94%; (g) **20**, EDCI·HCl, HOBt, triethylamine, THF, rt, 72 h, 8%; (h) Toluene, reflux, 72 h, 31%.

The cyclohexyl and cyclobutyl derivatives **15**, **15a**, and **16**, **16a** were synthesized by analogous convergent routes. Carbimidothioate building block **26** was prepared from N-(2-chlorophenyl)pyrimidine-4-carboxamide **24** by thionation with Lawesson's reagent to afford **25**, subsequent methylation with methyl tosylate in the presence of potassium *t*-butoxide resulted in **26**. The benzimidazolone ester intermediates (**30a-d**) were constructed using a three-step sequence consisting of nucleophilic aromatic substitution, reduction and cyclization. The nucleophilic aromatic substitution reaction of 4-fluoro-3-nitrobenzonitrile

(**28a**), as well as for 1-fluoro-2-nitrobenzene (**28b**), with **27a-b** respectively, was followed by a hydrogenation under standard conditions in the presence of palladium on charcoal to afford the corresponding diamines. Treatment with triphosgene yielded the benzimidazolone ester derivatives **30a-d**. These esters **30a-d** were then converted into the hydrazide **31a-d** using hydrazine hydrate and were then condensed with the carbimidothioate **26** to afford the final triazole derivatives **15**, **15a** and **16**, **16a**.

Scheme 2. Synthesis of compounds **15**, **15a** and **16**, **16a**




Reagents and conditions: (a) Lawesson's reagent, toluene, 7h, reflux, 63%; (b) *t*-BuOK, methyl tosylate, THF, rt, 18 h, 94%; (c) DIPEA, acetonitrile, reflux, 24 h, **29a**: 77%, **29b**: 91%, **29c**: 96%, **29d**: 95%; (d) Pd/C, H₂, EtOH, 2 h, a 86%, b 76%, c 72%, d 76%; (e)






1
2
3 triphosgene, dichloromethane, rt 24 h, reflux 24 h, **30a**: 93%, **30b**: 92%, **30c**: 95%, **30d**:
4 92%; (f) N₂H₄·H₂O, methanol, 20 °C, 20 h, **31c**: 64%; (g) N₂H₄·H₂O, ethanol, 80 °C, 3 h, **31a**:
5 90%, **31b**: 92%, **31d**: 90%; (h) **26**, TFA, DMA, 120 °C, 14 h, **15**: 10%, **15a**: 21%, **16**: 15%,
6 **16a**: 16%.
7
8
9

10 **Biochemical and Cellular Activity of the Inhibitors.**

11
12 The five hybrid compounds were tested for their ability to inhibit TNKS2 in a biochemical
13 assay and in the human embryonic kidney HEK293 as well as the in human colon SW480
14 cell line as functional assays of the WNT/ β -catenin signaling pathway. While compound **14**
15 with a phenyl linker displayed moderate activity in the biochemical assays (TNKS2: IC₅₀ 340
16 nM) only a weak activity was measured in the cellular assays. The trans-cyclohexane linker
17 in **15a** displayed moderate affinity versus tankyrase in the biochemical assay (TNKS2: IC₅₀
18 430 nM), however, it showed moderate activity in the cellular assay (HEK293: IC₅₀ 1.06 μ M;
19 SW480: IC₅₀ 1.8 μ M). By incorporation of a nitrile moiety at the benzimidazolone in **15**, a
20 10-fold increase in cellular potency (HEK 293: IC₅₀ 0.12 μ M; SW480: IC₅₀ 1.49 μ M) was
21 observed, while the biochemical potency was only slightly improved. A *trans* configured
22 cyclobutane linker in **16a** improved the cellular IC₅₀ by an order of magnitude compared to
23 **15a** (HEK293: IC₅₀ 0.2 μ M; SW480: IC₅₀ 0.41 μ M). When incorporating a nitrile group in **16**
24 a further 10-fold increase of activity was observed (HEK293: IC₅₀ 19 nM; SW480: IC₅₀ 70
25 nM) accompanied by a favorable biochemical IC₅₀ (TNKS2: IC₅₀ 6.3 nM), suggesting a
26 preserved overall binding mode with similar underlying contacts for **15** and **16** which was
27 subsequently confirmed by X-ray crystallography (Figure 3).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46
47 **Table 1.** Activity data in biochemical and cellular assays of **14-16**
48



Cmpd	R	L	TNKS2 IC ₅₀ (μM)	ST-Luc/Ren (HEK293) IC ₅₀ (μM)	ST-Luc/Ren (SW 480) IC ₅₀ (μM)
8 ¹⁸ (Reference)	-	-	0.025	0.05	4.2
14	-CN		0.34	>10	N.D.
15a	-H		0.43	1.06	1.80
15	-CN		0.33	0.12	1.49 ^a
16a	-H		0.0098	0.2	0.41
16	-CN		0.0063	0.019	0.070

^a partial precipitation under assay conditions.

Binding Mode of the Inhibitors **14** and **16**.


We obtained the co-crystal structures of **14** and **16** with the TNKS2 catalytic domain and these confirmed our hypothesis of the binding mode of these hybrid compounds. Both compounds were clearly visible in the electron density although the soaking approach used for obtaining **14** co-crystals was detrimental to the apo crystals and resolution of this data set was lower (Supplementary Table S1). Electron density for the compound is clear, but the density for the CN group is essentially missing (Figure 3d). **14** was also only found in one out of two TNKS2 proteins present in the crystallographic asymmetric unit. In comparison to co-crystal structure of **6** (Figure 3a), the pyrimidine of **14** has rotated and forms a hydrogen bond to a water molecule. This feature is also conserved in the binding mode of **16** (Figure 3e). The rigid phenyl linker in **14** positions the triazole deeper in the pocket and closer to the nicotinamide sub pocket. Despite the conserved hydrogen bond to Tyr1060 (TNKS2) backbone amide the distorted placement on this side of the compound is likely causing the observed lower potency (Table 1). The shorter linker found in **16** directs the triazole moiety in a slightly different orientation than in **14** and the binding mode is more similar to the

template compound **6** (Figure 3f). The pyrimidine is in a better position to efficiently form a displaced π - π stacking interaction with the sidechain of Tyr1060 yielding a better potency over the other tested linker moieties.

Profile of Tankyrase Inhibitor **16**.

To assess the biotarget specificity of **16**, biochemical inhibition of a panel of human PARP family members was tested. Potency for TNKS1 (29 nM) was slightly lower than for TNKS2 (6.3 nM) and none of the other tested PARP enzymes were inhibited by **16** (Table 2).

Table 2. Summary of activity and selectivity data as well as physicochemical and ADME properties of the tankyrase inhibitor **16**. For further details see Supporting Information.



Molecular weight	468.9
cLogP	3.39
Hydrogen bond donors	1
Hydrogen bond acceptors	6
Number of rotatable bonds	4
PSA	112.6 Å ²
Aqueous solubility (pH =7.4, 25°C)	31.2 μM
Ligand efficiency (LE)	0.34
IC ₅₀ TNKS1 (pIC ₅₀ ± SD) ^c	0.029 μM (7.54 ± 0.07)
IC ₅₀ TNKS2 (pIC ₅₀ ± SD) ^c	0.0063 μM (8.20 ± 0.03)
IC ₅₀ (ST-Luc/Ren in HEK293)	0.019 μM
IC ₅₀ (ST-Luc/Ren in SW480)	0.070 μM
IC ₅₀ ARTD1/PARP1	> 100 μM

IC ₅₀ ARTD2/PARP2	> 100 μM
IC ₅₀ ARTD3/PARP3	> 100 μM
IC ₅₀ ARTD4/PARP4	> 100 μM
IC ₅₀ ARTD7/PARP15	> 10 μM ^a
IC ₅₀ ARTD8/PARP14	> 10 μM ^a
IC ₅₀ ARTD10/PARP10	> 10 μM ^a
IC ₅₀ ARTD12/PARP12	> 10 μM ^a
IC ₅₀ ARTD15/PARP16	> 10 μM ^a
Kinase selectivity: kinases with >50% inhibition at 10 μM	4/320
Clearance human liver microsomes (Clint)	14.8 (μL/min/mg protein)
Clearance human hepatocyte (Clint)	6.44 (μL/min/10 ⁶ cells)
MDCK-MDR1 permeability	5.3 × 10 ⁻⁷ cm/s (A-B)
Papp at 10 μM	28.2 × 10 ⁻⁷ cm/s (B-A)
CYP (3A4/2C9/2C19/2D6/1A)	1.3 μM/11.9 μM/>25 μM/>25 μM/>25 μM
Mouse PK (p.o., 5 mg/kg)	F 47%; t _{1/2} 1.5 h; C _{max} 123.5 ng/mL; AUC(0-t) 144.7 hr×ng/mL; CL 34.02 L/kg
Rat PK (p.o., 14 mg/kg)	F 35%; t _{1/2} N.A.; C _{max} 843 ng/mL; AUC(0-t) 2765 hr×ng/mL; CL N.A.
Dog PK (p.o., 7 mg/kg)	F 91%; t _{1/2} 4.7 h; C _{max} 5851 ng/mL; AUC(0-t) 27134 hr×ng/mL; CL 0.14 L/kg

^aNo inhibition; concentration limited by low DMSO tolerance of the enzymes. ^bCalculated properties using ChemAxon package version 16.1., 2010-2016. ^cSD = standard deviation.

Furthermore, a kinase selectivity profiling on 320 wild-type protein kinases of **16** at 10 μM concentration was carried out revealing only 4 kinase hits with > 50% inhibition at 10 μM concentration (CLK2: 73%; MELK: 70%; PRKG1: 66% and TSF1: 52%), but no kinase with a > 90% inhibition (Supplementary Table S2). Cytochrome P450 inhibition was tested with a panel of CYP isoforms and *in vitro* metabolic stability was evaluated in human hepatocytes

and human liver microsomes. Of the tested CYP isoforms, 3A4 and 2C9 were significantly inhibited possibly due to the pyrimidine ring in **16**, while CYP2C19, CYP1A and CYP2D6 were not affected. Human hepatocyte clearance (Cl_{int}) was modest at 6.44 (μL/min/10⁶ cells) and 14.8 (μL/min/mg protein), respectively (Table 2). MDCK-MDR1 permeability of **16** showed a moderate influx ratio and a high mean efflux ratio leading to a significant efflux ratio of 53.2 indicating an active efflux and thus a low predicted blood-brain-barrier penetration (Table 2). However, an overall good oral bioavailability in mouse (F: 47%), rat (F: 35%) and dog (F: 91%) including a surprisingly low compound excretion in urine and feces in rat, underscored the suitability of **16** as a chemical tool for pharmacological in-vivo evaluation.




Figure 4. Anti-tumor activity of **16** in xenograft models. A) COLO320 colon cancer xenograft B) isogenic p388 leukemia mice model. Reduction of tumor volume (mm³) versus vehicle treated controls (blue) after once daily oral dosing of **16** at various depicted doses. Statistical significance is indicated: ANOVA on Ranks/Dunn's method, P < 0.05 (*), One Way ANOVA/Holm-Sidak method, P < 0.001 (**), and One-tailed P-value < 0.05 (***).

1
2
3 To evaluate the anti-tumor effects of **16** *in vivo*, we established xenografts using the human
4 colorectal cancer cell line COLO 320DM cells in male Balb/c nude mice. The tumor-bearing
5 mice were randomized into 4 treatment groups, 2 days after inoculation: i) vehicle control
6 (1% starch, n = 4), ii) 15 mg/kg **16** (n = 5), iii) 30 mg/kg **16** (n = 5) and iv) 60 mg/kg **16** (n =
7 5). After 10 days of once daily oral drug-administration, and caliper-based tumor size
8 measurements on day 12, 17 and 21, the experiment was terminated. Compared to vehicle
9 control, treatment with **16** resulted in 53%, 63% and 63% statistically significant tumor size
10 reductions at 15 mg/kg, 30 mg/kg and 60 mg/kg once daily oral administration, respectively
11 (Figure 4A). In a second tumor model, we used the syngeneic leukemic p388 mouse model.
12 Immunocompetent BDF1 (DBA2×C57Bl6j) mice were implanted with p388 cells and
13 randomized into 3 treatment groups consisting of 6 mice each after 2 days: Vehicle (1%
14 starch, n = 6) and two treatment groups for **16**: 10 mg/kg (n = 5) and 30 mg/kg (n = 5). After
15 10 days of once daily oral administration, the tumors sizes were measured using caliper and
16 the experiment was ended. Compared to vehicle control, treatment with **16** resulted in 32%
17 and 57% statistically significant tumor size reductions for 15 mg/kg and 30 mg/kg dosing,
18 respectively (Figure 4B). No animal discomforts or body weight differences were registered
19 throughout the experiment period. Collectively, the results show that **16** can significantly
20 decrease the growth of colorectal cancer and leukemia in immunodeficient and
21 immunocompetent models, respectively.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 CONCLUSIONS

47
48 A structure guided hybridization approach based on two known inhibitors allowed us to
49 design successfully a novel series of tankyrase inhibitors. The lead compound **16** shows high
50 selectivity towards TNKS1/2, enhanced IC₅₀ in biochemical assays (IC₅₀: TNKS1 29 nM,
51 TNKS2 6.3 nM) and *in vitro* cellular assays (HEK293: IC₅₀ 19 nM; SW480: IC₅₀ 70 nM). In
52
53
54
55
56
57
58
59
60

1
2
3 addition, good pharmacokinetic properties in mice, and dogs and efficacy in a tumor
4 xenograft model were achieved. The novel tankyrase inhibitor **16** expands the available small
5 molecules space to selectively inhibit TNKS1/2 *in vitro* and *in vivo*.
6
7
8
9

10 11 12 13 **EXPERIMENTAL SECTION**

14
15 **Chemistry.** All chemicals were purchased from commercial suppliers: Activate Scientific,
16 Sigma-Aldrich and Alfa Aesar and used as received unless otherwise specified. NMR spectra
17 were recorded at either 295 K (300 MHz) or 300 K (600 MHz) at either Bruker AV 300 (300
18 MHz, 75 MHz) or Bruker AV 600 (600 MHz, 151 MHz) spectrometers. Chemical shifts are
19 reported in ppm (δ) referenced to TMS ($\delta = 0.00$ ppm), DMSO (2.50 ppm) and CHCl_3 (7.26
20 ppm). Melting points were recorded in open capillaries on a Büchi B-545 Melting Point
21 Apparatus. Temperatures are expressed in degrees Celsius ($^{\circ}\text{C}$) and are uncorrected.
22
23
24
25
26
27
28
29

30
31 LC/MS analysis was performed on an Agilent LC/MS 1260 analytical HPLC with DAD
32 coupled to an Agilent 6120 single quadrupole mass spectrometer (ESI-SQ) equipped with a
33 Thermo Fisher Scientific Accucore C18 column, 2.1 x 30 mm, 2.6 μm . Method: ESI+, flux:
34 0.8 ml/min, 5-95% CH_3CN in H_2O + 0.1% FA, total runtime: 2.5 min. High resolution mass
35 spectra were recorded on an Agilent 6220A accurate-mass time-of-flight mass spectrometer
36 (ESI-TOF) with Agilent 1200 HPLC/DAD front-end. The HPLC was equipped with an
37 Agilent Poroshell 120, C18 column, 2.1 x 100 mm, 1.8 μm . Method: ESI+, flux: 0.6 ml/min,
38 5-99% CH_3CN in H_2O + 0.1% FA, total runtime: 4.5 min.
39
40
41
42
43
44
45
46
47

48 Purity and characterization of all final compounds was established by a combination of LC-
49 MS, LC-HRMS and NMR analytical techniques. All compounds were found to be >95% pure
50 by LC-MS and LC-HRMS analysis.
51
52
53
54
55
56
57
58
59
60

Preparation of 1-(4-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-yl)phenyl)-2,3-dihydro-2-oxo-1H-benzo[d]imidazole-5-carbonitrile (14).

(i) **N-(2-Chlorophenyl)pyrimidine-4-carboxamide (24).** To a flask charged with pyrimidine-4-carboxylic acid (**17**) (1.4 g, 11.28 mmol) were added dichloromethane (50 ml), N,N-diisopropylethylamine (5.89 ml, 33.84 mmol), HATU (4.75 g, 12.4 mmol) and 2-chlorobenzenamine (**18**) (1.18 ml, 11.28 mmol) respectively. The resulting mixture was stirred overnight at room temperature. Water was added to reaction mixture and the organic layer was separated using a separating funnel. The organic layer was dried over magnesium sulfate, filtered and concentrated in vacuo, and subjected to column chromatography on silica gel using a cyclohexane and ethyl acetate gradient as eluent to afford N-(2-chlorophenyl)pyrimidine-4-carboxamide (**24**) as an amorphous solid. Yield 1.79 g (68%). MS (ESI) for $C_{11}H_8ClN_3O + H^+$, $[M+H]^+$ m/z, calcd 234.0, found 234.1. 1H NMR (300 MHz, $CDCl_3$) δ 10.57 (s, 1H), 9.37 (s, 1H), 9.05 (d, $J = 5.0$ Hz, 1H), 8.61 (dd, $J = 8.2, 1.6$ Hz, 1H), 8.22 (d, $J = 5.0$ Hz, 1H), 7.44 (dd, $J = 8.1, 1.5$ Hz, 1H), 7.35 (td, $J = 7.9, 1.6$ Hz, 1H), 7.19 – 7.06 (m, 1H). ^{13}C NMR (75 MHz, $CDCl_3$) δ 160.3, 159.4, 157.8, 156.1, 133.9, 129.2, 127.7, 125.3, 123.6, 121.1, 118.5.

(ii) **(Z)-N''-(2-Chlorophenyl)pyrimidine-4-carboximidhydrazide (20).** To a solution of N-(2-chlorophenyl)pyrimidine-4-carboxamide (**24**) (450 mg, 1.92 mmol) in toluene (20 ml) was added phosphorous pentachloride (1.2 g, 5.77 mmol) and phosphoryl chloride (540 μ l, 5.77 mmol) and the mixture stirred at reflux for 7 hours. The solvents were then evaporated to achieve (Z)-N-(2-chlorophenyl)pyrimidine-4-carbimidoyl chloride (**19**) as intermediate and 3 ml of hydrazine hydrate was added and stirred for 4 hours at room temperature. Solvents were evaporated and subjected to column chromatography on silica gel using a cyclohexane and ethyl acetate gradient as eluent to afford (Z)-N''-(2-chlorophenyl)pyrimidine-4-carboximidhydrazide (**20**) as a yellow foam. Yield: 230 mg

1
2
3 (48%) ^1H NMR (300 MHz, DMSO- d_6) δ 9.05 – 8.98 (m, 1H), 8.66 (d, J = 5.5 Hz, 1H), 7.86
4 (dd, J = 5.5, 1.0 Hz, 1H), 7.44 – 7.35 (m, 1H), 7.30 (s, 2H), 7.18 – 7.04 (m, 2H), 6.87 – 6.72
5 (m, 1H), 6.22 (d, J = 7.9 Hz, 1H). ^{13}C NMR (75 MHz, DMSO- d_6) δ 160.2, 158.5, 156.6,
6 138.8, 133.2, 129.5, 127.9, 120.6, 120.5, 116.6, 116.1.
7
8
9
10

11 **(iii) 4-(6-Cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)benzoic acid (22).** To a
12 solution of ethyl 4-(4-cyano-2-nitrophenylamino)benzoate (**21**) (891 mg, 2.86 mmol) in
13 EtOH (20 mL) was added SnCl₂ (5.55 g, 29.2 mmol). The orange mixture was refluxed for 45
14 min. It was then poured into EtOAc. The organic layer was washed with 1N HCl (50 mL),
15 brine, and satd. NaHCO₃, and dried over Na₂SO₄. After evaporation of the solvents under
16 reduced pressure the residue was purified by column chromatography on silica gel with 25%
17 EtOAc in hexanes to afford ethyl 4-(2-amino-4-cyanophenylamino)benzoate as a pale yellow
18 foam. Yield: 743 mg (92%). ^1H -NMR (300 MHz, CDCl₃) δ ppm 7.98-7.95 (m, 2H), 8.25 (d,
19 J = 6.0 Hz, 1 H), 7.11-7.06 (m, 2H), 6.89-6.85 (m, 2H), 5.71 (s, 1H), 4.38-4.32 (m, 2H),
20 4.00-3.70 (m, 2H), 1.40-1.36 (m, 3H). To a mixture of ethyl 4-(2-amino-4-
21 cyanophenylamino)benzoate (320 mg, 1.21 mmol) in dichloromethane (50 mL) under ice-
22 water bath cooling triphosgene (573 mg, 1.93 mmol) was added. The mixture was stirred at 0
23 °C for 1 h. Then it was allowed to warm to room temperature and stirred overnight. The
24 resulting mixture was then refluxed overnight. It was diluted with dichloromethane, washed
25 with satd. NaHCO₃, and brine, and evaporated to give a white solid ethyl 4-(6-cyano-1,2-
26 dihydro-2-oxobenzo[d]imidazol-3-yl)benzoate. Yield: 254 mg (72%). ^1H -NMR (300 MHz,
27 CDCl₃) δ ppm 9.55 (s, 1H), 8.29-8.26 (m, 2H), 7.66-7.30 (m, 2H), 7.45-7.42 (m, 2H), 7.18-
28 7.15 (m, 1H), 4.45 (q, J = 5.4 Hz, 2H), 1.44 (t, J = 5.4 Hz, 3H). To a mixture of ethyl 4-(6-
29 cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)benzoate (400 mg, 1.30 mmol) in THF/H₂O
30 (4:1 v/v, 15 mL) was added LiOH monohydrate (133 mg, 3.16 mmol). The mixture was
31 stirred overnight at room temperature and was then acidified with 1N HCl. The resulting
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

precipitate was collected by filtration to give 4-(6-cyano-1,2-dihydro-2-oxobenzodimidazol-3-yl)benzoic acid (**22**) as a greasy solid. Yield: 340 mg (94%). ¹H-NMR (300 MHz, CDCl₃) δ ppm 8.09 (d, *J* = 6.6 Hz, 2H), 7.67 (d, *J* = 6.6 Hz, 2H), 7.50-7.44 (m, 2H), 7.19 (d, *J* = 6.0 Hz, 1H), 7.08-7.02 and 6.82-6.76 (m, 1H).

(iv) **N'-((Z)-(2-Chlorophenylimino)(pyrimidin-4-yl)methyl)-4-(6-cyano-1,2-dihydro-2-oxobenzodimidazol-3-yl)benzohydrazide (23)**. To a mixture of 4-(6-cyano-1,2-dihydro-2-oxobenzodimidazol-3-yl)benzoic acid (**22**) (271 mg, 0.97 mmol) in THF (35 mL) was sequentially added (Z)-N''-(2-chlorophenyl)pyrimidine-4-carboximidhydrazide (**20**) (331 mg, 1.34 mmol), EDCI•HCl (323 mg, 1.68 mmol), HOBT (175 mg, 1.30 mmol) and Et₃N (1.5 mL, 10.8 mmol). The mixture was stirred for 3 d. It was diluted with dichloromethane, washed with water, satd. NaHCO₃, dried over Na₂SO₄. After filtration and evaporation of the solvents under reduced pressure the residue was purified by column chromatography on silica gel with 5% MeOH in dichloromethane as an eluent to afford N'-((Z)-(2-chlorophenylimino)(pyrimidin-4-yl)methyl)-4-(6-cyano-1,2-dihydro-2-oxobenzodimidazol-3-yl)benzohydrazide (**23**) as a pale yellow foam. Yield: 40 mg (8%). ¹H-NMR (300 MHz, DMSO-*d*₆) δ ppm 11.65 (s, 1H), 9.04 (s, 1H), 8.89-8.87 (m, 1H), 8.01-7.92 (m, 1H), 7.91-7.82 (m, 2 H), 7.64-7.58 (m, 3H), 7.50-7.35 (m, 4H), 7.18-7.02 (m, 2H), 7.00-6.92 (m, 1H), 6.75-6.68 (m, 1H).

(v) **1-(4-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-yl)phenyl)-2,3-dihydro-2-oxo-1H-benzodimidazole-5-carbonitrile (14)**. A mixture of N'-((Z)-(2-chlorophenylimino)(pyrimidin-4-yl)methyl)-4-(6-cyano-1,2-dihydro-2-oxobenzodimidazol-3-yl)benzohydrazide (**23**) (20 mg, 0.039 mmol) in toluene (10 mL) was refluxed for 3 d. The mixture was evaporated and subjected to column chromatography on silica gel with 2% MeOH in dichloromethane to give the crude product, which was washed with Et₂O to afford the title compound 1-(4-(4-(2-chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-

1
2
3 yl)phenyl)-2,3-dihydro-2-oxo-1H-benzo[d]imidazole-5-carbonitrile (**14**) as a pale yellow
4
5 solid. Yield: 6 mg (31%); mp 182 °C. MS for C₂₆H₁₅ClN₈O -H⁺, [M-H]⁻ m/z, calcd 489.1,
6
7 found 489.3. ¹H NMR (300 Hz, CDCl₃+CD₃OD) δ 8.82-8.00 (m, 2H), 8.28-8.26 (m, 1H),
8
9 7.66-7.60 (m, 2H), 7.54-7.44 (m, 4H), 7.40-7.32 (m, 4H), 7.08-7.06 (m, 1H).

11
12 **Preparation of 1-((trans)-4-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-**
13
14 **yl)cyclohexyl)-1H-benzo[d]imidazol-2(3H)-one (15a).**

15
16 (i) **N-(2-Chlorophenyl)pyrimidine-4-carbothioamide (25).** To a solution of N-(2-
17
18 chlorophenyl)pyrimidine-4-carboxamide (**24**) (2.0 g, 8.55 mmol) in toluene (20 ml)
19
20 Lawesson's reagent (2.42 g, 5.99 mmol) was added and the mixture was refluxed for 7 hours.
21
22 After the reaction mixture was cooled to room temperature, the solvents were evaporated.
23
24 The crude product was purified by chromatography on silica gel eluting with a gradient of
25
26 cyclohexane /ethyl acetate. The fractions containing the product were combined and the
27
28 solvent evaporated under reduced pressure to yield N-(2-chlorophenyl)pyrimidine-4-
29
30 carbothioamide (**25**) as an amorphous solid. Yield: 1.34 g (63%). HRMS for C₁₁H₈ClN₃S
31
32 +H⁺, [M+H]⁺ m/z, calcd 250.0200, found 250.0206. ¹H NMR (300 MHz, CDCl₃) δ 12.39 (s,
33
34 1H), 9.32 (s, 1H), 9.09 (d, *J* = 8.2 Hz, 1H), 9.02 (d, *J* = 5.2 Hz, 1H), 8.65 (d, *J* = 5.2 Hz, 1H),
35
36 7.54 (dd, *J* = 8.0, 1.5 Hz, 1H), 7.41 (t, *J* = 7.8 Hz, 1H), 7.27 (td, *J* = 7.7, 1.5 Hz, 1H). ¹³C
37
38 NMR (75 MHz, CDCl₃) δ 185.5, 158.6, 157.0, 156.5, 134.9, 129.7, 127.5, 127.1, 126.7,
39
40 123.4, 120.3.

41
42
43 (ii) **Methyl(Z)-N-(2-chlorophenyl)pyrimidine-4-carbimidothioate (26).** To a flask
44
45 charged with N-(2-chlorophenyl)pyrimidine-4-carbothioamide (**25**) (249.7 mg, 1.0 mmol) in
46
47 tetrahydrofuran (6 ml) was added potassium tert-butoxide (112.2 mg, 1.0 mmol) and stirred
48
49 for 15 min. Methyl tosylate (151 μl, 1.0 mmol) was then added dropwise and the reaction
50
51 mixture was stirred at room temperature for further 18 hours. The reaction mixture was then
52
53 partitioned between water and ethyl acetate, the layers were separated and the organic layer
54
55
56
57
58
59
60

1
2
3 was washed twice with water and brine, dried over magnesium sulfate, filtered and
4
5 evaporated under reduced pressure to provide crude product. The crude greasy product
6
7 methyl(*Z*)-*N*-(2-chlorophenyl)pyrimidine-4-carbimidothioate (**26**) was directly used in the
8
9 next reaction step without any purification. Yield: 248 mg (94%). HRMS for C₁₂H₁₀ClN₃S
10
11 +H⁺, [M+H]⁺ m/z, calcd 264.0357, found 264.0369.

12
13
14 **(iii) (*trans*)-Methyl-4-(2-nitrophenylamino)cyclohexane carboxylate (29b).** To a
15
16 solution of (*trans*)-methyl-4-aminocyclohexane (**27a**) carboxylate hydrochloride (1 g, 5.16
17
18 mmol) and 1-fluoro-2-nitrobenzene (**28b**) (544 μ l, 5.16 mmol) in acetonitrile (50 ml)
19
20 potassium carbonate (1.06 g, 77.4 mmol) was added at ambient temperature. The resulting
21
22 reaction mixture was heated for 8 hours at reflux. After cooling to room temperature, the
23
24 acetonitrile was removed under reduced pressure and the crude solid was dissolved in
25
26 dichloromethane. The dichloromethane layer was washed with water, dried over magnesium
27
28 sulfate, filtered and then concentrated under reduced pressure. The crude solid product was
29
30 washed with cold methanol to yield (*trans*)-methyl-4-(2-nitrophenylamino)-cyclohexane
31
32 carboxylate (**29b**) as a yellow solid which was used in the next reaction step without any
33
34 further purification. Yield: 1.3 g (91%); mp 81 - 83 °C. HRMS for C₁₄H₁₈N₂O₄ +H⁺, [M+H]⁺
35
36 m/z, calcd 279.1339, found 279.1350. ¹H NMR (600 MHz, DMSO-*d*₆) δ 8.08 – 8.00 (m, 1H),
37
38 7.89 (d, *J* = 8.0 Hz, 1H), 7.51 (q, *J* = 9.4, 8.8 Hz, 1H), 7.14 (t, *J* = 9.2 Hz, 1H), 6.67 (q, *J* =
39
40 10.4, 9.1 Hz, 1H), 3.67 – 3.53 (m, 4H), 2.40 – 2.31 (m, 1H), 2.09 – 2.03 (m, 2H), 1.95 (d, *J* =
41
42 14.4 Hz, 2H), 1.61 – 1.51 (m, 2H), 1.38 (q, *J* = 15.4, 13.0 Hz, 2H). ¹³C NMR (151 MHz,
43
44 DMSO-*d*₆) δ 175.1, 144.3, 136.6, 130.8, 126.3, 115.3, 114.9, 51.3, 49.8, 41.3, 31.0, 27.2.

45
46
47 **(iv) (*trans*)-Methyl-4-(1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexane**
48
49 **carboxylate (30b).** A reaction flask with a mixture of (*trans*)-methyl-4-(2-
50
51 nitrophenylamino)-cyclohexane carboxylate (**29b**) (1.85 g, 9.55 mmol) and a catalytic
52
53 amount of 10% palladium on charcoal in ethanol (400 ml) was equipped with a hydrogen
54
55
56
57
58
59
60

balloon and hydrogen gas was bubbled into reaction mixture at atmospheric pressure. After 2 hours, the balloon was removed and the reaction mixture purged with nitrogen, filtered through a pad of celite and washed twice with ethanol. The filtrate was concentrated under reduced pressure to afford (*trans*)-methyl-4-(2-aminophenylamino)cyclohexane carboxylate as a yellow greasy solid. Yield: 1.25 g (76%). HRMS for $C_{14}H_{20}N_2O_2 + H^+$, $[M+H]^+$ m/z , calcd 249.1598, found 249.1599. 1H NMR (600 MHz, DMSO- d_6) δ 6.58 (d, $J = 7.7$ Hz, 1H), 6.51 (t, $J = 4.1$ Hz, 2H), 6.42 (td, $J = 7.9, 3.8$ Hz, 1H), 4.88 (s, 3H), 3.60 (t, $J = 4.4$ Hz, 3H), 3.17 (q, $J = 7.8, 4.7$ Hz, 1H), 2.31 (tt, $J = 15.4, 4.8$ Hz, 1H), 2.07 – 2.00 (m, 2H), 1.97 – 1.91 (m, 2H), 1.47 (qd, $J = 13.3, 3.9$ Hz, 2H), 1.21 (dq, $J = 17.8, 7.2, 4.6$ Hz, 2H). ^{13}C NMR (151 MHz, DMSO- d_6) δ 175.3, 134.7, 134.3, 118.1, 117.2, 115.1, 111.5, 51.3, 50.8, 42.0, 31.5, 27.6. To a mixture of (*trans*)-methyl-4-(2-aminophenylamino)cyclohexane carboxylate (1.00 g, 4.02 mmol) in dichloromethane (70 ml) at 0°C (ice bath) was added triphosgene (1.79 g, 6.03 mmol). The mixture was stirred at 0°C for 1 h. Then it was allowed to warm to room temperature and stirred for 24 hours. Next the resulting mixture was refluxed for 24 hours. The reaction mixture was diluted with dichloromethane and washed with saturated sodium bicarbonate, and brine, dried over magnesium sulfate filtered and evaporated to give (*trans*)-methyl-4-(1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexane carboxylate (**30b**) as an amorphous solid. Yield: 950 mg (86%). HRMS for $C_{15}H_{18}N_2O_3 + H^+$, $[M+H]^+$ m/z , calcd 275.1390, found 275.1403. 1H NMR (600 MHz, DMSO- d_6) δ 10.80 (d, $J = 5.1$ Hz, 1H), 7.30 (t, $J = 6.6$ Hz, 1H), 6.95 (d, $J = 5.6$ Hz, 3H), 4.19 – 4.11 (m, 1H), 3.64 – 3.58 (m, 3H), 2.21 (q, $J = 12.8$ Hz, 2H), 2.04 (d, $J = 13.2$ Hz, 2H), 1.72 (d, $J = 12.5$ Hz, 2H), 1.53 (q, $J = 12.9$ Hz, 2H). ^{13}C NMR (151 MHz, DMSO- d_6) δ 175.1, 153.7, 129.2, 128.3, 120.4, 120.2, 108.9, 108.7, 51.4, 50.7, 41.0, 28.1.

(v) (*trans*)-4-(1,2-Dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexanecarbohydrazide (**31b**). To a mixture of (*trans*)-methyl-4-(1,2-dihydro-2-oxobenzo[d]imidazol-3-

1
2
3 yl)cyclohexane carboxylate (**30b**) (500 mg, 1.82 mmol) in ethanol was added hydrazine
4 hydrate and heated at 120°C for 3 hours under microwave irradiation in a sealed vial (Biotage
5 initiator⁺). Upon completion of reaction, the solvent was removed under reduced pressure and
6 the crude solid was washed with a mixture of dichloromethane and methanol to yield (*trans*)-
7 4-(1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexanecarbohydrazide (**31b**) as a solid
8 powder which was used in the next reaction step without further purification. Yield: 460 mg
9 (92%), mp 273 °C (decomposition). HRMS for C₁₄H₁₈N₄O₂ +H⁺, [M+H]⁺ m/z, calcd
10 275.1503, found 275.1511.
11
12
13
14
15
16
17
18

19
20 (vi) 1-((*trans*)-4-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-
21 yl)cyclohexyl)-1H-benzo[d]imidazol-2(3H)-one (**15a**). The (*trans*)-4-(1,2-dihydro-2-
22 oxobenzo[d]imidazol-3-yl)cyclohexanecarbohydrazide (**31b**) (200 mg, 0.729 mmol) was
23 added to a solution of methyl(*Z*)-N-(2-chlorophenyl)pyrimidine-4-carbimidothioate (**26**) (211
24 mg, 0.801 mmol) in N,N-dimethylacetamide (1 ml). Trifluoroacetic acid (27.8 µl) was added
25 and the reaction mixture was heated at 120°C for 14 hours. After cooling down to room
26 temperature, the reaction mixture was filtered and the residue washed with dichloromethane
27 and methanol. The filtrate was concentrated under reduced pressure to remove the methanol
28 and dichloromethane. A mixture of water and dichloromethane was added and the reaction
29 mixture portioned using a separating funnel. The organic layer was washed with water, brine,
30 dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude
31 product was purified by chromatography on silica gel eluting with a gradient of
32 dichloromethane / methanol. The fractions containing the product were combined and the
33 solvent evaporated under reduced pressure to yield the title compound **15a** as a white solid.
34 Yield: 72 mg (21%); mp 274 - 276 °C. HRMS for C₂₅H₂₂ClN₇O +H⁺, [M+H]⁺ m/z, calcd
35 472.1647, found 472.1654. ¹H NMR (600 MHz, DMSO-*d*₆) δ 10.87 (s, 1H), 9.00 (d, *J* = 5.3
36 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
37 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
38 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
39 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
40 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
41 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
42 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
43 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
44 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
45 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
46 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
47 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
48 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
49 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
50 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
51 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
52 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
53 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
54 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
55 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
56 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
57 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
58 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
59 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,
60 Hz, 1H), 8.93 (d, *J* = 1.4 Hz, 1H), 8.27 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.86 (dd, *J* = 7.8, 1.7 Hz,

1
2
3 1H), 7.81 (dd, $J = 8.0, 1.4$ Hz, 1H), 7.72 (td, $J = 7.8, 1.7$ Hz, 1H), 7.67 (td, $J = 7.6, 1.5$ Hz,
4 1H), 7.35 (td, $J = 4.7, 4.1, 2.2$ Hz, 1H), 7.01 (d, $J = 2.7$ Hz, 3H), 4.29 (tt, $J = 12.2, 3.9$ Hz,
5 1H), 2.62 (tdd, $J = 11.9, 7.3, 3.7$ Hz, 1H), 2.19 (dtd, $J = 30.7, 14.8, 13.9, 3.5$ Hz, 3H), 2.02 –
6
7 1.90 (m, 3H), 1.81 (ddd, $J = 15.6, 8.8, 2.9$ Hz, 2H). ^{13}C NMR (151 MHz, DMSO- d_6) δ 160.1,
8
9 158.3, 158.0, 153.7, 153.2, 150.0, 132.5, 131.6, 131.0, 130.1, 129.8, 129.4, 128.5, 128.2,
10
11 120.4, 120.2, 118.9, 108.7, 108.6, 50.8, 32.7, 30.6, 30.1, 28.5.

12
13
14
15
16 **Preparation of 1-((*trans*)-4-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-**
17
18 **3-yl)cyclohexyl)-2,3-dihydro-2-oxo-1H-benzo[d]imidazole-5-carbonitrile (15)**

19
20 (i) (*trans*)-Methyl-4-(4-cyano-2-nitrophenylamino)cyclohexane carboxylate (29a). To
21 a solution of methyl-4-aminocyclohexane carboxylate hydrochloride (27a) (1.5 g, 7.74
22 mmol) and 4-fluoro-3-nitrobenzonitrile (28a) (1.28 g, 7.74 mmol) in acetonitrile (50 ml) was
23 added N,N-diisopropylethylamine (2.69 ml, 15.49 mmol) at room temperature. The resulting
24 reaction mixture was stirred for 18 hours at room temperature. Acetonitrile was removed
25 under reduced pressure and the crude solid was dissolved in dichloromethane and the
26 dichloromethane layer was washed with water, dried over magnesium sulfate, filtered and
27 then concentrated under reduced pressure. The crude solid (*trans*)-methyl-4-(4-cyano-2-
28 nitrophenylamino)cyclohexane carboxylate (29a) was washed with cold methanol and used in
29 the next step without any further purification. Yield 1.8 g (77%); mp 138 - 140 °C. HRMS
30 for $\text{C}_{15}\text{H}_{17}\text{N}_3\text{O}_4 + \text{H}^+$, $[\text{M}+\text{H}]^+$ m/z , calcd = 304.1292, found 304.1307. ^1H NMR (600 MHz,
31 DMSO- d_6) δ 8.49 (d, $J = 2.1$ Hz, 1H), 8.20 (d, $J = 7.9$ Hz, 1H), 7.81 (dd, $J = 9.1, 2.1$ Hz, 1H),
32 7.30 (d, $J = 9.2$ Hz, 1H), 3.72 (tdt, $J = 11.3, 7.8, 3.9$ Hz, 1H), 3.61 (s, 3H), 2.35 (tt, $J = 11.9,$
33 3.6 Hz, 1H), 2.05 – 1.99 (m, 2H), 1.98 – 1.92 (m, 2H), 1.56 (qd, $J = 13.0, 3.1$ Hz, 2H), 1.49 –
34 1.41 (m, 2H). ^{13}C NMR (151 MHz, DMSO- d_6) δ 175.0, 146.0, 137.7, 132.0, 130.6, 118.2,
35 116.1, 96.5, 51.4, 50.3, 41.2, 30.6, 27.1.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 (ii) **(*trans*)-Methyl-4-(6-cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexane**
4
5 **carboxylate (30a)**. A reaction flask with a mixture of (*trans*)-methyl-4-(4-cyano-2-
6 nitrophenylamino) cyclohexane carboxylate (**29a**) (2.38 g, 7.85 mmol), catalytic amount of
7 10% palladium on charcoal in ethanol (450 ml) was equipped with a hydrogen balloon and
8 hydrogen gas bubbled through the reaction mixture at atmospheric pressure. After 2 hours,
9 the balloon was removed and the reaction mixture purged with nitrogen, filtered through a
10 pad of celite and washed twice with ethanol. Then the filtrate was concentrated under reduced
11 pressure to afford (*trans*)-methyl-4-(2-amino-4-cyanophenylamino)cyclohexane carboxylate
12 as an amorphous solid. Yield: 1.84 g (86%). HRMS for $C_{15}H_{19}N_3O_2 + H^+$, $[M+H]^+$ m/z, calcd
13 274.1550, found 274.1553. 1H NMR (600 MHz, DMSO- d_6) δ 6.89 (d, $J = 8.8$ Hz, 1H), 6.75
14 (d, $J = 9.7$ Hz, 1H), 6.51 (t, $J = 9.8$ Hz, 1H), 5.09 (d, $J = 8.5$ Hz, 1H), 4.97 (s, 2H), 3.59 (q, J
15 = 12.0 Hz, 3H), 2.32 (q, $J = 11.2, 10.0$ Hz, 1H), 2.04 – 1.97 (m, 2H), 1.93 (t, $J = 11.3$ Hz,
16 2H), 1.49 (dt, $J = 22.1, 12.1$ Hz, 2H), 1.23 (dt, $J = 22.3, 12.1$ Hz, 2H). ^{13}C NMR (151 MHz,
17 DMSO- d_6) δ 175.2, 138.8, 135.0, 122.7, 121.1, 115.1, 108.9, 96.1, 51.3, 50.0, 41.8, 31.3,
18 27.5. To a mixture of (*trans*)-methyl-4-(2-amino-4-cyanophenylamino)cyclohexane
19 carboxylate (1.87 g, 6.84 mmol) in dichloromethane (65 ml) was added triphosgene (3.06 g,
20 10.31 mmol) at 0°C (ice water bath). The mixture was stirred at 0 °C for 1 h. Then it was
21 allowed to warm to room temperature and stirred for 24 hours. Next the resulting mixture was
22 heated to reflux for 24 hours. The reaction mixture was diluted with dichloromethane and
23 washed with saturated sodium bicarbonate, and brine, dried over magnesium sulfate and
24 evaporated to give (*trans*)-methyl-4-(6-cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-
25 yl)cyclohexane carboxylate (**30a**) as a glassy solid. Yield: 1.90 g (93%). HRMS for
26 $C_{16}H_{17}N_3O_3 + H^+$, $[M+H]^+$ m/z, calcd 300.1343, found 300.1356. 1H NMR (600 MHz,
27 DMSO- d_6) δ 11.30 (s, 1H), 7.53 (d, $J = 8.3$ Hz, 1H), 7.44 (dd, $J = 8.3, 1.6$ Hz, 1H), 7.35 (d, J
28 = 1.6 Hz, 1H), 4.20 (tt, $J = 12.4, 4.0$ Hz, 1H), 3.62 (s, 3H), 2.20 (qd, $J = 13.0, 3.7$ Hz, 2H),
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 2.03 (ddd, $J = 11.9, 5.0, 2.8$ Hz, 2H), 1.74 (dt, $J = 13.4, 3.7$ Hz, 2H), 1.53 (qd, $J = 13.2, 3.5$
4 Hz, 2H). ^{13}C NMR (151 MHz, DMSO- d_6) δ 175.1, 153.6, 133.0, 128.6, 125.5, 119.7, 111.5,
5 109.5, 102.3, 51.4, 51.2, 40.9, 28.0, 27.9.
6
7

8
9 (iii) **(trans)-4-(6-Cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexanecarbohydrazide (31a)**. To a mixture of (trans)-methyl-4-(6-cyano-1,2-
10 dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexane carboxylate (**30a**) (299.3 mg, 1.00 mmol)
11 in ethanol (7 ml) was added hydrazine hydrate (7 ml) and heated to 80 °C for 3 hours under
12 microwave irradiation in a sealed vial (Biotage initiator+). Upon completion of reaction, the
13 reaction mixture was filtered and the crude solid was washed with methanol, filtered and
14 dried to afford (trans)-3-(1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclobutanecarbohydrazide (**31a**) as a white solid. Yield: 269 mg (90%); mp 348 °C
15 (decomposition). HRMS for $\text{C}_{15}\text{H}_{17}\text{N}_5\text{O}_2 + \text{H}^+$, $[\text{M}+\text{H}]^+$ m/z, calcd 300.1455, found 300.1466.
16
17
18
19
20
21
22
23
24
25
26
27
28

29 (iv) **1-((trans)-4-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-yl)cyclohexyl)-2,3-dihydro-2-oxo-1H-benzo[d]imidazole-5-carbonitrile (15)**. (trans)-4-(6-
30 Cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclohexanecarbohydrazide (**31a**) (142 mg,
31 0.47 mmol) was added to a solution of methyl(Z)-N-(2-chlorophenyl)pyrimidine-4-
32 carbimidothioate (**26**) (138 mg, 0.52 mmol) in N,N-dimethylacetamide (2 ml). Trifluoroacetic
33 acid (18.2 μl , 0.23 mmol) was added and the reaction mixture was heated at 120 °C for 14
34 hours. The reaction mixture was then cooled down to room temperature and filtered. Water
35 was added to the filtrate and the reaction mixture was extracted with dichloromethane. The
36 combined organic layers were washed with water, brine, dried over magnesium sulfate,
37 filtered and concentrated under reduced pressure. The crude solid was then washed with an
38 acetonitrile/water mixture (1:1, 3 ml) followed by cold methanol to afford compound 1-
39 ((trans)-4-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-yl)cyclohexyl)-2,3-
40 dihydro-2-oxo-1H-benzo[d]imidazole-5-carbonitrile (**15**) as a light yellow solid. Yield: 24
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 mg (10%); mp 267 °C. HRMS for C₂₆H₂₁ClN₈O, +H⁺, [M+H]⁺ m/z, calcd = 497.1600, found
4 497.1609. ¹H NMR (600 MHz, DMSO-*d*₆) δ 11.44 (s, 1H), 9.08 (d, *J* = 5.3 Hz, 1H), 9.01 (d,
5 *J* = 1.4 Hz, 1H), 8.34 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.93 (dd, *J* = 7.8, 1.7 Hz, 1H), 7.89 (dd, *J* =
6 8.1, 1.4 Hz, 1H), 7.80 (td, *J* = 7.8, 1.7 Hz, 1H), 7.75 (td, *J* = 7.7, 1.5 Hz, 1H), 7.67 (d, *J* = 8.3
7 Hz, 1H), 7.58 (dd, *J* = 8.2, 1.6 Hz, 1H), 7.48 (d, *J* = 1.5 Hz, 1H), 4.43 (tt, *J* = 12.3, 3.9 Hz,
8 1H), 2.73 – 2.66 (m, 1H), 2.31 – 2.19 (m, 3H), 2.10 – 1.98 (m, 3H), 1.95 – 1.85 (m, 2H). ¹³C
9 NMR (151 MHz, DMSO-*d*₆) δ 160.0, 158.3, 158.0, 153.6, 153.2, 150.0, 133.2, 132.5, 131.6,
10 131.0, 130.1, 129.8, 128.5, 128.4, 125.5, 119.7, 118.9, 111.4, 109.4, 102.3, 51.3, 32.6, 30.5,
11 30.0, 28.2.

12
13
14
15
16
17
18
19
20
21
22 **Preparation of 1-((*trans*)-3-(4-(2-chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-**
23 **yl)cyclobutyl)-1H-benzo[d]imidazol-2(3H)-one (16a).**

24
25
26
27 (i) (*trans*)-Methyl-3-(2-nitrophenylamino)cyclobutanecarboxylate (29d). To a
28 solution of (*trans*)-methyl-3-aminocyclobutanecarboxylate hydrochloride (27b) (1.0 g, 6.05
29 mmol) and 1-fluoro-2-nitrobenzene (28b) (1.09 g, 7.74 mmol) in acetonitrile (50 ml) was
30 added potassium carbonate (1.60 g, 11.6 mmol) at ambient temperature. The resulting
31 reaction mixture was heated for 16 hours at reflux. Acetonitrile was removed under reduced
32 pressure and the crude solid was dissolved in dichloromethane and the dichloromethane layer
33 was washed with water, dried over magnesium sulfate and then concentrated under reduced
34 pressure. Unreacted 1-fluoro-2-nitrobenzene was removed by co-distillation with toluene.
35 The crude product was purified by chromatography on silica gel eluting with a gradient of
36 cyclohexane /ethyl acetate. The fractions containing the product were combined and the
37 solvent evaporated under reduced pressure to yield 1.43g of (*trans*)-methyl-(2-
38 nitrophenylamino)cyclobutanecarboxylate (29d) as a greasy solid. Yield: 1.43 g (95%).
39 HRMS for C₁₂H₁₄N₂O₄ +H⁺, [M+H]⁺ m/z, calcd 251.1026, found 251.1044. ¹H NMR (300
40 MHz, DMSO-*d*₆) δ 8.10 – 8.04 (m, 1H), 8.01 (d, *J* = 5.8 Hz, 1H), 7.53 (ddd, *J* = 8.7, 6.9, 1.9
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Hz, 1H), 6.81 (d, $J = 8.7$ Hz, 1H), 6.79 – 6.67 (m, 1H), 4.24 (h, $J = 7.1$ Hz, 1H), 3.66 (s, 3H),
4
5 3.18 (tt, $J = 9.6, 4.4$ Hz, 1H), 2.75 – 2.60 (m, 2H), 2.34 (dtd, $J = 12.9, 6.9, 2.5$ Hz, 2H). ^{13}C
6
7 NMR (75 MHz, DMSO- d_6) δ 175.3, 143.7, 136.7, 131.4, 126.2, 115.9, 114.9, 51.7, 45.5,
8
9 32.3, 32.2.

10
11 (ii) **(*trans*)-Methyl-3-(1,2-dihydro-2-oxobenzo[d]imidazol-3-**
12 **yl)cyclobutanecarboxylate (30d)**. A reaction flask with a mixture of (*trans*)-methyl-3-(2-
13 nitrophenylamino)cyclobutane carboxylate (**29d**) (1.4 g, 5.59 mmol), a catalytic amount of
14 10% palladium on charcoal in ethanol was equipped with a hydrogen balloon and hydrogen
15 gas was bubbled into reaction mixture at atmospheric pressure. After 2 hours, the balloon was
16 removed and the reaction mixture purged with nitrogen, filtered through a pad of celite and
17 washed twice with ethanol. The filtrate was concentrated under reduced pressure to afford
18 (*trans*)-methyl-3-(2-aminophenylamino)cyclobutanecarboxylate as a solid. The crude product
19 was used in next step without any further purification. Yield: 936 mg. (76%). HRMS for
20 $\text{C}_{12}\text{H}_{16}\text{N}_2\text{O}_2 + \text{H}^+$, $[\text{M}+\text{H}]^+$ m/z , calcd 221.1285, found 221.1286. ^1H NMR (600 MHz,
21 DMSO- d_6) δ 6.53 (d, $J = 6.8$ Hz, 1H), 6.44 (dt, $J = 19.4, 7.2$ Hz, 2H), 6.21 (d, $J = 7.5$ Hz,
22 1H), 4.71 (d, $J = 6.5$ Hz, 1H), 4.50 (s, 2H), 3.93 (q, $J = 6.9$ Hz, 1H), 3.65 (s, 3H), 3.14 (dt, $J =$
23 9.8, 5.1 Hz, 1H), 2.56 (ddd, $J = 12.5, 7.4, 4.3$ Hz, 2H), 2.15 (ddd, $J = 12.6, 9.5, 6.0$ Hz, 2H).
24 ^{13}C NMR (151 MHz, DMSO- d_6) δ 175.7, 135.5, 134.3, 117.4, 114.0, 110.5, 51.5, 46.3, 32.6,
25 32.5. To a mixture of (*trans*)-methyl-3-(2-aminophenylamino)cyclobutanecarboxylate (1.12
26 g, 5.08 mmol) in dichloromethane was added triphosgene (2.26 g, 7.63 mmol) at 0°C (ice
27 water bath). The mixture was stirred at 0 °C for 1 h. Then it was allowed to warm to room
28 temperature and stirred for 24 hours. Next the resulting mixture was refluxed for 24 hours.
29 The reaction mixture was diluted with dichloromethane and washed with saturated sodium
30 bicarbonate, and brine, dried over magnesium sulfate and evaporated under reduced pressure
31 to give (*trans*)-methyl-3-(1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclobutanecarboxylate
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **(30d)** as a white foam. Yield: 1.15 g (92%). HRMS for $C_{13}H_{14}N_2O_3 + H^+$, $[M+H]^+$ m/z, calcd
4 247.1077, found 247.1076. 1H NMR (300 MHz, $CDCl_3$) δ 10.60 (s, 1H), 7.22 – 7.04 (m, 4H),
5 5.36 – 5.03 (m, 1H), 3.80 (s, 3H), 3.33 (q, $J = 7.5$ Hz, 3H), 2.82 – 2.65 (m, 2H). ^{13}C NMR
6 (75 MHz, $CDCl_3$) δ 176.0, 155.5, 129.3, 128.0, 121.4, 121.0, 109.9, 108.3, 52.0, 45.3, 32.3,
7
8
9
10
11
12 30.3.

13
14 **(iii) (*trans*)-3-(1,2-Dihydro-2-oxobenzo[d]imidazol-3-yl)cyclobutanecarbohydrazide**

15
16 **(31d)**. To a mixture of (*trans*)-methyl-3-(1,2-dihydro-2-oxobenzo[d]imidazol-3-
17 yl)cyclobutanecarboxylate **(30d)** (400 mg, 1.62 mmol) in ethanol (7 ml) was added hydrazine
18 hydrate (7 ml) and heated to 100 °C for 3 hours under microwave irradiation in a sealed vial
19 (Biotage initiator⁺). Upon completion of reaction, the solvent was removed under reduced
20 pressure and the crude solid was washed with methanol resulted in (*trans*)-3-(1,2-dihydro-2-
21 oxobenzo[d]imidazol-3-yl)cyclobutanecarbohydrazide **(31d)** as a solid product which was
22 used in the next step without any further purification. Yield: 360 mg. (90%); mp > 250 °C
23 (decomposition). HRMS for $C_{12}H_{14}N_4O_2 + H^+$, $[M+H]^+$ m/z, calcd 247.1190, found 247.1202.
24
25
26
27
28
29
30
31
32

33 **(iv) 1-((*trans*)-3-(4-(2-chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-
34 yl)cyclobutyl)-1H-benzo[d]imidazol-2(3H)-one (16a).** (*trans*)-3-(1,2-Dihydro-2-
35 oxobenzo[d]imidazol-3-yl)cyclobutanecarbohydrazide **(31d)** (123 mg, 0.5 mmol) was added
36 to a solution of methyl(*Z*)-N-(2-chlorophenyl)pyrimidine-4-carbimidothioate **(26)** (158 mg,
37 0.6 mmol) in N,N-dimethylacetamide (1 ml). Trifluoroacetic acid (19 μ l, 0.25 mmol) was
38 added and the reaction mixture was heated to 120°C for 14 hours. Water was added and the
39 reaction mixture was extracted with dichloromethane. The organic layer was washed with
40 water, brine, dried over magnesium sulfate, filtered and concentrated. The crude solid was
41 purified by preparative HPLC (C18 reverse phase column, elution with a water/MeCN
42 gradient with 0.1% TFA). The fractions containing the product were evaporated and
43 lyophilized to yield 1-((*trans*)-3-(4-(2-chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 yl)cyclobutyl)-1H-benzo[d]imidazol-2(3H)-one (**16a**) as a white solid. The product was
4
5 obtained as its trifluoroacetate salt. Yield: 46 mg (16%); mp 160 °C. HRMS for C₂₃H₁₈ClN₇O
6
7 +H⁺, [M+H]⁺ m/z, calcd 444.1334, found 444.1342. ¹H NMR (600 MHz, CDCl₃) δ 9.41 (s,
8
9 1H), 8.87 – 8.84 (m, 2H), 8.33 (dd, *J* = 5.3, 1.4 Hz, 1H), 7.58 (dd, *J* = 8.1, 1.5 Hz, 1H), 7.53
10
11 (td, *J* = 7.8, 1.6 Hz, 1H), 7.47 (td, *J* = 7.7, 1.6 Hz, 1H), 7.34 (dd, *J* = 7.9, 1.6 Hz, 1H), 7.17 (d,
12
13 *J* = 7.8 Hz, 1H), 7.13 (dt, *J* = 7.8, 4.2 Hz, 1H), 7.10 (dd, *J* = 4.0, 0.9 Hz, 2H), 5.40 – 5.24 (m,
14
15 1H), 3.50 – 3.41 (m, 2H), 3.41 – 3.34 (m, 1H), 3.01 – 2.96 (m, 1H), 2.95 – 2.90 (m, 1H). ¹³C
16
17 NMR (151 MHz, DMSO-*d*₆) δ 159.6, 158.4, 158.1, 153.7, 153.1, 150.7, 132.6, 131.5, 130.8,
18
19 130.0, 129.7, 129.1, 128.5, 128.2, 120.8, 120.5, 118.9, 108.8, 108.6, 44.0, 31.2, 30.6, 23.6.

22
23 **Preparation of 1-((*trans*)-3-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-**
24
25 **yl)cyclobutyl)-2,3-dihydro-2-oxo-1H-benzo[d]imidazole-5-carbonitrile (**16**).**

26
27 (i) (*trans*)-Methyl-3-(4-cyano-2-nitrophenylamino)cyclobutanecarboxylate (**29c**). To
28
29 a solution of (*trans*)-methyl-3-aminocyclobutanecarboxylate hydrochloride (**27b**) (331.2 mg,
30
31 2.0 mmol) and 4-fluoro-3-nitrobenzotrile (**28a**) (332 mg, 2.0 mmol) in acetonitrile (4 ml)
32
33 was added N,N-diisopropylethylamine (1.04 ml, 6.0 mmol) at ambient temperature. The
34
35 resulting reaction mixture was stirred for 24 hours at room temperature. After completion of
36
37 the reaction, the acetonitrile was removed under reduced pressure and the crude solid was
38
39 dissolved in dichloromethane. The dichloromethane layer was washed twice with water,
40
41 brine, dried over magnesium sulfate, concentrated under reduced pressure and the
42
43 precipitated product washed with cold methanol to yield (*trans*)-methyl-3-(4-cyano-2-
44
45 nitrophenylamino)cyclobutanecarboxylate (**29c**) as a light yellow solid. Yield: 318 mg
46
47 (96%); mp 127 - 131 °C. HRMS for C₁₃H₁₃N₃O₄ +H⁺, [M+H]⁺ m/z, calcd 276.0979, found
48
49 276.0992. ¹H NMR (300 MHz, CDCl₃) δ 8.53 – 8.42 (m, 2H), 7.61 (dd, *J* = 9.0, 1.9 Hz, 1H),
50
51 6.77 (d, *J* = 9.0 Hz, 1H), 4.38 (h, *J* = 7.3 Hz, 1H), 3.76 (s, 3H), 3.22 (td, *J* = 9.6, 4.7 Hz, 1H),
52
53
54
55
56
57
58
59
60

2.94 – 2.76 (m, 2H), 2.37 (dtd, $J = 13.1, 7.1, 2.7$ Hz, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ 175.4, 145.7, 137.7, 132.0, 131.4, 117.7, 115.2, 98.6, 52.1, 45.9, 33.0, 32.8.

(ii) **(*trans*)-Methyl-3-(6-cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclobutanecarboxylate (30c)**. A reaction flask charged with a mixture of (*trans*)-methyl-3-(4-cyano-2-nitrophenylamino)cyclobutanecarboxylate (**29c**) (1.5 g, 5.45 mmol), a catalytic amount of 10% palladium on charcoal in ethanol (400 ml) was equipped with a hydrogen balloon and hydrogen gas was bubbled through the reaction mixture at atmospheric pressure. After 2 hours, the balloon was removed and the reaction mixture was purged with nitrogen, filtered through a pad of celite and washed twice with ethanol. The filtrate was concentrated under reduced pressure to afford (*trans*)-methyl-3-(2-amino-4-cyanophenylamino)cyclobutanecarboxylate as a greasy solid. Yield: 962 mg (72%). MS (ESI) for $\text{C}_{13}\text{H}_{15}\text{N}_3\text{O}_2 + \text{H}^+$, $[\text{M}+\text{H}]^+$ m/z , calcd = 246.1, found 246.2. To a mixture of (*trans*)-methyl-3-(2-amino-4-cyanophenylamino)cyclobutanecarboxylate (1 g, 4.07 mmol) in dichloromethane (60 ml) at 0°C (ice water bath) was added triphosgene (1.8 g, 6.12 mmol). The mixture was stirred at 0°C for 1 h and was then allowed to warm to room temperature and stirred for 24 hours. Next the resulting mixture was refluxed for 24h. The reaction mixture was diluted with dichloromethane and washed with saturated sodium bicarbonate, and brine, dried over magnesium sulfate and evaporated under reduced pressure to give (*trans*)-methyl-3-(6-cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclobutanecarboxylate (**30c**) as an amorphous solid. Yield: 1.05 g (95%). MS (ESI) for $\text{C}_{14}\text{H}_{13}\text{N}_3\text{O}_3 + \text{H}^+$, $[\text{M}+\text{H}]^+$ m/z , calcd 272.1, found 272.2. ^1H NMR (300 MHz, CDCl_3) δ 10.62 (s, 1H), 7.43 (d, $J = 8.2$ Hz, 2H), 7.21 (d, $J = 8.2$ Hz, 1H), 5.21 – 5.02 (m, 1H), 3.80 (s, 3H), 3.30 (td, $J = 8.6, 8.0, 3.1$ Hz, 3H), 2.84 – 2.63 (m, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ 175.8, 155.4, 132.9, 127.9, 126.2, 119.2, 112.9, 108.7, 104.6, 52.2, 45.9, 32.3, 30.3.

1
2
3 (iii) **(trans)-3-(5-Cyano-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)cyclobutane-1-**
4 **carbohydrazide (31c)**. To a mixture of *(trans)*-methyl-3-(6-cyano-1,2-dihydro-2-
5 oxobenzo[d]imidazol-3-yl)cyclobutanecarboxylate (**30c**) (360.0 mg, 1.32 mmol) in methanol
6 (7 ml) was added hydrazine hydrate in ethanol (7 ml) and stirred at 20 °C for 20 hours. Upon
7 completion of reaction, the reaction mixture was filtered and the crude solid was washed with
8 methanol, filtered and dried to afford *(trans)*-3-(6-cyano-1,2-dihydro-2-oxobenzo[d]imidazol-
9 3-yl)cyclobutanecarbohydrazide (**31c**) as an amorphous powder. Yield: 230 mg (64%). MS
10 (ESI) for C₁₃H₁₃N₅O₂, +H⁺, [M+H]⁺ m/z, calcd 272.1, found 272.2.
11
12
13
14
15
16
17
18
19

20 (iv) **1-((trans)-3-(4-(2-Chlorophenyl)-5-(pyrimidin-4-yl)-4H-1,2,4-triazol-3-**
21 **yl)cyclobutyl)-2,3-dihydro-2-oxo-1H-benzo[d]imidazole-5-carbonitrile (16)**. *(trans)*-3-(6-
22 Cyano-1,2-dihydro-2-oxobenzo[d]imidazol-3-yl)cyclobutanecarbohydrazide (**31c**) (68.5 mg,
23 0.25 mmol) was added to a solution of Methyl(Z)-N-(2-chlorophenyl)pyrimidine-4-
24 carbimidothioate (**26**) (80 mg, 0.30 mmol) in N,N-dimethylacetamide (2 ml). Trifluoroacetic
25 acid (9.6 µl, 0.125 mmol) was added and the reaction mixture was heated to 120°C for 14
26 hours. Water was added and the reaction mixture was extracted with ethyl acetate. The
27 organic layer was washed with brine, dried over magnesium sulfate and concentrated under
28 reduced pressure. The crude solid was purified by preparative HPLC (C₁₈ reverse phase
29 column, elution with a water/MeCN gradient with 0.1% TFA). The fractions containing the
30 product were evaporated and lyophilized to yield 1-((*trans*)-3-(4-(2-chlorophenyl)-5-
31 (pyrimidin-4-yl)-4H-1,2,4-triazol-3-yl)cyclobutyl)-2,3-dihydro-2-oxo-1H-benzo[d]imidazole-
32 5-carbonitrile (**16**) as a white solid. The product was obtained as its trifluoroacetate salt.
33 Yield: 22.5 mg (15%); mp 252 - 255°C. HRMS for C₂₄H₁₇ClN₈O +H⁺, [M+H]⁺ m/z, calcd
34 469.1287, found 469.1275. ¹H NMR (600 MHz, DMSO-*d*₆) δ 11.37 (s, 1H), 8.97 (d, *J* = 5.2
35 Hz, 1H), 8.90 (s, 1H), 8.25 (d, *J* = 5.0 Hz, 1H), 7.74 (dd, *J* = 8.0, 1.7 Hz, 2H), 7.63 (td, *J* =
36 7.8, 1.6 Hz, 1H), 7.57 (dt, *J* = 7.6, 3.5 Hz, 2H), 7.45 (dd, *J* = 8.3, 1.6 Hz, 1H), 7.38 (s, 1H),
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 5.27 (p, $J = 8.9$ Hz, 1H), 3.48 (dq, $J = 10.0, 5.1, 3.7$ Hz, 1H), 3.10 (q, $J = 10.2$ Hz, 2H), 2.83
4
5 (ddt, $J = 12.3, 8.4, 3.7$ Hz, 1H), 2.73 (ddt, $J = 12.7, 8.5, 4.0$ Hz, 1H). ^{13}C NMR (151 MHz,
6
7 DMSO- d_6) δ 160.0, 158.9, 158.6, 154.2, 153.6, 151.2, 133.5, 133.1, 132.0, 131.3, 130.5,
8
9 130.2, 129.0, 128.9, 126.2, 120.1, 119.4, 112.1, 109.8, 103.2, 44.9, 31.7, 31.1, 24.2.
10
11
12
13

14 ASSOCIATED CONTENT

15
16
17 **Supporting Information.** The Supporting Information is available free of charge on the
18 ACS Publications website at DOI: XXXX. Experimental procedures for X-ray
19 crystallography, biochemical and pharmacological characterization, as well as analytical data
20 for all compounds. Molecular formula strings (CSV).
21
22
23
24
25

26
27 **Accession codes.** Coordinates and structure factors are deposited at the Protein Data Bank
28 with codes 5NSP (**14**) and 5NOB (**16**). The authors will release the atomic coordinates and
29 experimental data upon article publication.
30
31
32
33
34
35

36 AUTHOR INFORMATION

37 Corresponding Author

38
39
40
41 * M.N: Phone: +49 30 9406-3083. E-mail: nazare@fmp-berlin.de; S.K.: Phone: +47 23 07 90
42
43 17. E-mail: Stefan.KRAUSS@rr-research.no.
44
45

46 Present Addresses

47
48 [Dr. Upendra Rao Anumala, Research and Development, Amoli Organics Pvt. Ltd., ECP
49 Canal Road, Taluka Padra District, Vadodara, India.
50
51
52
53

54 Author Contributions

1
2
3 The manuscript was written through contributions of all authors. All authors have given
4 approval to the final version of the manuscript.
5
6
7

8 **Funding Sources**

9
10 The research was funded by Research Council of Norway “Extended biotarget validation
11 studies for the specific inhibitor of Wnt/ β -catenin signaling OD270”, the Jane and Aatos
12 Erkkö Foundation, Sigrid Jusélius Foundation, Biocenter Oulu and Academy of Finland
13 (grant No. 287063 and 294085 for L.L.) and the Russian Ministry of Education and Science
14 (grant No 14.N08.11.0056 for A.G.M. and S.L.).
15
16
17
18
19
20
21

22 **Notes**

23
24 The authors declare no competing financial interest.
25
26

27 **ACKNOWLEDGMENT**

28
29
30 Protein crystallography experiments were performed at the Diamond Light Source (Didcot,
31 UK) on beamline I02 and at European Synchrotron Radiation Facility (ESRF, Grenoble,
32 France) on beamline ID23-1. We are grateful to Local Contacts for providing assistance in
33 using beamlines. The research leading to these results has received funding from the
34 European Community's Seventh Framework Programme (FP7/2007-2013) under BioStruct-X
35 (grant agreement N°283570). The use of the facilities of the Biocenter Oulu Protein
36 crystallography core facility, a member of Biocenter Finland and Instruct-FI is gratefully
37 acknowledged. The authors thank Bernd Rupp und Raed Al-Yamori (cheminformatics and
38 software development) for the technical and scientific support in ChemAxon software
39 deployment.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 **ABBREVIATIONS**

1
2
3 ARTD, ADP ribosyl transferase; AXIN, axis inhibition protein; Balb/c, Bagg albino/c;
4
5 CLK2, CDC like kinase 2; COLO320, colon adenocarcinoma cell line 320; EDCI, 1-ethyl-3-
6
7 (3-dimethylaminopropyl)carbodiimide; GRB, genomic regulatory blocks; IRAP, insulin-
8
9 regulated aminopeptidase; MELK, maternal embryonic leucine zipper kinase; NuMA,
10
11 nuclear mitotic apparatus; PARP, poly (ADP-ribose) polymerase; PRKG1, protein kinase G1;
12
13 SAM sterile alpha motif; ST-Luc/Ren, superTOP-Luciferase/Renilla; SW480, human colon
14
15 cancer cell line; TNKS, telomere-associated poly-ADP ribose polymerase tankyrase; TRF,
16
17 telomere restriction fragment; TSF1, serine/threonine-protein kinase.
18
19
20

21 REFERENCES

- 22
23 1. Durkacz, B. W.; Irwin, J.; Shall, S. The Effect of Inhibition of (ADP-ribose)_n
24
25 Biosynthesis on DNA Repair Assayed by the Nucleoid Technique. *Eur. J. Biochem.* **1981**,
26
27 *121*, 65-69.
28
29
30 2. Durkacz, B. W.; Omidiji, O.; Gray, D. A.; Shall, S. (ADP-ribose)_n Participates in
31
32 DNA Excision Repair. *Nature* **1980**, *283*, 593-596.
33
34
35 3. Rouleau, M.; Patel, A.; Hendzel, M. J.; Kaufmann, S. H.; Poirier, G. G. PARP
36
37 Inhibition: PARP1 and Beyond. *Nat. Rev. Cancer* **2010**, *10*, 293-301.
38
39
40 4. Purnell, M. R.; Whish, W. J. Novel Inhibitors of Poly(ADP-ribose) Synthetase.
41
42 *Biochem. J.* **1980**, *185*, 775-777.
43
44
45 5. Hottiger, M. O.; Hassa, P. O.; Lüscher, B.; Schüler, H.; Koch-Nolte, F. Toward a
46
47 Unified Nomenclature for Mammalian ADP-Ribosyltransferases. *Trends Biochem. Sci.* **2010**,
48
49 *35*, 208-219.
50
51
52 6. Haikarainen T., Krauss S., Lehtio L. Tankyrases: Structure, Function and Therapeutic
53
54 Implications in Cancer. *Curr. Pharm. Des.* **2014**, *20*, 6472-6488.
55
56
57
58
59
60

- 1
2
3 7. Lehtiö, L.; Chi, N.-W.; Krauss, S. Tankyrases as Drug Targets. *FEBS J.* **2013**, *280*,
4 3576-3593.
5
6
7
8 8. Riffell, J. L.; Lord, C. J.; Ashworth, A. Tankyrase-targeted Therapeutics: Expanding
9 Opportunities in the PARP Family. *Nat. Rev. Drug Discovery* **2012**, *11*, 923-936.
10
11
12
13 9. Guettler, S.; LaRose, J.; Petsalaki, E.; Gish, G.; Scotter, A.; Pawson, T.; Rottapel, R.;
14 Sicheri, F. Structural Basis and Sequence Rules for Substrate Recognition by Tankyrase
15 Explain the Basis for Cherubism Disease. *Cell* **2011**, *147*, 1340-1354.
16
17
18
19
20 10. Cook, B. D.; Dynek, J. N.; Chang, W.; Shostak, G.; Smith, S. Role for the Related
21 Poly(ADP-Ribose) Polymerases Tankyrase 1 and 2 at Human Telomeres. *Mol. Cell. Biol.*
22 **2002**, *22*, 332-342.
23
24
25
26
27 11. Tanaka, N.; Mashima, T.; Mizutani, A.; Sato, A.; Aoyama, A.; Gong, B.; Yoshida, H.;
28 Muramatsu, Y.; Nakata, K.; Matsuura, M.; Katayama, R.; Nagayama, S.; Fujita, N.;
29 Sugimoto, Y.; Seimiya, H. APC Mutations as a Potential Biomarker for Sensitivity to
30 Tankyrase Inhibitors in Colorectal Cancer. *Mol. Cancer Ther.* **2017**, *16*, 752-762.
31
32
33
34
35
36
37 12. Lau, T.; Chan, E.; Callow, M.; Waaler, J.; Boggs, J.; Blake, R. A.; Magnuson, S.;
38 Sambrone, A.; Schutten, M.; Firestein, R.; Machon, O.; Korinek, V.; Choo, E.; Diaz, D.;
39 Merchant, M.; Polakis, P.; Holsworth, D. D.; Krauss, S.; Costa, M. A Novel Tankyrase
40 Small-Molecule Inhibitor Suppresses *APC* Mutation-Driven Colorectal Tumor Growth.
41 *Cancer Res.* **2013**, *73*, 3132-3144.
42
43
44
45
46
47
48 13. Huang, S.-M. A.; Mishina, Y. M.; Liu, S.; Cheung, A.; Stegmeier, F.; Michaud, G.
49 A.; Charlat, O.; Wiелlette, E.; Zhang, Y.; Wiessner, S.; Hild, M.; Shi, X.; Wilson, C. J.;
50 Mickanin, C.; Myer, V.; Fazal, A.; Tomlinson, R.; Serluca, F.; Shao, W.; Cheng, H.; Shultz,
51 M.; Rau, C.; Schirle, M.; Schlegl, J.; Ghidelli, S.; Fawell, S.; Lu, C.; Curtis, D.; Kirschner,
52
53
54
55
56
57
58
59
60

1
2
3 M. W.; Lengauer, C.; Finan, P. M.; Tallarico, J. A.; Bouwmeester, T.; Porter, J. A.; Bauer,
4 A.; Cong, F. Tankyrase Inhibition Stabilizes Axin and Antagonizes Wnt Signalling. *Nature*
5 **2009**, *461*, 614-620.
6
7

8
9
10 14. Zhong, L.; Ding, Y.; Bandyopadhyay, G.; Waaler, J.; Börgeson, E.; Smith, S.; Zhang,
11 M.; Phillips, S. A.; Mahooti, S.; Mahata, S. K.; Shao, J.; Krauss, S.; Chi, N.-W. The
12 Parsylation Activity of Tankyrase in Adipose Tissue Modulates Systemic Glucose
13 Metabolism in Mice. *Diabetologia* **2016**, *59*, 582-591.
14
15
16

17
18
19 15. Schneikert, J.; Behrens, J. The Canonical Wnt Signalling Pathway and its APC
20 Partner in Colon Cancer Development. *Gut* **2007**, *56*, 417-425.
21
22
23

24
25 16. Akhmetshina, A.; Palumbo, K.; Dees, C.; Bergmann, C.; Venalis, P.; Zerr, P.; Horn,
26 A.; Kireva, T.; Beyer, C.; Zwerina, J.; Schneider, H.; Sadowski, A.; Riener, M.-O.;
27 MacDougald, O. A.; Distler, O.; Schett, G.; Distler, J. H. W. Activation of Canonical Wnt
28 Signalling is Required for TGF- β -Mediated Fibrosis. *Nat. Commun.* **2012**, *3*, 735.
29
30
31
32

33
34 17. Angelova, M.; Zvezdaryk, K.; Ferris, M.; Shan, B.; Morris, C. A.; Sullivan, D. E.
35 Human Cytomegalovirus Infection Dysregulates the Canonical Wnt/ β -catenin Signaling
36 Pathway. *PLoS Pathog.* **2012**, *8*, e1002959.
37
38
39
40

41
42 18. Voronkov, A.; Holsworth, D. D.; Waaler, J.; Wilson, S. R.; Ekblad, B.; Perdreau-
43 Dahl, H.; Dinh, H.; Drewes, G.; Hopf, C.; Morth, J. P.; Krauss, S. Structural Basis and SAR
44 for G007-LK, a Lead Stage 1,2,4-Triazole Based Specific Tankyrase 1/2 Inhibitor. *J. Med.*
45 *Chem.* **2013**, *56*, 3012-3023.
46
47
48
49

50
51 19. Huang, H.; Guzman-Perez, A.; Acquaviva, L.; Berry, V.; Bregman, H.; Dovey, J.;
52 Gunaydin, H.; Huang, X.; Huang, L.; Saffran, D.; Serafino, R.; Schneider, S.; Wilson, C.;
53
54
55
56
57
58
59
60

1
2
3 DiMauro, E. F. Structure-Based Design of 2-Aminopyridine Oxazolidinones as Potent and
4
5 Selective Tankyrase Inhibitors. *ACS Med Chem Lett.* **2013**, *4*, 1218-1223.
6

7
8 20. Bregman, H.; Chakka, N.; Guzman-Perez, A.; Gunaydin, H.; Gu, Y.; Huang, X.;
9
10 Berry, V.; Liu, J.; Teffera, Y.; Huang, L.; Egge, B.; Mullady, E. L.; Schneider, S.; Andrews,
11
12 P. S.; Mishra, A.; Newcomb, J.; Serafino, R.; Strathdee, C. A.; Turci, S. M.; Wilson, C.;
13
14 DiMauro, E. F. Discovery of Novel, Induced-Pocket Binding Oxazolidinones as Potent,
15
16 Selective, and Orally Bioavailable Tankyrase Inhibitors. *J. Med. Chem.* **2013**, *56*, 4320-4342.
17

18
19 21. Shultz, M. D.; Majumdar, D.; Chin, D. N.; Fortin, P. D.; Feng, Y.; Gould, T.; Kirby,
20
21 C. A.; Stams, T.; Waters, N. J.; Shao, W. Structure-Efficiency Relationship of [1,2,4]Triazol-
22
23 3-ylamines as Novel Nicotinamide Isosteres that Inhibit Tankyrases. *J. Med. Chem.* **2013**, *56*,
24
25 7049-7059.
26

27
28 22. Shultz, M. D.; Cheung, A. K.; Kirby, C. A.; Firestone, B.; Fan, J.; Chen, C. H.-T.;
29
30 Chen, Z.; Chin, D. N.; DiPietro, L.; Fazal, A.; Feng, Y.; Fortin, P. D.; Gould, T.; Lagu, B.;
31
32 Lei, H.; Lenoir, F.; Majumdar, D.; Ochala, E.; Palermo, M. G.; Pham, L.; Pu, M.; Smith, T.;
33
34 Stams, T.; Tomlinson, R. C.; Touré, B. B.; Visser, M.; Wang, R. M.; Waters, N. J.; Shao, W.
35
36 Identification of NVP-TNKS656: The Use of Structure–Efficiency Relationships To
37
38 Generate a Highly Potent, Selective, and Orally Active Tankyrase Inhibitor. *J. Med. Chem.*
39
40 **2013**, *56*, 6495-6511.
41
42
43

44
45 23. Haikarainen, T.; Koivunen, J.; Narwal, M.; Venkannagari, H.; Obaji, E.; Joensuu, P.;
46
47 Pihlajaniemi, T.; Lehtio, L. Para-Substituted 2-Phenyl-3,4-dihydroquinazolin-4-ones as
48
49 Potent and Selective Tankyrase Inhibitors. *ChemMedChem* **2013**, *8*, 1978-1985.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 24. Haikarainen, T.; Waaler, J.; Ignatev, A.; Nkizinkiko, Y.; Venkannagari, H.; Obaji, E.;
4
5 Krauss, S.; Lehtiö, L. Development and Structural Analysis of Adenosine Site Binding
6
7 Tankyrase Inhibitors. *Bioorg. Med. Chem. Lett.* **2016**, *26*, 328-333.
8

9
10 25. Thorsell, A.-G.; Ekblad, T.; Karlberg, T.; Löw, M.; Pinto, A. F.; Trésaugues, L.;
11
12 Moche, M.; Cohen, M. S.; Schüler, H. Structural Basis for Potency and Promiscuity in
13
14 Poly(ADP-ribose) Polymerase (PARP) and Tankyrase Inhibitors. *J. Med. Chem.* **2017**, *60*,
15
16 1262-1271.
17

18
19 26. Bregman, H.; Gunaydin, H.; Gu, Y.; Schneider, S.; Wilson, C.; DiMauro, E. F.;
20
21 Huang, X. Discovery of A Class of Novel Tankyrase Inhibitors that Bind to Both the
22
23 Nicotinamide Pocket and The Induced Pocket. *J. Med. Chem.* **2013**, *56*, 1341-1345
24

25
26 27. Hua, Z. H.; Bregman, H.; Buchanan, J. L.; Chakka, N.; Guzman-Perez, A.; Gunaydin,
27
28 H.; Huang, X.; Gu, Y.; Berry, V.; Liu, J. Z.; Teffera, Y.; Huang, L. Y.; Egge, B.; Emkey, R.;
29
30 Mullady, E. L.; Schneider, S.; Andrews, P. S.; Acquaviva, L.; Dovey, J.; Mishra, A.;
31
32 Newcomb, J.; Saffran, D.; Serafino, R.; Strathdee, C. A.; Turci, S. M.; Stanton, M.; Wilson,
33
34 C.; DiMauro, E. F. Development of Novel Dual Binders as Potent, Selective, and Orally
35
36 Bioavailable Tankyrase Inhibitors. *J. Med. Chem.* **2013**, *56*, 10003-10015.
37

38
39 28. Nathubhai, A.; Haikarainen, T.; Koivunen, J.; Murthy, S.; Koumanov, F.; Lloyd, M.
40
41 D.; Holman, G. D.; Pihlajaniemi, T.; Tosh, D.; Lehtiö, L.; Threadgill, M. D. Highly Potent
42
43 and Isoform Selective Dual Site Binding Tankyrase/Wnt Signaling Inhibitors That Increase
44
45 Cellular Glucose Uptake and Have Antiproliferative Activity. *J. Med. Chem.* **2017**, *60*, 814-
46
47 820.
48

49
50 29. Waaler, J.; Machon, O.; von Kries, J. P.; Wilson, S. R.; Lundenes, E.; Wedlich, D.;
51
52 Gradl, D.; Paulsen, J. E.; Machonova, O.; Dembinski, J. L.; Dinh, H.; Krauss, S. Novel
53
54
55

1
2
3 Synthetic Antagonists of Canonical Wnt Signaling Inhibit Colorectal Cancer Cell Growth.
4

5 *Cancer Res.* **2011**, *71*, 197-205.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Graphical TOC

