New Bis(mercaptoimidazolyl)(pyrazolyl)borate Ligands and Their Zinc Complex Chemistry

Mouhai Shu,^[a] Rainer Walz,^[a] Biao Wu,^[a] Jan Seebacher,^[a] and Heinrich Vahrenkamp*^[a]

Keywords: Tripodal ligands / Zinc / Bioinorganic chemistry

Nine new tripodal NS₂ ligands of the bis(mercaptoimidazolyl)(pyrazolyl)borate type with varying 3-R-mercaptoimidazolyl moieties were prepared as their potassium salts. Treatment with zinc salts yielded the complex types L·Zn–Cl, L·Zn–I, L·Zn–ONO₂, L·Zn–OClO₃ and [L·Zn(imidazole)]ClO₄. Attempts at the formation of L·Zn–OH or cationic L·Zn complexes resulted in dismutation and formation of ZnL₂ complexes. Hydrolytic destruction yielded one [OZn₄(thiooimidazolate)₆] complex. The ZnS₂NO coordination which is present in the enzyme-substrate complex of alcohol dehydrogenase could be successfully modelled by an $[L-Zn(C_2H_5OH)]^+$ complex. The L-Zn-X complexes showed very low catalytic activity in the dehydrogenation of 2-propanol or the hydrogenation of *p*-nitrobenzaldehyde. The new compounds were identified by a total of 12 structure determinations.

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2003)

Introduction

The importance of tris(pyrazolyl)borate ligands for coordination chemistry can only be compared with that of cyclopentadienyl ligands in organometallic chemistry. This holds specifically for the classical or bioinorganic coordination chemistry of $zinc^{[1,2]}$ to which we have made contributions.^[3] The advantage of these tripodal ligands is that they quite reliably support an N₃ZnX coordination motif which has given them then name "tetrahedral enforcers" and which has made them so valuable for the bioinorganic chemistry of zinc.^[3,4]

The N₃Zn motif represents the attachment of a zinc ion to three histidine residues in a protein. Considering that cysteine is equally important for zinc binding in enzymes, one would like to have sulfur-containing tripodal ligands with qualities similar to those of the pyrazolylborates for a model complex chemistry. The accumulated experience with tripodal ligands containing thiolate functions has shown, however, that these are not suitable because the bridging tendency of the thiolates cannot be suppressed. In this context Reglinski's discovery of the tris(mercaptoimidazolyl)borates^[5] must be considered a breakthrough. As shown since by Reglinski himself^[6,7] and his competitors,^[8,9] including ourselves,^[10] these tripods are very good "tetrahedral enforcers" and allow a fruitful biomimetic zinc

 [a] Institut für Anorganische und Analytische Chemie der Universität Freiburg Albertstr. 21, 79104 Freiburg, Germany Fax: (internat.) + 49-761/203-6001 E-mail: vahrenka@uni-freiburg.de model complex chemistry. The reason for this is the fact that their sulfur donors are "tame" thiolates with a low bridging tendency due to their thiol-thione tautomerism as depicted below.



Having the easily accessible tris(pyrazolyl)- and tris(mercaptoimidazolyl)borates available as N_3 and S_3 donors, we decided to aim at their combination in the form of N_2S and NS_2 donors. Specifically, the NS_2 tripods should be valuable as they allow us to mimic the $(NS_2)ZnX$ coordination in alcohol dehydrogenase. With this in mind both the research group of Parkin and ourselves set out to study the chemistry of the bis(mercaptoimidazolyl)(pyrazolyl)borates. Parkin has published the synthesis of one such ligand and some basic metal complex chemistry of it.^[11–13] We have communicated the first correct structural model of the enzymesubstrate complex of alcohol dehydrogenase in the form of a (NS₂)Zn(C₂H₅OH) complex.^[14]

This paper summarises our recent work on bis(mercaptoimidazolyl)(pyrazolyl)borate ligands, their basic zinc complex chemistry and attempts to apply them for the modelling of the zinc enzyme alcohol dehydrogenase. Nine new ligands \mathbf{L} were prepared and their zinc complexes identified by a total of twelve structure determinations.



Results and Discussion

Ligand Synthesis

From our work with tris(pyrazolyl)borates we have the experience that their biomimetic zinc complex chemistry requires a good encapsulation of the metal ion. This is achieved by bulky substituents at the 3-positions of the pyrazole rings. We applied this for the new NS₂ ligands L by incorporating 5-methyl-3-phenylpyrazole units in L¹ to L⁸ and 3-*tert*-butyl-5-methylpyrazole in L⁹. Likewise, the thioimidazole units were made bulky with substituents at the outer nitrogen atoms, mostly with aromatic groups (L² to L⁸), but also with methyl (L¹) and *tert*-butyl (L⁹). Thus, all our NS₂ ligands are more bulky than the one with *N*-methylthioimidazole and unsubstituted pyrazole groups reported previously.^[13]

The synthesis of the tripodal ligands L from KBH₄ and the nitrogen heterocycles is favoured by the fact that the thioimidazoles react more readily with KBH₄ than the pyrazoles. Thus, lower reaction temperatures are possible which ensures better yields and which allows us to run the preparations in a one-pot procedure. Mixtures of KBH₄, pyrazole and thioimidazole (1:1:2) were allowed to react in a melt at ca. 160 °C to form L¹, L², L⁵ and L⁹. Later it turned out that the preparations in boiling toluene gave better yields and cleaner products. Furthermore, a 10% excess of the pyrazole was found to improve the yield of ligand L considerably. Ligands L³, L⁴, L⁶, L⁷ and L⁸ were prepared this way. The isolated compounds K[L] are colourless powders which are somewhat air- and light-sensitive, sparingly soluble in aromatic hydrocarbons, and which decompose upon melting between 160 and 200 °C.

Simple Zinc Complexes

Like the tris(pyrazolyl)- and tris(mercaptoimidazolyl)borates, the mixed ligands L easily form stable zinc halide complexes. This was applied for the products 1a-1f by treating the corresponding ligands with the zinc halides. FULL PAPER

The structures of 1c and 1d were determined, of which the one of 1c is displayed here (Figure 1). There is very close agreement in terms of bond lengths and angles between 1c and 1d, and the coordination environment of the zinc ion also resembles that in the previously reported complex L·ZnI of a related NS₂ ligand.^[11] The deviations from tetrahedral symmetry at the zinc ion are small. The largest bond angle is N-Zn-Cl, reflecting the steric hindrance between the phenyl substituent on the pyrazole and the Zn-Cl unit.



Figure 1. Molecular structure of complex 1c; selected bond lengths [Å] and angles [°]: Zn-Cl 2.201(1), Zn-Sl 2.335(1), Zn-S2 2.329(1), Zn-N 2.041(2); Cl-Zn-Sl 110.91(3), Cl-Zn-S2 109.11(3), Cl-Zn-N 120.09(6), Sl-Zn-S2 107.60(3), Sl-Zn-N 108.24(6), S2-Zn-N 99.82(6)

Other than the zinc halide and the zinc thiolate species,^[15] the simple L·Zn complexes with anions of oxo acids turned out to be quite labile. Attempts to prepare acetates invariably led to the formation of ZnL_2 complexes (see below). The same was observed when trying to prepare nitrates. Only those ligands L with sterically demanding substituents, typically in the *ortho* positions of the aromatic rings, yielded nitrate complexes, of which 2a-2c were isolated.

L ³ •ZnONO ₂	L^{6} -ZnONO ₂	L ⁸ •ZnONO ₂
2a	2b	2c

The structure determination of **2a** (Figure 2) proved the composition of the nitrate complexes and the semibidentate nature of the nitrate ligand. Again, the tetrahedral ligand environment is not seriously distorted, and again the N–Zn–O angle is unusually large due to steric hindrance. The semibidentate attachment of the nitrate coupled with its orientation with respect to the two sulfur donors makes the S1–Zn–O1 angle rather large and the S2–Zn–O1 angle rather small. In all these respects **2a** resembles the Zn–NO₃ complex of a related tris(mercaptoimidazolyl)borate ligand.^[10]



Figure 2. Molecular structure of complex **2a**; selected bond lengths [Å] and angles [°]: Zn–O1 1.965(5), Zn–O2 2.782(6), Zn–S1 2.307(2). Zn–S2 2.315(2), Zn–N 2.017(6); O1–Zn–S1 117.8(2), O1–Zn–S2 105.0(2), O1–Zn–N 115.3(2), S1–Zn–S2 108.6(1), S1–Zn–N 99.8(2), S2–Zn–N 110.2(2)

ZnL₂ Complexes

It has already been observed that the L·ZnX complexes of these NS₂ ligands are amenable to dismutation, with formation of the corresponding ZnL₂ complexes.^[13] Our investigations were also plagued frequently by this tendency. Many attempts to prepare L·Zn complexes with hard coligands did not yield the desired L·ZnX species, but instead ZnL₂ and, presumably, ZnX₂. The reason for this is the highly preferred ZnS₄ coordination in the uncharged ZnL₂ molecules. In the following we describe those reactions that led to ZnL₂ species, even though we performed no designed (viz. trivial) synthesis of a ZnL₂ complex.

According to one of the goals of this work, namely the modelling of zinc enzymes, we intended to prepare $[\mathbf{L}\cdot\mathbf{Zn}-\mathbf{OH}_2]^+$ or $\mathbf{L}\cdot\mathbf{Zn}-\mathbf{OH}$ complexes. The most straightforward way to do this^[3] is the reaction between a tripodal ligand **L** and hydrated zinc perchlorate, either directly or in combination with KOH. This was done here with \mathbf{L}^1 , \mathbf{L}^5 and \mathbf{L}^9 . Although some evidence could be gathered for the formation of $\mathbf{L}\cdot\mathbf{Zn}-\mathbf{OH}$ species (see below), they could not

be isolated in a pure form, and the final products of the reactions were the ZnL_2 complexes **3a**, **3d** and **3f**. Also, as mentioned above, the L·Zn-OAc complexes were not stable enough to be isolated, and in the case of L³ a clean dismutation yielded **3b**. Furthermore, the L·Zn-NO₃ complexes of those ligands L that are sterically less hindered, evaded their isolation by dismutation. In case of L⁴ and L⁷ this resulted in the isolation of pure **3c** and **3e**.

ZnL_2^1	ZnL^3_2	ZnL ⁴ 2
3a	3b	3c
$ZnL^{5}2$	ZnL ⁷ 2	ZnL ⁹ 2
3d	3e	3f

Structure determinations of **3a**, **3b**, **3c**, **3d** and **3f** revealed that in all cases the ligands L act as bidentate sulfur donors, i.e. the pyrazolyl arm of the tripod is uncoordinated. This corresponds to the observations made previously for similar ZnL_2 complexes^[13] and for those derived from the tris(mercaptoimidazolyl)borates.^[10] The molecular structures of **3a** and **3f** are shown as representative examples in Figures 3 and 4, respectively. In all complexes **3** the Zn-S bond lengths are in the range 2.33–2.40 Å, which is rather long but typical for this kind of ZnS₄ species.^[10,13] The bond angles at the zinc ion show quite a range around the tetrahedral angle, the extremes being 89° in **3a** and 125° in **3c**.

Hydrolysis Products

During our attempts to prepare $\mathbf{L} \cdot \mathbf{Zn} - \mathbf{OH}$ or $[\mathbf{L} \cdot \mathbf{Zn}]^+$ species by treating \mathbf{L} with $Zn(ClO_4)_2$ it also became obvious that the (mercaptoimidazolyl)borates are not only more easily formed than the pyrazolylborates but also more easily hydrolysed. Thus, the acidic conditions brought about by dissolving hydrated zinc perchlorate in alcohols frequently led to the partial destruction of the ligands \mathbf{L} , and free mercaptoimidazoles were found in the solutions. In three cases zinc complexes of the mercaptoimidazoles were the result of such hydrolytic cleavages.



Figure 3. Molecular structure of **3a**; selected bond lengths [Å] and angles [°]: Zn–S1 2.336(2), Zn–S2 2.382(2), Zn–S3 2.346(2), Zn–S4 2.401(2); S1–Zn–S2 110.77(8), S1–Zn–S3 115.68(8), S1–Zn–S4 114.64(8), S2–Zn–S3 112.93(7), S2–Zn–S4 89.28(6), S3–Zn–S4 110.70(7)

Figure 4. Molecular structure of **3f**; selected bond lengths [Å] and angles [°]: Zn-S1 2.359(2), Zn-S2 2.371(2), Zn-S3 2.368(2), Zn-S4 2.403(2); S1-Zn-S2 109.35(6), S1-Zn-S3 100.35(5), S1-Zn-S4 111.09(5), S2-Zn-S3 110.72(5), S2-Zn-S4 118.99(5), S3-Zn-S4 104.65(5)

In two of these cases the free mercaptoimidazole was incorporated as a coligand in the intended L·Zn complexes: Zn(ClO₄)₂·6H₂O and L³ yielded complex **4a**, and with L⁶ complex **4b** was formed. Both **4a** and **4b** were characterised by structure determinations. The structure of **4a** is shown in Figure 5. In **4a** and **4b** the tripod ligand is bound to the zinc ion in the normal fashion; the mercaptoimidazole is attached as a neutral molecule. Its bonding situation is that of a thioimidazoline, i.e. of a thiourea derivative. This is a very common situation in zinc complex chemistry, and the Zn-S bond length (2.33 Å in **4a**, 2.31 Å in **4b**) corresponds to expectation.^[16] The deviations from tetrahedral symmetry at the zinc ion (smallest angle 99°, largest angle 123°) are in the same range as observed above.



 $[L^{3}\cdot Zn(N-toly]$ thioimidazoline)]ClO₄ $[L^{6}\cdot Zn(N-cumy]$ thioimidazoline)]ClO₄ 4a 4b

FULL PAPER

In the case of L^1 the hydrolytic destruction resulted in a zinc complex that bears only mercaptoimidazolate ligands. After addition of KOH to the reaction mixture, $[OZn_4(me-thimazolate)_6]$ (5) was obtained, another crystalline form of which has already been described.^[17] Complex 5 is an analogue of $OZn_4(OAc)_6$,^[18] i.e. a variant of the classical basic beryllium acetate structure with the six chelating N,S li-



Figure 6. Molecular structure of 5.; selected bond lengths [Å]: Zn-O 1.955(2), Zn-N 2.000(4), Zn-S 2.325(1) and 2.340(1)

gands spanning the edges of the Zn_4 tetrahedron. Figure 6 shows the structure of **5**, all details of which have already been discussed.^[17]

Attempts at Enzyme Modelling

With the intention to prepare $L \cdot Zn - OH$, $L \cdot Zn(H_2O)$ or $[L \cdot Zn]^+$ complexes, i.e. functional models for the resting state of zinc enzymes, we performed a number of reactions between the ligands L and zinc salts of noncoordinating anions under varying pH conditions. So far, however, none of the intended species, which would be the starting point for modelling alcohol dehydrogenase chemistry, could be obtained in a pure form. However, in two cases evidence for the existence of a L·Zn–OH species could be gathered. Treatment of zinc perchlorate with L¹ and KOH allowed the precipitation of a compound whose elemental analyses and spectra may be in agreement with its formulation as 6a. Likewise, a species which may be formulated as 6b was obtained with L². Both compounds, however, show considerable amounts of impurities in their NMR spectra, and both could not be purified due to their lability toward hydrolysis or dismutation as described above.

Only with one ligand L, namely the *o*-methoxy-substituted L^8 , could stable complexes be prepared, which come close to containing naked $[L\cdot Zn]^+$. Treatment with zinc perchlorate in methanol yielded 7, and under the same conditions in ethanol 8 was obtained. The identification of 7 rests on its spectra. The presence of zinc-bound perchlorate is evident from the sharp and intense IR band at 1093

Figure 5. Molecular structure of **4a**; selected bond lengths [Å] and angles [°]: Zn-S1 2.323(2), Zn-S2 2.332(2), Zn-S3 2.326(2), Zn-N 2.027(5); N-Zn-S1 98.6(1), N-Zn-S2 105.6(1), N-Zn-S3 122.6(1), S1-Zn-S2 112.9(1), S1-Zn-S3 104.0(1), S2-Zn-S3 112.5(1)

Eur. J. Inorg. Chem. 2003, 2502-2511

www.eurjic.org

cm⁻¹, in analogy to the structurally characterised [tris(mercaptoimidazolyl)borate]zinc perchlorate complex.^[10] The ethanol complex **8** is a relative of two [tris(mercaptoimidazolyl)borate]zinc–alcohol complexes.^[10,19]

L8•ZnOClO3	$[\mathbf{L^{8\bullet}Zn}(\mathbf{C_{2}H_{5}OH})]\mathbf{ClO_{4}}$
7	8

The structure of 8 is shown in Figure 7. The ethanol ligand occupies the fourth coordination site of the zinc ion, the coordination geometry of which is tetrahedral to a good approximation. The Zn-O bond length is very similar to that in the nitrate complexes (see above), indicating a strong Zn-O(ethanol) bond. The ethanol ligand in the crystalline compound is connected by a hydrogen bond to a second ethanol molecule, which explains the absence of characteristic OH bands in the 3300-3700 cm⁻¹ region in the IR spectrum. The molecular shape of 8 does not offer a convincing clue why it is just ligand L^8 that makes complexes like 7 and 8 stable. Yet it is evident that the o-methoxy substituents enforce an orientation of the three phenyl groups that creates a bowl-like shape of the whole ligand, which may provide just enough stabilization of the Zn-EtOH unit in its interior.

Complex 8 is the first zinc complex in which an alcohol molecule is bound to a zinc ion with NS₂ ligation, creating a tetrahedral ZnONS₂ system. This bonding pattern reproduces the attachment of ethanol to the active centre of alcohol dehydrogenase.^[20] In our preliminary communication on $8^{[14]}$ we have elaborated the close geometrical relation between the ZnONS₂ cores of 8 and of the enzyme structure of highest resolution. Hence, in terms of bioinorganic chemistry, the formation and structure of 8 are the most relevant result of this paper (Figure 7).



Figure 7. Molecular structure of **8**; selected bond lengths [Å] and angles [°]: Zn-O 1.970(3), Zn-S1 2.282(1), Zn-S2 2.314(1), Zn-N 2.012(3); O-Zn-S1 105.4(1), O-Zn-S2 109.8(1), O-Zn-N 116.9(1), S1-Zn-S2 113.2(1), S1-Zn-N 111.7(1), S2-Zn-N 100.2(1)

Having achieved structural modelling of alcohol dehydrogenase we also attempted functional modelling. As done before,^[10,21,22] we used (benzyl)nicotinamide chloride ([BNA]Cl) as a mimic of NAD⁺ and (dihydrobenzyl)nicotinamide (BNAH) as a mimic of NADH. For attempts at oxidation we used 2-propanol as an electron-rich alcohol and for attempts at reduction p-nitrobenzaldehyde as an electron-poor aldehyde. Thus, the model reactions were:

$(CH_3)_2CHOH + BNA^+ \rightarrow (CH_3)_2CO + BNAH$	(1)
p -O ₂ N-C ₆ H ₄ CHO + BNAH \rightarrow p -O ₂ N-C ₆ H ₄ CH ₂ OH + BNA ⁺	(2)

It is known that reaction (1) does not proceed in the absence of a catalyst, but reaction (2) proceeds very slowly. Trying some of the L·Zn-X complexes described here we found that they show no or very low catalytic efficiency for both processes. The best results, obtained with complex 2a, consisted in a 2-3% yield of acetone and traces of *p*-nitrobenzyl alcohol after 24 h of heating. These are the poorest results in our studies on the functional modelling of alcohol dehydrogenase, although the applied "catalysts" are the closest structural models of the enzyme. Conversely, the best catalytic results were obtained by us with zinc complexes whose structure has no resemblance at all with that of the enzyme.^[22]

Conclusions

This work has shown that the tripodal NS_2 ligands L are easily accessible. They fill half of the gap between the analogous N_3 (i.e. pyrazolylborate) and S_3 (i.e. mercaptoimidazolylborate) ligands. Provided that the as-yet-unknown tripodal N_2S ligands of this type are also accessible, the door would be open for a complete set of comparative investigations of metal complexes with homologous tripodal N_3 , N_2S , NS_2 , and S_3 ligands. We intend to undertake this investigation with L·ZnX complexes, putting emphasis on enzyme model compounds.

The preliminary L-ZnX chemistry in this paper has revealed that the resulting complexes are not as inert as their tris(pyrazolylborate)·ZnX counterparts. One reason for this seems to be the high stability of the dismutation products ZnL₂ which comes from their ZnS₄ coordination in an uncharged complex. The other reason seems to be the geometrical disposition of the mercaptoimidazolyl moieties, which provides less encapsulation for the ZnX units than that of the pyrazolyl moieties, because their aromatic substituents point away from rather than enclosing the centre of the complexes. The relative stabilities of the L·ZnX complexes, however, give a clue on how to improve this situation: it seems that *ortho* substituents on the phenyl groups induce an orientation of them that creates a bowl-shaped environment around the ZnX units and hence the desired encapsulation.

The perchlorate complex 7 and the ethanol complex 8 are the best examples for this kind of stabilization. Complex 7 is as close as possible to being "naked" $[L \cdot Zn]^+$, and 8 is the best model so far for the enzyme-substrate complex of alcohol dehydrogenase. Both 7 and 8 also demonstrate the subtle stability differences in these complexes of weakly bound ligands: perchlorate is a better ligand than methanol (7 crystallises from methanol) but a weaker ligand than ethanol (8 crystallises from ethanol). Thus 7 and 8 on one side show the way to improve the ligand design for making more stable $\mathbf{L}\cdot\mathbf{Z}n$ complexes of this type, on the other side they offer themselves as starting materials for a functional and possibly catalytic $\mathbf{L}\cdot\mathbf{Z}n$ chemistry.

Experimental Section

General: For general working and measuring procedures, see ref.^[23] All procedures for the ligand syntheses were performed under an inert gas. 5-Methy-3-phenylpyrazole^[24] and 3-*tert*-butyl-5-methylpyrazole^[25] were prepared according to the published procedures. All *N*-substituted mercaptoimidazoles were prepared from the corresponding amino acetals RNHCH₂CH(OEt)₂ and HSCN according to an established procedure.^[26,27] They were characterised by ¹H NMR spectroscopy in CDCl₃ as follows:

R = methyl: ¹H NMR (CHCl₃): δ = 3.66 ppm (s, 3 H, NMe), 6.59 (d, J = 2.0 Hz, 1 H, 4-H), 6.78 (d, J = 2.2 Hz, 1 H, 5-H). R = phenyl: ¹H NMR (CHCl₃): $\delta = 6.75$ ppm (m, 2 H, 4-H, 5-H), 7.31-7.56 (m, 5 H, Ph). R = o-tolyl: ¹H NMR (CHCl₃): δ = 2.19 ppm (s, 3 H, Me), 6.66 (d, J = 2.4 Hz, 1 H, 4-H), 6.86 (d, J =2.4 Hz, 1 H, 5-H), 7.00-7.43 (m, Ar). R = p-tolyl: ¹H NMR (CHCl₃): $\delta = 2.41$ ppm (s, 3 H, Me), 6.80 (d, J = 1.6 Hz, 1 H, 4-H), 6.84 (d, J = 1.6 Hz, 1 H, 5-H), 7.28 (m, 2 H, Ar), 7.45 (m, 2 H, Ar). R = 2,4-xylyl: ¹H NMR (CHCl₃): δ = 2.16 ppm (s, 3 H, Me), 2.38 (s, 3 H, Me), 6.69 (d, J = 2.4 Hz, 1 H, 4-H), 6.81 (d, J = 2.4 Hz, 1 H, 5-H), 7.15 (m, 3 H, Ar), 7.26 (s, 1 H, Ar). R = *p*-cumyl: ¹H NMR (CHCl₃): $\delta = 1.28$ ppm (d, J = 6.8 Hz, 6 H, Me), 2.97 (sept, J = 6.8 Hz, 1 H, CH), 6.80 (d, J = 2.4 Hz, 1 H, 4-H), 6.84 (d, J = 2.4 Hz, 1 H, 5-H), 7.34 (d, J = 6.6 Hz, 2 H, Ar), 7.50 (d, J = 6.6 Hz, 2 H, Ar). R = *p*-anisyl: ¹H NMR (CHCl₃): $\delta = 3.85 \text{ ppm}$ (s, 3 H, OMe), 6.80 (d, J = 2.4 Hz, 1 H, 4-H), 6.98 (d, J = 2.2 Hz, 1 H, 5-H), 7.00 (d, J = 6.8 Hz, 2 H, Ar), 7.47 (d, J = 6.8 Hz, 2 H, Ar), 7.47 (d, J = 6.8 Hz, 2 H, Ar)J = 6.8 Hz, 2 H, Ar). R = o-anisyl: ¹H NMR (CHCl₃): δ = 3.74 ppm (s, 3 H, OMe), 6.81 (d, J = 2.1 Hz, 1 H, 4-H), 6.88 (d,J = 2.1 Hz, 1 H, 5-H), 6.98 (m, 2 H, Ar), 7.39 (m, 2 H, Ar). R = *tert*-butyl: ¹H NMR (CHCl₃): $\delta = 1.81$ ppm (s, 9 H, *t*Bu), 6.67 (d, J = 2.4 Hz, 1 H, 4-H), 6.84 (d, J = 2.4 Hz, 1 H, 5-H).

Ligand Synthesis: All ligands L were prepared by thermal reaction between KBH_4 , the pyrazole and the imidazole. Initially, the reactions were performed by melting the reagents (procedure A) which resulted in low yields and unsatisfactory purity of the products. Then it was found that the ligands L also form in boiling toluene (procedure B) which is the preferred procedure.

Procedure A: A 1:1 mixture of KBH₄ and the pyrazole was heated to 120 °C (thermometer immersed into the melt). The melt was stirred for 30 min. Then, 2 mol-equiv. of the mercaptoimidazole was added and the temperature raised slowly to 140 °C and kept at this temperature (30–60 min) until hydrogen evolution stopped. After cooling to room temp., the resulting solid was powdered and the raw product crystallised from ethanol at -30 °C.

Procedure B: A 1:1.1:2 mixture of KBH₄, the pyrazole and the imidazole was dried in vacuo at 60 °C for 2 h. 125 mL of dry toluene was added and the mixture refluxed for 4 days by heating with an oil bath (150 °C). Then, the volume of the solution was reduced in vacuo to 50 mL, and the mixture was poured into 200 mL of petroleum ether (boiling range 60–70 °C). The resulting precipitate was washed with a small amount of benzene and then with petroleum ether and dried in vacuo. L¹: By procedure A from KBH₄ (1.70 g, 31.6 mmol), pyrazole (5.00 g, 31.6 mmol) and the imidazole (7.21 g, 63.2 mmol). Yield 4.10 g (30%); m.p. 200 °C (dec.). $C_{18}H_{20}BKN_6S_2$ (434.44): calcd. C 49.77, H 4.64, N 19.34; found C 48.59, H 4.90, N 17.60. ¹H NMR (CDCl₃): $\delta = 2.33$ [s, 3 H, Me (pz)], 3.58 (s, 6 H, NMe), 6.30 [s, 1 H, H (pz)], 6.45 [d, J = 2.0 Hz, 2 H, H (im)], 6.63 [d, J = 2.0 Hz, 2 H, H (im)], 7.31 (m, 3 H, Ph), 7.51 (d, J = 7.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2392$ m (BH) cm⁻¹.

L²: By procedure A from KBH₄ (920 mg, 17.1 mmol), the pyrazole (2.70 g, 17.1 mmol) and the imidazole (6.03 g, 34.2 mmol). Yield 4.9 g (50%); m.p. 180 °C (dec.). $C_{28}H_{24}BKN_6S_2$ (558.58): calcd. C 60.21, H 4.33, N 15.05; found C 56.50, H 4.83, N 13.69. ¹H NMR (CDCl₃): $\delta = 2.29$ [s, 3 H, Me (pz)], 6.29 [s, 2 H, H (pz)], 6.72 (d, J = 2.4 Hz, 2 H, H (im)], 6.75 [d, J = 2.4 Hz, 2 H, H (im)], 7.30–7.51 (m, 13 H, Ar), 7.53 (d, J = 7.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2483$ m (BH) cm⁻¹.

L³: By procedure B from KBH₄ (1.08 g, 20.0 mmol), the pyrazole (3.47 g, 22.0 mmol) and the imidazole (7.60 g, 40.0 mmol). Yield 10.1 g (86%); m.p. 205 °C (dec.). $C_{30}H_{28}BKN_6S_2$ (586.63): calcd. C 61.42, H 4.81, N 14.33, S 10.93; found C 61.36, H 5.12, N 14.23, S 10.47. ¹H NMR (CD₃OD): $\delta = 2.16$ [s, 6 H, Me (tol)], 2.43 [s, 3 H, Me (pz)], 6.31 [s, 1 H, H (pz)], 6.59 [d, J = 2.2 Hz, 2 H, H (im)], 6.79 [d, J = 2.2 Hz, 2 H, H (im)], 7.27 (m, 8 H, Ar), 7.75 (d, J = 7.0 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2451$ m (BH) cm⁻¹.

L⁴: By procedure B from KBH₄ (1.08 g, 20.0 mmol), the pyrazole (3.47 g, 22.0 mmol) and the imidazole (7.60 g, 40.0 mmol). Yield 9.86 g (84%); m.p. 165 °C (dec.). $C_{30}H_{28}BKN_6S_2$ (586.63): calcd. C 61.42, H 4.81, N 14.33, S 10.93; found C 61.29, H 5.25, N 14.25, S 9.91. ¹H NMR (CD₃OD): $\delta = 2.37$ [s, 6 H, Me (tol)], 2.42 [s, 3 H, Me (pz)], 6.30 [s, 1 H, H (pz)], 6.57 [d, J = 2.2 Hz, 2 H, H (im)], 6.92 [d, J = 2.2 Hz, 2 H, H (im)], 7.28 (m, 8 H, Ar), 7.45 [d, J = 8.2 Hz, 4 H, H (tol)], 7.74 (d, J = 7.0 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{\nu} = 2482$ m (BH) cm⁻¹.

L⁵: By procedure A from KBH₄ (0.54 g, 10.0 mmol), the pyrazole (3.16 g, 20.0 mmol) and the imidazole (8.16 g, 40.0 mmol). Yield 0.74 g (12%); m.p. 185 °C (dec.). $C_{32}H_{32}BKN_6S_2$ (614.69): calcd. C 62.53, H 5.25, N 13.67; found C 58.72, H 5.31, N 12.33. ¹H NMR (CDCl₃): $\delta = 2.12$ [s, 6 H, Me (xyl)], 2.34 [s, 6 H, Me (xyl)], 2.75 [s, 3 H, Me (pz)], 6.36 [s, 1 H, H (pz)], 6.54 [d, J = 2.0 Hz, 2 H, H (im)], 6.68 [d, J = 2.0 Hz, 2 H, H (im)], 7.04–7.15 (m, 6 H, xyl), 7.40–7.43 (m, 3 H, Ph), 7.69–7.75 (m, 2 H, Ph) ppm. IR (KBr): $\tilde{\nu} = 2487$ m (BH) cm⁻¹.

L⁶: By procedure B from KBH₄ (1.08 g, 20.0 mmol), the pyrazole (3.47 g, 22.0 mmol) and the imidazole (8.72 g, 40.0 mmol). Yield 4.63 g (36%); m.p. 165 °C (dec.). $C_{34}H_{36}BKN_6S_2$ (642.74): calcd. C 63.54, H 5.65, N 13.08, S 9.98; found C 63.21, H 5.91, N 13.07, S 10.08. ¹H NMR (CD₃OD): $\delta = 1.22$ [d, J = 6.8 Hz, 18 H, Me (*i*Pr)], 2.08 [s, 3 H, Me (pz)], 2.89 [sept, J = 6.8 Hz, 2 H, CH (*i*Pr)], 6.26 [s, 1 H, H (pz)], 6.37 [d, J = 2.2 Hz, 2 H, H (im)], 6.75 [d, J = 2.2 Hz, 2 H, H (im)], 7.24 (m, 8 H, Ar), 7.42 (d, J = 8.3 Hz, 4 H, Cum), 7.68 (d, J = 7.0 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2476$ m (BH) cm⁻¹.

L⁷: By procedure B from KBH₄ (0.54 g, 10.0 mmol), the pyrazole (1.74 g, 11.0 mmol) and the imidazole (4.12 g, 20.0 mmol). Yield 5.57 g (90%); m.p. 175 °C (dec.). $C_{30}H_{28}BKN_6O_2S_2$ (618.63): calcd. C 58.25, H 4.56, N 13.58, S 10.37; found C 57.68, H 4.42, N 13.20, S 10.13. ¹H NMR (CD₃OD): $\delta = 2.44$ [s, 3 H, Me (pz)], 3.82 (s, 6 H, OMe), 6.29 [s, 1 H, H (pz)], 6.57 [d, J = 2.2 Hz, 2 H, H (im)], 6.90 [d, J = 2.2 Hz, 2 H, H (im)], 7.00 (d, J = 8.4 Hz, 4 H, C₆H₄),

FULL PAPER

7.31 (m, 3 H, Ph), 7.44 (d, J = 8.3 Hz, 4 H, C₆H₄), 7.72 (d, J = 7.0 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2486$ m (BH) cm⁻¹.

L⁸: By procedure B from KBH₄ (0.70 g, 13.0 mmol), the pyrazole (2.26 g, 14.3 mmol) and the imidazole (5.35 g, 26.0 mmol). Yield 6.79 g (85%); m.p. 185 °C (dec.). $C_{30}H_{28}BKN_6O_2S_2$ (618.63): calcd. C 58.25, H 4.56, N 13.58, S 10.37; found C 57.99, H 4.60, N 13.54, S 10.43. ¹H NMR ([D₆]DMSO): $\delta = 2.43$ [s, 3 H, Me (pz)], 3.73 (s, 6 H, OMe), 6.24 [s, 1 H, H (pz)], 6.76 [d, J = 2.1 Hz, 2 H, H (im)], 6.82 [d, J = 2.1 Hz, 2 H, H (im)], 6.85–7.49 (m, 1 H, Ar), 7.74 (d, J = 7.1 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2441$ w (BH) cm⁻¹.

L⁹: By procedure A from (0.54 g, 10.0 mmol) of KBH₄, (3.00 g, 20.0 mmol) of the pyrazole and (6.24 g, 40.0 mmol) of the imidazole. Yield 0.51 g (10%), m.p. 160 °C (dec.). The product contained considerable amounts of impurities. ¹H NMR ([D₆]DMSO): δ = 1.18 [s, 9 H, *t*Bu (pz)], 1.73 [s, 18 H, *t*Bu (im)], 2.15 [s, 3 H, Me (pz)], 5.59 [s, 1 H, H (pz)], 6.26 [d, *J* = 2.0 Hz, 2 H, H (im)], 6.79 [d, *J* = 2.0 Hz, 2 H, H (im)] ppm. IR (KBr): \tilde{v} = 2496m (BH) cm⁻¹.

Zinc Halide Complexes. General Procedure: A solution of the zinc halide in 10 mL of methanol was added dropwise, with stirring, to a solution of the ligand in 50 mL of dichloromethane. After 2 h of stirring, the solution was filtered, the filtrate reduced to half of its volume in vacuo and then cooled to -30 °C overnight. The resulting precipitate of the raw product was filtered off, washed with 1-2 mL of methanol and dried in vacuo. If necessary it was recrystallised from CH₂Cl₂/CH₃OH or CH₂Cl₂/CH₃CN. All L·ZnHal complexes were obtained as colourless powders.

1a: From L¹ (200 mg, 0.46 mmol) and ZnCl₂ (63 mg, 0.46 mmol). Yield 57 mg (25%); m.p. 225 °C (dec.). $C_{18}H_{20}BCIN_6S_2Zn$ (496.20): calcd. C 43.57, H 4.06, N 16.94; found C 42.86, H 4.13, N 16.32. ¹H NMR (CDCl₃): $\delta = 2.42$ [s, 3 H, Me (pz)], 3.57 [s, 6 H, Me (im)], 6.22 [s, 1 H, H (pz)], 6.60 [d, J = 2.0 Hz, 2 H, H (im)], 7.15 [d, J = 2.0 Hz, 2 H, H (im)], 7.28 (m, 3 H, Ph), 7.74 (d, J = 7.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2503$ m (BH) cm⁻¹.

1b: From L² (300 mg, 0.52 mmol) and ZnCl₂ (71 mg, 0.52 mmol). Yield 156 mg (48%); m.p. 215 °C (dec.). $C_{28}H_{24}BCIN_6S_2Zn$ (620.33): calcd. C 54.21, H 3.90, N 13.55; found C 54.68, H 4.05, N 13.57. ¹H NMR (CDCl₃): $\delta = 2.29$ [s, 3 H, Me (pz)], 6.29 [s, 1 H, H (pz)], 6.75 [d, J = 2.4 Hz, 2 H, H (im)], 6.81 [d, J = 2.4 Hz, 2 H, H (im)], 7.30–7.51 (m, 13 H, Ar), 7.53 (d, J = 7.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2507$ m (BH) cm⁻¹.

1c: From L³ (196 mg, 0.33 mmol) and ZnCl₂ (46 mg, 0.34 mmol). Yield 180 mg (84%); m.p. 270 °C (dec.). $C_{30}H_{28}BCIN_6S_2Zn$ (648.38): calcd. C 55.57, H 4.35, N 12.96, S 9.89; found C 55.23, H 4.58, N 13.33, S 9.34. ¹H NMR (CDCl₃): $\delta = 2.06$ [s, 3 H, Me¹ (tol)], 2.28 [s, 3 H, Me² (tol)], 2.49 [s, 3 H, Me (pz)], 6.39 [s, 1 H, H (pz)], 6.88 [s, 2 H, H (im)], 7.05 [s, 2 H, H (im)], 7.25-7.37 (m, 11 H, Ar), 7.53 (d, J = 7.2 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{\nu} = 2559$ w (BH) cm⁻¹.

1d: From L⁵ (614 mg, 1.00 mmol) and ZnCl₂ (136 mg, 1.00 mmol). Yield 0.25 mg (37%), m.p. 220 °C (dec.). $C_{32}H_{32}BCIN_6S_2Zn$ (676.43): calcd. C 56.82, H 4.77, N 12.42; found C 56.99, H 4.70, N 12.36. ¹H NMR (CDCl₃): $\delta = 2.14$ [s, 6 H, Me (xyl)], 2.37 [s, 6 H, Me (xyl)], 2.48 [s, 3 H, Me (pz)], 6.38 [s, 1 H, H (pz)], 6.83 [d, J = 2.0 Hz, 2 H, H (im)], 6.99 [d, J = 2.0 Hz, 2 H, H (im)], 7.00–7.17 (m, 6 H, xyl), 7.31–7.37 (m, 3 H, Ph), 7.53–7.58 (m, 2 H, Ph) ppm. IR (KBr): $\tilde{\nu} = 2505$ m (BH) cm⁻¹.

1e: From L^{6} (206 mg, 0.33 mmol) and ZnCl₂ (46 mg, 0.34 mmol). Yield 169 mg (67%), m.p. 240 °C (dec.). $C_{36}H_{40}BClN_6S_3Zn$ (764.60): calcd. C 56.55, H 5.27, N 10.99, S 12.58; found C 56.31, H 5.60, N 11.02, S 12.25. ¹H NMR (CDCl₃): δ = 1.26 [d, *J* = 7.0 Hz, 12 H, Me (*i*Pr)], 2.93 [sept, *J* = 7.0 Hz, 2 H, CH (*i*Pr)], 6.38 [s, 1 H, H (pz)], 7.01 [d, *J* = 2.0 Hz, 2 H, H (im)], 7.07 [d, *J* = 2.0 Hz, 2 H, H (im)], 7.28 (m, 8 H, Ar), 7.45 (m, 3 H, Ar), 7.62 (d, *J* = 7.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{\nu}$ = 2512 w (BH) cm⁻¹.

1f: From L⁸ (301 mg, 0.49 mmol) and ZnI₂ (155 mg, 0.49 mmol). Yield 257 mg (69%), m.p. 230 °C (dec.). $C_{30}H_{28}BIN_{6}$ -O₂S₂Zn·0.5(CH₂Cl₂) (771.83 +42.47): calcd. C 44.99, H 3.59, N 10.32, S 7.88; found C 45.79, H 3.95, N 10.75, S 8.25. ¹H NMR (CDCl₃): δ = 2.49 [s, 3 H, Me (pz)], 3.85 (s, 6 H, OMe), 5.32 (s, 1 H, CH₂Cl₂), 6.15 [s, 1 H, H (pz)], 6.91–7.95 [m, 15 H, H (im) + Ar] ppm. IR (KBr): $\tilde{\nu} = 2511$ w (BH) cm⁻¹.

Zinc Nitrate Complexes

2a: A solution of L₃ (196 mg, 0.33 mmol) in 5 mL of dichloromethane was added dropwise with stirring to a solution of Zn(NO₃)₂·6H₂O (95 mg, 0.34 mmol) in 10 mL of methanol. After 4 h of stirring, the precipitate was filtered off. Recrystallisation from dichloromethane/acetonitrile yielded 158 mg (71%) of **2a**, m.p. 220 °C (dec.). C₃₀H₂₈BN₇O₃S₂Zn (674.93): calcd. C 53.39, H 4.18, N 14.53, S 9.50; found C 53.29, H 4.29, N 14.42, S 9.30. ¹H NMR (CD₂Cl₂): $\delta = 1.96$ [s, 3 H, Me¹ (tol)], 2.25 [s, 3 H, Me² (tol)], 2.53 [s, 3 H, Me (pz)], 6.46 [s, 1 H, H (pz)], 6.96 [d, J = 2.2 Hz, 2 H, H (im)], 7.08 [d, J = 2.2 Hz, 2 H, H (im)], 7.37 (m, 11 H, Ar), 7.52 (d, J = 6.0 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2479$ w (BH), 1382 vs (NO₃), 1294 vs (NO₃) cm⁻¹.

2b: Like **2a** from L⁶ (214 mg, 0.33 mmol) and Zn(NO₃)₂·6H₂O (95 mg, 0.34 mmol). Yield 169 mg (73%); m.p. 180 °C (dec.). C₃₄H₃₆BN₇O₃S₂Zn (731.04): calcd. C 55.86, H 4.96, N 13.41, S 8.77; found C 56.26, H 5.46, N 12.87, S 8.01. ¹H NMR (CDCl₃): $\delta = 1.27$ [d, J = 6.9 Hz, 12 H, Me (*i*Pr)], 2.96 [sept, J = 6.9 Hz, 2 H, CH (*i*Pr)], 6.40 [s, 1 H, H (pz)], 7.00 [s, 4 H, H (im)], 7.36 (s, 8 H, Cum), 7.38 (m, 3 H, Ph), 7.56 (d, J = 5.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2486$ w (BH), 1373 s (NO₃), 1292 s (NO₃) cm⁻¹.

2c: Like **2a** from L⁸ (307 mg, 0.52 mmol) and Zn(NO₃)₂·4H₂O (137 mg, 0.52 mmol). Yield 115 mg (31%), m.p. 215 °C (dec.). C₃₀H₂₈BN₇O₅S₂Zn (706.93): calcd. C 50.97, H 3.99, N 13.87, S 9.07; found C 50.43, H 4.28, N 13.51, S 8.93. ¹H NMR (CDCl₃): $\delta = 2.44$ [s, 3 H, Me (pz)], 3.75 (s, 6 H, OMe), 6.32 [s, 1 H, H (pz)], 6.86 [d, J = 2.2 Hz, 2 H, H (im)], 6.96 [d, J = 2.2 Hz, 2 H, H (im)], 7.10–7.48 (m, 11 H, Ar), 7.54–7.60 (m, 2 H, Ph) ppm. IR (KBr): $\tilde{\nu} = 2489$ w (BH), 1384 vs (NO₃), 1294 vs (NO₃) cm⁻¹.

ZnL₂ Complexes

3a: A solution of (171 mg, 0.46 mmol) of Zn(ClO₄)₂·6H₂O in 20 mL of methanol was added with stirring to a solution of (200 mg, 0.46 mmol) of L^1 in 50 mL of dichloromethane. After 30 min, powdered KOH (29 mg, 0.50 mmol) was added. After 20 h of stirring and filtration, the volume of the solution was reduced to one half in vacuo. Cooling to -4 °C overnight produced a colourless precipitate of 6a (characterisation see below). Attempts to purify this product by dissolution in dichloromethane and slow concentration yielded, after 4 weeks, 18 mg (4%) of colourless crystalline 3a; m.p. 230 °C (dec.). $C_{36}H_{40}B_2N_{12}S_4Zn \cdot 3H_2O$ (856.14 +54.02): calcd. C 47.51, H 5.09, N 18.47; found C 46.40, H 5.03, N 19.25. ¹H NMR (CDCl₃): $\delta = 1.49$ (s, 6 H, H₂O), 2.38 [s, 6 H, Me (pz)], 3.43 [s, 12 H, Me (im)], 6.22 [s, 2 H, H (pz)], 6.67 [d, J =2.0 Hz, 4 H, H (im)], 7.17 [d, J = 2.0 Hz, 4 H, H (im)], 7.31 (m, 6 H, Ph), 7.75 (d, J = 7.4 Hz, 4 H, Ph) ppm. IR (KBr): $\tilde{v} = 2500$ m (BH) cm^{-1} .

3b: A solution of L³ (196 mg, 0.33 mmol) in methanol (5 mL) was added dropwise with stirring to a solution of $Zn(CH_3COO)_2$ (62 mg, 0.34 mmol) in methanol (10 mL). After 4 h of stirring, the precipitate of raw **3b** was filtered off. Crystallisation from methanol/dichloromethane by slow concentration yielded 146 mg (76%) of **3b** as colourless crystals; m.p. 250 °C (dec.). $C_{60}H_{56}B_2N_{12}S_4Zn$ (1160.46): calcd. C 62.10, H 4.86, N 14.48, S 11.05; found C 61.90, H 4.90, N 14.56, S 11.27. ¹H NMR (CDCl₃): $\delta = 1.97$ [s, 3 H, Me (tol)], 2.04 [s, 3 H, Me (tol)], 2.53 [s, 3 H, Me (pz)], 6.33 [s, 1 H, H (pz)], 6.59 [d, J = 2.0 Hz, 2 H, H (im)], 6.68 [d, J = 2.0 Hz, 2 H, H (im)], 7.18–7.41 (m, 11 H, Ar), 7.89 (d, J = 7.2 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2521$ m (BH) cm⁻¹.

3c: Like **3b** from L⁴ (196 mg, 0.33 mmol) and Zn(NO₃)₂·6H₂O (95 mg, 0.34 mmol). Yield 159 mg (83%) of **3c** as colourless crystals; m.p. 240 °C (dec.). $C_{60}H_{56}B_2N_{12}S_4Zn$ (1160.46): calcd. C 62.10, H 4.86, N 14.48, S 11.05; found C 61.10, H 4.74, N 14.24, S 10.56. ¹H NMR (CDCl₃): $\delta = 2.43$ [s, 6 H, Me (tol)], 2.75 [s, 3 H, Me (pz)], 6.38 [s, 1 H, H (pz)], 6.84 [d, J = 2.2 Hz, 2 H, H (im)], 6.94 [d, J = 2.2 Hz, 2 H, H (im)], 7.10–7.45 (m, 11 H, Ar), 7.74 (d, J = 7.2 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2477$ w (BH) cm⁻¹.

3d: A solution of $Zn(ClO_4)_2 \cdot 6H_2O$ (372 mg, 1.00 mmol) in methanol (10 mL) was slowly added dropwise with stirring to a solution of L⁵ (614 mg, 1.00 mmol) in dichloromethane (10 mL). After 2 h of stirring, all volatiles were removed in vacuo. The residue was extracted with four 5-mL portions of dichloromethane. The combined extracts were concentrated to dryness and the residue dissolved in a minimum amount of benzene. Slow concentration produced a precipitate of crystalline solvated 3d. Filtration and drying for 3 h in vacuo yielded 200 mg (32%) of 3d as a colourless powder; m.p. 230 °C (dec.). C₆₄H₆₄B₂N₁₂S₄Zn (1216.58): calcd. C 63.19, H 5.30, N 13.82; found C 63.26, H 5.66, N 12.80. ¹H NMR (CDCl₃): $\delta = 2.00$ [s, 6 H, Me (xyl)], 2.21 [s, 6 H, Me (xyl)], 2.94 [s, 3 H, Me (pz)], 6.15 [s, 1 H, H (pz)], 6.25 [d, J = 2.0 Hz, 2 H, H (im)], 6.55 [d, J = 2.0 Hz, 2 H, H (im)], 6.79 - 7.00 (m, 6 H, xyl), 7.45 - 7.53(m, 3 H, Ph), 8.32–8.35 (m, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2513$ m (BH) cm^{-1} .

3e: Like **3b** from L⁷ (206 mg, 0.33 mmol) and Zn(NO₃)₂·6H₂O (95 mg, 0.34 mmol). Yield 127 mg (63%) of **3e** as colourless crystals; m.p. 245 °C (dec.). $C_{60}H_{56}B_2N_{12}O_4S_4Zn$ (1224.46): calcd. C 58.86, H 4.61, N 13.73, S 10.48; found C 58.54, H 4.83, N 13.52, S 10.12. ¹H NMR (CD₂Cl₂): $\delta = 2.75$ [s, 3 H, Me (pz)], 3.83 (s, 6 H, OMe), 6.38 [s, 1 H, H (pz)], 6.83 [s, 2 H, H (im)], 6.96 (d, J = 8.4 Hz, 4 H, anisyl), 7.44 (d, J = 8.4 Hz, 4 H, anisyl), 7.46 (m, 3 H, Ph), 7.74 (d, J = 7.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2488$ m (BH) cm⁻¹.

3f: Like **3d** from Zn(ClO₄)₂·6H₂O (372 mg, 1.00 mmol) and L⁹ (500 mg, 1.00 mmol). Recrystallisation from acetonitrile yielded 120 mg (24%) of **3f** as colourless crystals; m.p. 180 °C (dec.). $C_{44}H_{72}B_2N_{12}S_4Zn$ (984.40): calcd. C 53.69, H 7.37, N 17.07, S 13.03; found C 52.61, H 6.84, N 16.98, S 12.56. ¹H NMR (CD₂Cl₂): $\delta = 1.28$ [s, 9 H, *t*Bu (pz)], 1.70 [s, 18 H, *t*Bu (im)], 2.44 [s, 3 H, Me (pz)], 5.78 [s, 1 H, H (pz)], 6.82 [d, J = 2.0 Hz, 2 H, H (im)], 6.87 [d, J = 2.2 Hz, 2 H, H (im)] ppm. IR (KBr): $\tilde{v} = 2520$ m (BH) cm⁻¹.

Hydrolysis Products

4a: A solution of L^3 (196 mg, 0.33 mmol) in dichloromethane (5 mL) was added dropwise, with stirring, to a solution of $Zn(ClO_4)_2$ ·6H₂O (124 mg, 0.34 mmol) in ethanol (10 mL). After stirring for 24 h, all volatiles were removed in vacuo, the residue extracted with dichloromethane, filtered and the filtrate concen-

trated to dryness again. Recrystallisation from dichloromethane/ ethanol yielded 110 mg (35%) of **4a** as colourless crystals; m.p. 210 °C (dec.). $C_{40}H_{38}BCIN_8O_4S_3Zn$ (902.64): calcd. C 53.23, H 4.24, N 12.41, S 10.66; found C 51.94, H 4.36, N 11.95, S 10.02. ¹H NMR (CD₂Cl₂): $\delta = 1.78$ [s, 3 H, Me (tol)], 1.96 [s, 3 H, Me (tol)], 2.12 [s, 3 H, Me (tol)], 2.49 [s, 3 H, Me (pz)], 6.40 [s, 1 H, H (pz)], 6.72 [s, 1 H, H (im)], 6.73 [d, J = 2.2 Hz, 1 H, H (im)], 6.80 [s, 1 H, H (im)], 6.84 [s, 1 H, H (im)], 7.00 [s, 2 H, H (im)], 7.10–7.39 (m, 17 H, Ar), 10.64 (s, 1 H, NH) ppm. IR (KBr): $\tilde{v} = 2463$ w (BH) cm⁻¹.

4b: Like **4a** from L⁶ (214 mg, 0.33 mmol) and Zn(ClO₄)₂·6H₂O (124 mg, 0.34 mmol). Yield 88 mg (27%) of **4b** as colourless crystals, m.p. 180 °C (dec.). C₄₆H₅₀BClN₈O₄S₃Zn (986.80): calcd. C 55.99, H 5.11, N 11.36, S 9.75; found C 55.27, H 5.52, N 11.10, S 9.07. ¹H NMR (CDCl₃): $\delta = 1.28$ [d, J = 7.0 Hz, 18 H, Me(*i*Pr)], 2.52 [s, 3 H, Me (pz)], 2.97 [sept, J = 7.0 Hz, 3 H, CH(*i*Pr)], 6.39 [s, 1 H, H (pz)], 6.83 [s, 1 H, H (im)], 6.94 [s, 1 H, H (im)], 6.99 [s, 2 H, H (im)], 7.14 [s, 2 H, H (im)], 7.29 (d, J = 8.0 Hz, 6 H, C₆H₄), 7.39 (m, 3 H, Ph), 7.43 (d, J = 8.0 Hz, 6 H, C₆H₄), 7.47 (d, J = 7.8 Hz, 2 H, Ph), 10.92 (s, 1 H, NH) ppm. IR (KBr): $\tilde{v} = 2468$ w (BH) cm⁻¹.

5: A solution of L¹ (350 mg, 0.80 mmol) in dichloromethane (50 mL) was added dropwise, with stirring, to a solution of Zn(ClO₄)₂·6H₂O (300 mg, 0.18 mmol) in methanol (100 mL). After 2 h of stirring, powdered KOH (46 mg, 0.80 mmol) was added and the mixture stirred for another 20 h. After filtration, the filtrate was concentrated to half its volume. Within a few weeks 74 mg (9%) of colourless crystalline **5**, m.p. 167 °C, was obtained. C₂₄H₃₀N₁₂OS₆Zn₄·6H₂O (956.50 + 108.06): calcd. C 27.08, H 3.98, N 15.79; found C 27.03, H 3.26, N 16.59. ¹H NMR (CDCl₃): δ = 1.51 (s, 2 H, H₂O), 3.49–3.69 (m, 3 H, Me), 6.78–7.10 (m, 2 H, H (im) ppm.

Enzyme Models

6a: The precipitate of **6a**, which is an intermediate in the formation of **3a** (see above), was dried in vacuo. 46 mg (21%) of raw **6a** remained as a colourless powder; m.p. 220 °C (dec.). $C_{18}H_{21}BN_6OS_2Zn$ (477.74): calcd. C 45.25, H 4.43, N 17.59, S 13.42; found C 43.44, H 4.16, N 17.56, S 12.66. ¹H NMR (CDCl₃): $\delta = 2.36$ [s, 3 H, Me (pz)], 3.50 [s, 6 H, Me (im)], 6.22 [s, 1 H, H (pz)], 6.61 [d, J = 2.0 Hz, 2 H, H (im)], 7.14 [d, J = 2.0 Hz, 2 H, H (im)], 7.28 (m, 3 H, Ph), 7.77 (d, J = 7.4 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{\nu} = 3630$ vw (OH), 2503 m (BH) cm⁻¹.

6b: A solution of Zn(ClO₄)₂·6H₂O (130 mg, 0.35 mmol) in methanol (20 mL) was added with stirring to a solution of L^2 (200 mg, 0.35 mmol) in dichloromethane (100 mL). After 30 min, powdered KOH (22 mg, 0.39 mmol) was added. After stirring for 20 h, the mixture was filtered and the filtrate reduced in vacuo to half its volume. Cooling to -30 °C for 3 days precipitated 96 mg (44%) of raw 6b as colourless powder, m.p. 220 °C (dec.) which started decomposing upon attempts of further purification. C28H25BN6OS2Zn (601.88): calcd. C 55.88, H 4.19, N 13.96, Zn 10.86; found C 54.68, H 4.05, N 13.57, Zn 10.71. ¹H NMR $(CDCl_3)$: $\delta = 2.29$ [s, 3 H, Me (pz)], 6.29 [s, 2 H, H (pz)], 6.73 [d, J = 2.4 Hz, 2 H, H (im)], 6.80 [d, J = 2.4 Hz, 2 H, H (im)], 7.30-7.51 (m, 13 H, Ar), 7.53 (d, J = 7.3 Hz, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 3650 \text{ w}$ (OH), 2499 m (BH) cm⁻¹.

7: A solution of L^5 (665 mg, 1.08 mmol) and anhydrous methanol (60 mL) was slowly added dropwise, with stirring, to a solution of Zn(ClO₄)₂·6H₂O (400 mg, 1.08 mmol) in anhydrous methanol (60 mL). After stirring for 1 h, the precipitate was filtered off and the filtrate reduced to 80 mL in vacuo. The resulting precipitate was also filtered off and the filtrate concentrated to dryness. The residue was extracted with dichloromethane (14 mL). After filtration, the extract was concentrated to dryness and the residue dried in vacuo. 438 mg (55%) of 7 remained as a colourless powder; m.p. 230 °C (dec.). $C_{30}H_{28}BCIN_6O_6S_2Zn$ (744.37): calcd. C 48.41, H 3.79, N 11.29, S 8.62; found C 48.57, H 4.30, N 11.28, S 8.43. ¹H NMR (CDCl₃): $\delta = 2.28$ [s, 3 H, Me (pz)], 3.68 (s, 6 H, OMe), 6.49 [s, 1 H, H (pz)], 6.94–7.70 [m, 15 H, H (im), Ar], 7.93 [d, J = 7.2 Hz, 2 H, Ph] ppm. IR (KBr): $\tilde{v} = 2476$ w (BH), 1179 vs, 1093 vs (ClO₄) cm⁻¹.

8: A solution of L⁸ (309 mg, 0.50 mmol) in anhydrous ethanol (25 mL) was slowly added dropwise, with stirring, to a solution of Zn(ClO₄)₂·6H₂O (186 mg, 0.50 mmol) in anhydrous ethanol (25 mL) and the mixture stirred for 15 h. After filtration, the volume was reduced to one half in vacuo and the resulting precipitate filtered off. Recrystallisation from anhydrous ethanol yielded 313 mg (71%) of **8** as colourless crystals, m.p. 200 °C (dec.). $C_{32}H_{34}BClN_6O_7S_2Zn\cdotC_2H_5OH$ (790.44 + 46.07): calcd. C 48.82, H 4.82, N 10.05, S 7.67; found C 47.75, H 4.87, N 9.74, S 7.33. ¹H NMR (CDCl₃): $\delta = 1.18$ [t, J = 7.0 Hz, 6 H, CH₃ (Et)], 2.50 [s, 3

Table 1. Crystallogaphic details

	1c	1d	2a	3a	3b	3c
Empirical formula	C ₃₀ H ₂₈ BClN ₆ S ₂ Zn	C ₃₂ H ₃₂ BClN ₆ S ₂ Zn · 2CH ₃ CN	C ₃₀ H ₂₈ BN ₇ O ₃ S ₂ Zn	$C_{36}H_{40}B_2N_{12}S_4Zn \cdot 3H_2O$	$C_{60}H_{56}B_2N_{12}S_4Zn \cdot CH_2CN\cdot 2H_2O$	$C_{60}H_{56}B_2N_6S_4Zn \cdot H_2O$
Molecular mass	648.33	758.49	674.89	910.11	1237.54	1178.47
Crystal size [mm]	$0.15\times0.2\times0.35$	0.4 imes 0.3 imes 0.3	0.1 imes 0.1 imes 0.2	0.4 imes 0.3 imes 0.2	0.05 imes 0.1 imes 0.2	0.1 imes 0.1 imes 0.3
Space group	$P2_1/c$	$P\overline{1}$	$P2_1/n$	$P\overline{1}$	$P\overline{1}$	$P\bar{1}$
Z	4	2	4	2	2	2
a [A]	11.322(2)	10.087(2)	10.243(1)	8.98(2)	11.234(2)	10.788(1)
b [A]	14.625(2)	11.908(2)	29.169(3)	15.07(3)	17.060(2)	15.874(1)
	18.481(2)	17.035(3)	11.475(1)	17.55(4)	17.377(2)	19.024(2)
	90	102.41(3)	90	//.35(3)	/5.29/(3)	106.094(2)
p []	93.349(3)	95.54(5)	115.550(5)	80.10.(5) 78.80(2)	81.242(5) 87.280(2)	104.902(2) 105.176(2)
γ [] V [\mathbf{A}^{3}]	90 3045 9(7)	1845.0(6)	3147.8(6)	70.00(3) 2252 $A(8)$	3183 7(8)	103.170(2) 2822 5(4)
$d(calcd)$ [$\sigma \cdot cm^{-3}$]	1 41	1 37	1 42	1 34	1 55	1 37
$\mu(Mo-K_a) [mm^{-1}]$	1.06	0.89	0.96	0.78	1.14	0.64
hkl range	h: -15 to 9	h: -12 to 0	h: -13 to 5	h: -11 to 0	h: -14 to 14	h: -13 to 14
	k: -19 to 19	k: - 13 to 14	k: −38 to 38	k: −18 to 18	k: -22 to 22	<i>k</i> : −20 to 21
	<i>l</i> : -24 to 24	<i>l</i> : -20 to 20	<i>l</i> : -15 to 14	<i>l</i> : -21 to 21	<i>l</i> : -23 to 23	<i>l</i> : -25 to 24
Measured reflections	19180	7657	20120	9427	29078	25673
Independent reflections	7362	7230	7680	8831	14881	13177
Observed refl. $[I > 2\sigma(I)]$	5294	5589	2749	5204	5199	6271
Parameters	370	448	398	523	746	712
Refined reflections	7362	7230	7680	8831	14881	13177
R_1 (obsd. refl.)	0.035	0.046	0.072	0.066	0.079	0.056
WR_2 (all refl.)	0.113	0.132	0.256	0.238	0.282	0.145
[e/Å ³]	+0.3/-0.4	+1.3/-0.0	+0.8/-0.8	+1.3/-0.9	+1.1/-0.0	+0.77-0.7
	3d	3f	4a	4b	5	8
Empirical formula	3d C ₆₄ H ₆₄ B ₂ N ₁₂ S ₄ Zn• 8C ₆ H ₆	3f $C_{44}H_{72}B_2N_{12}S_4Zn$	4a C ₄₀ H ₃₈ BClN ₈ O ₄ S ₃ Zn• CH ₂ Cl ₂	4b C ₄₆ H ₅₀ BClN ₈ O ₄ S ₃ Zn• CH ₂ Cl ₂	5 C ₂₄ H ₃₀ N ₁₂ OS ₆ Zn ₄ · 6H ₂ O	8 C ₃₂ H ₃₄ BClN ₆ O ₇ S ₂ Zn• 2C ₂ H ₅ OH
Empirical formula Molecular mass	3d C ₆₄ H ₆₄ B ₂ N ₁₂ S ₄ Zn• 8C ₆ H ₆ 1841.47	3f C ₄₄ H ₇₂ B ₂ N ₁₂ S ₄ Zn 984.37	4a C ₄₀ H ₃₈ BClN ₈ O ₄ S ₃ Zn• CH ₂ Cl ₂ 987.52	4b C ₄₆ H ₅₀ BClN ₈ O ₄ S ₃ Zn• CH ₂ Cl ₂ 1071.68	5 C ₂₄ H ₃₀ N ₁₂ OS ₆ Zn ₄ · 6H ₂ O 1064.56	8 C ₃₂ H ₃₄ BClN ₆ O ₇ S ₂ Zn• 2C ₂ H ₅ OH 882.58
Empirical formula Molecular mass Crystal size [mm]	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn \\ 8C_6H_6 \\ 1841.47 \\ 0.2\times0.25\times0.35 \end{array}$	$\begin{array}{c} \textbf{3f} \\ \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ \\ \textbf{984.37} \\ 0.5\times0.4\times0.4 \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn^{\bullet} \\ CH_2Cl_2 \\ 987.52 \\ 0.15\times0.15\times0.25 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline \\ C_{46}H_{50}BClN_8O_4S_3Zn^{\bullet} \\ CH_2Cl_2 \\ 1071.68 \\ 0.1\times0.05\times0.05 \end{array}$	$\begin{array}{c} 5 \\ \\ \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{O}\mathbf{S}_{6}\mathbf{Z}\mathbf{n}_{4}\textbf{\cdot} \\ 6\mathbf{H}_{2}\mathbf{O} \\ 1064.56 \\ 0.2\times0.1\times0.1 \end{array}$	$\begin{array}{c} 8 \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn^{\star}\\ 2C_2H_5OH\\ 882.58\\ 0.25\times 0.15\times 0.1 \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn \\ 8C_6H_6 \\ 1841.47 \\ 0.2 \times 0.25 \times 0.35 \\ C2/c \end{array}$	$\begin{array}{c} \textbf{3f} \\ \\ \hline C_{44} H_{72} B_2 N_{12} S_4 Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn^{\bullet} \\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline \\ C_{46}H_{50}BClN_8O_4S_3Zn^{\bullet} \\ CH_2Cl_2 \\ 1071.68 \\ 0.1\times0.05\times0.05 \\ P\bar{1} \end{array}$	$\begin{array}{c} 5 \\ \\ \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{O}\mathbf{S}_{6}\mathbf{Z}\mathbf{n}_{4}\textbf{\cdot} \\ 6\mathbf{H}_{2}\mathbf{O} \\ 1064.56 \\ 0.2\times0.1\times0.1 \\ R\bar{3} \end{array}$	$\begin{array}{c} {\bf 8} \\ \hline {\bf C}_{32}{\bf H}_{34}{\bf B}{\bf C}{\bf 1}{\bf N}_6{\bf O}_7{\bf S}_2{\bf Z}{\bf n}^*\\ {\bf 2}{\bf C}_2{\bf H}_5{\bf O}{\bf H}\\ {\bf 882.58}\\ {\bf 0.25\times 0.15\times 0.1}\\ P\bar{\bf 1} \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn \\ 8C_6H_6 \\ 1841.47 \\ 0.2 \times 0.25 \times 0.35 \\ C2/c \\ 4 \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^{\bullet}\\ CH_2Cl_2\\ 1071.68\\ 0.1\times0.05\times0.05\\ P\bar{1}\\ 2 \end{array}$	$\begin{array}{c} 5 \\ \\ \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{O}\mathbf{S}_{6}\mathbf{Z}\mathbf{n}_{4}^{\star} \\ 6\mathbf{H}_{2}\mathbf{O} \\ 1064.56 \\ 0.2\times0.1\times0.1 \\ R\overline{3} \\ 6 \end{array}$	$\begin{array}{c} 8 \\ \hline \mathbf{C}_{32}\mathbf{H}_{34}\mathbf{B}\mathbf{C}\mathbf{1N}_{6}\mathbf{O}_{7}\mathbf{S}_{2}\mathbf{Z}\mathbf{n}^{*}\\ \mathbf{2C}_{2}\mathbf{H}_{5}\mathbf{O}\mathbf{H}\\ 882.58\\ 0.25\times0.15\times0.1\\ P\bar{1}\\ 2 \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a $\begin{bmatrix} A \\ A \end{bmatrix}$	3d $C_{64}H_{64}B_2N_{12}S_4Zn$ $8C_6H_6$ 1841.47 $0.2 \times 0.25 \times 0.35$ C2/c 4 26.507(3) 80.607(3)	3f $C_{44}H_{72}B_2N_{12}S_4Zn$ 984.37 $0.5 \times 0.4 \times 0.4$ $P\bar{1}$ 2 10.576(1) 16.00(2)	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn^{\bullet}\\ CH_2Cl_2\\ 987.52\\ 0.15\times 0.15\times 0.25\\ P2_1/c\\ 4\\ 11.399(1)\\ \cdots \cdots \cdots \cdots \cdots \\ \textbf{4} \end{array}$	4b $C_{46}H_{50}BClN_8O_4S_3Zn$ CH_2Cl_2 1071.68 $0.1 \times 0.05 \times 0.05$ $P\bar{I}$ 2 10.897(1) 15.6897(1)	5 $C_{24}H_{30}N_{12}OS_6Zn_4 \cdot 6H_2O$ 1064.56 $0.2 \times 0.1 \times 0.1$ R_3 6 15.993(2) 16.000000000000000000000000000000000000	$\begin{array}{c} 8 \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn \\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ PI \\ 2 \\ 10.962(1) \\ 14.102(0) \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] b [Å]	3d $C_{64}H_{64}B_2N_{12}S_4Zn$ $8C_6H_6$ 1841.47 $0.2 \times 0.25 \times 0.35$ C2/c 4 26.507(3) 20.022(2) 20.014(2)	3f $C_{44}H_{72}B_2N_{12}S_4Zn$ 984.37 0.5 × 0.4 × 0.4 $P\bar{1}$ 2 10.576(1) 16.243(1) 16.2043(1)	4a $C_{40}H_{38}BCIN_8O_4S_3Zn$ CH_2Cl_2 987.52 $0.15 \times 0.15 \times 0.25$ $P2_1/c$ 4 11.399(1) 11.133(1) 25 (51(2))	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 1071.68 \\ 0.1 \times 0.05 \times 0.05 \\ P\bar{I} \\ 2 \\ 10.897(1) \\ 15.635(2) \\ 17.12(2) \end{array}$	5 $C_{24}H_{30}N_{12}OS_6Zn_4 \cdot 6H_2O$ 1064.56 $0.2 \times 0.1 \times 0.1$ R_3^3 6 15.993(2) 15.993(2) 20.715(6)	$\begin{array}{c} 8 \\ \hline \mathbf{C}_{32}\mathbf{H}_{34}\mathbf{B}\mathbf{C}\mathbf{IN}_{6}\mathbf{O}_{7}\mathbf{S}_{2}\mathbf{Z}\mathbf{n} \\ \mathbf{2C}_{2}\mathbf{H}_{5}\mathbf{OH} \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14, 131(2) \\ 14, 92(2) \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] b [Å] c [Å] c [Å]	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^{*}\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90 \end{array}$	3f $C_{44}H_{72}B_2N_{12}S_4Zn$ 984.37 $0.5 \times 0.4 \times 0.4$ $P\bar{1}$ 2 10.576(1) 16.243(1) 16.809(1) 105.25(2)	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn+\\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \\ 11.399(1) \\ 11.133(1) \\ 35.651(3) \\ 90 \end{array}$	4b $C_{46}H_{50}BClN_8O_4S_3Zn$ CH_2Cl_2 1071.68 $0.1 \times 0.05 \times 0.05$ $P\bar{1}$ 2 10.897(1) 15.635(2) 17.131(2) 66.421(2)	$\begin{array}{c} 5 \\ \\ \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{OS}_{6}\mathbf{Zn}_{4}\boldsymbol{\cdot} \\ \mathbf{6H}_{2}\mathbf{O} \\ 1064, 56 \\ 0.2\times0.1\times0.1 \\ \mathbf{R}_{3}^{3} \\ 6 \\ 15.993(2) \\ 15.993(2) \\ 33.715(5) \\ 90 \end{array}$	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn^{\text{.}}\\ 2C_2H_5OH\\ 882.58\\ 0.25\times0.15\times0.1\\ P\bar{1}\\ 2\\ 10.962(1)\\ 14.131(2)\\ 14.834(2)\\ 101.011(2)\\ \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] c [Å] a [°] a [°]	3d $C_{64}H_{64}B_2N_{12}S_4Zn$ $8C_6H_6$ 1841.47 $0.2 \times 0.25 \times 0.35$ C2/c 4 26.507(3) 20.022(2) 20.914(3) 90 112.828(2)	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \end{array}$	4a $C_{40}H_{38}BCIN_8O_4S_3Zn$ - CH_2Cl_2 987.52 $0.15 \times 0.15 \times 0.25$ $P2_1/c$ 4 11.399(1) 11.133(1) 35.651(3) 90 97.979(2)	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^4\\ CH_2Cl_2\\ 1071.68\\ 0.1\times0.05\times0.05\\ P\bar{1}\\ 2\\ 10.897(1)\\ 15.635(2)\\ 17.131(2)\\ 66.431(2)\\ 74.600(2)\\ \end{array}$	$\begin{array}{c} 5 \\ \\ \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{OS}_{6}\mathbf{Zn}_{4}\boldsymbol{\cdot} \\ \mathbf{6H}_{2}\mathbf{O} \\ 1064, 56 \\ 0.2\times0.1\times0.1 \\ \mathbf{R}_{3}^{3} \\ 6 \\ 15.993(2) \\ 15.993(2) \\ 33.715(5) \\ 90 \\ 90 \end{array}$	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn \\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 100.58(2) \\ \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°]	$\begin{array}{c} \textbf{3d} \\\\\hline C_{64}H_{64}B_2N_{12}S_4Zn^{*}\\ 8C_6H_6\\ 1841.47\\ 0.2\times0.25\times0.35\\ \textit{C2/c}\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90 \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \\ 11.399(1) \\ 11.133(1) \\ 35.651(3) \\ 90 \\ 97.978(2) \\ 90 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^{\bullet}\\ CH_2Cl_2\\ 1071.68\\ 0.1\times0.05\times0.05\\ P\bar{1}\\ 2\\ 10.897(1)\\ 15.635(2)\\ 17.131(2)\\ 66.431(2)\\ 74.600(2)\\ 80.263(3)\\ \end{array}$	$\begin{array}{c} 5 \\ & \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{OS}_{6}\mathbf{Zn}_{4}\boldsymbol{\cdot} \\ & \mathbf{6H}_{2}\mathbf{O} \\ & 1064.56 \\ & 0.2\times0.1\times0.1 \\ & R\overline{3} \\ & 6 \\ & 15.993(2) \\ & 15.993(2) \\ & 15.993(2) \\ & 33.715(5) \\ & 90 \\ & 90 \\ & 120 \end{array}$	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn^{\bullet}\\ 2C_2H_5OH\\ 882.58\\ 0.25\times0.15\times0.1\\ P\bar{1}\\ 2\\ 10.962(1)\\ 14.131(2)\\ 14.834(2)\\ 101.011(2)\\ 109.568(2)\\ 94.234(2) \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] $\beta [°]$ $\gamma [°]$ $V [Å^3]$	3d $C_{64}H_{64}B_2N_{12}S_4Zn$ $8C_6H_6$ 1841.47 $0.2 \times 0.25 \times 0.35$ C2/c 4 26.507(3) 20.022(2) 20.914(3) 90 112.828(2) 90 10229 $9(18)$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline C_{40}H_{38}BClN_8O_4S_3Zn^4\\ CH_2Cl_2\\ 987.52\\ 0.15\times0.15\times0.25\\ P2_1/c\\ 4\\ 11.399(1)\\ 11.133(1)\\ 35.651(3)\\ 90\\ 97.978(2)\\ 90\\ 4480\ 6(6) \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^{\bullet}\\ CH_2Cl_2\\ 1071.68\\ 0.1\times0.05\times0.05\\ P\bar{1}\\ 2\\ 10.897(1)\\ 15.635(2)\\ 17.131(2)\\ 66.431(2)\\ 74.600(2)\\ 80.263(3)\\ 2572.6(5) \end{array}$	5 $C_{24}H_{30}N_{12}OS_6Zn_4$ · $6H_2O$ 1064.56 $0.2 \times 0.1 \times 0.1$ R_3^3 6 15.993(2) 15.993(2) 33.715(5) 90 90 120 7468.0(15)	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn^{\bullet}\\ 2C_2H_5OH\\ 882.58\\ 0.25\times0.15\times0.1\\ P\bar{1}\\ 2\\ 10.962(1)\\ 14.131(2)\\ 14.834(2)\\ 101.011(2)\\ 109.568(2)\\ 94.234(2)\\ 2101.6(4) \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] c [Å] $\alpha [°]$ $\beta [°]$ $\gamma [°]$ $\gamma [°]$ $V [Å^3]$ d(calcd.) [gcm-3]	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^{*}\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ \textit{C2/c}\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline C_{40}H_{38}BClN_8O_4S_3Zn^4\\ CH_2Cl_2\\ 987.52\\ 0.15\times 0.15\times 0.25\\ P2_1/c\\ 4\\ 11.399(1)\\ 11.133(1)\\ 35.651(3)\\ 90\\ 97.978(2)\\ 90\\ 4480.6(6)\\ 1.46 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^{*}\\ CH_2Cl_2 \\ 1071.68 \\ 0.1 \times 0.05 \times 0.05 \\ P\bar{1} \\ 2 \\ 10.897(1) \\ 15.635(2) \\ 17.131(2) \\ 66.431(2) \\ 74.600(2) \\ 80.263(3) \\ 2572.6(5) \\ 1.38 \end{array}$	$\begin{array}{c} {\bf 5} \\ \\ {\bf C}_{24}{\bf H}_{30}{\bf N}_{12}{\bf OS}_6{\bf Zn}_4 \cdot \\ {\bf 6H}_2{\bf O} \\ {\bf 1064,56} \\ {\bf 0.2 \times 0.1 \times 0.1} \\ {\bf R}\bar{\bf 3} \\ {\bf 6} \\ {\bf 15.993(2)} \\ {\bf 15.993(2)} \\ {\bf 33.715(5)} \\ {\bf 90} \\ {\bf 90} \\ {\bf 120} \\ {\bf 7468.0(15)} \\ {\bf 1.40} \end{array}$	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn^{*}\\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] c [Å] a [°] $\beta [°]$ $\gamma [°]$ $V [A^3]$ d(calcd.) [gcm-3] $\mu(Mo-K_0) [mm-1]$	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2\times0.25\times0.35\\ \textit{C2/c}\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline C_{40}H_{38}BClN_8O_4S_3Zn^4\\ CH_2Cl_2\\ 987.52\\ 0.15\times0.15\times0.25\\ P2_1/c\\ 4\\ 11.399(1)\\ 11.133(1)\\ 35.651(3)\\ 90\\ 97.978(2)\\ 90\\ 4480.6(6)\\ 1.46\\ 0.92\\ \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline \\ C_{46}H_{50}BClN_8O_4S_3Zn^{*}\\ CH_2Cl_2 \\ 1071.68 \\ 0.1 \times 0.05 \times 0.05 \\ P\bar{1} \\ 2 \\ 10.897(1) \\ 15.635(2) \\ 17.131(2) \\ 66.431(2) \\ 74.600(2) \\ 80.263(3) \\ 2572.6(5) \\ 1.38 \\ 0.81 \\ \end{array}$	$\begin{array}{c} 5 \\ \\ \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{OS}_{6}\mathbf{Zn}_{4} \\ \mathbf{6H}_{2}\mathbf{O} \\ 1064.56 \\ 0.2 \times 0.1 \times 0.1 \\ R\bar{3} \\ 6 \\ 15.993(2) \\ 15.993(2) \\ 33.715(5) \\ 90 \\ 90 \\ 120 \\ 7468.0(15) \\ 1.40 \\ 2.20 \end{array}$	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn^{*}\\ 2C_2H_5OH\\ 882.58\\ 0.25\times0.15\times0.1\\ P\bar{1}\\ 2\\ 10.962(1)\\ 14.131(2)\\ 14.834(2)\\ 101.011(2)\\ 109.568(2)\\ 94.234(2)\\ 2101.6(4)\\ 1.40\\ 0.81\\ \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å] d(calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35 \text{ to } 33 \end{array}$	$\begin{array}{c} \textbf{3f} \\ \\ \hline C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline C_{40}H_{38}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \\ 11.399(1) \\ 11.133(1) \\ 35.651(3) \\ 90 \\ 97.978(2) \\ 90 \\ 4480.6(6) \\ 1.46 \\ 0.92 \\ h: -15 \text{ to } 6 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 1071.68 \\ 0.1 \times 0.05 \times 0.05 \\ P\overline{l} \\ 2 \\ 10.897(1) \\ 15.635(2) \\ 17.131(2) \\ 66.431(2) \\ 74.600(2) \\ 80.263(3) \\ 2572.6(5) \\ 1.38 \\ 0.81 \\ h: -14 \text{ to } 14 \end{array}$	$\begin{array}{c} 5 \\ & \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{OS}_{6}\mathbf{Zn}_{4} \cdot \\ \mathbf{6H}_{2}\mathbf{O} \\ & 1064.56 \\ & 0.2 \times 0.1 \times 0.1 \\ \mathbf{R3} \\ & 6 \\ & 15.993(2) \\ & 15.993(2) \\ & 33.715(5) \\ & 90 \\ & 90 \\ & 120 \\ & 7468.0(15) \\ & 1.40 \\ & 2.20 \\ & h: -19 \text{ to } 19 \end{array}$	$\begin{array}{c} {\bf 8} \\ \\ {\bf C}_{32}{\bf H}_{34}{\bf B}{\bf C}{\bf 1}{\bf N}_6{\bf O}_7{\bf S}_2{\bf Z}{\bf n}\cdot\\ {\bf 2C}_2{\bf H}_5{\bf O}{\bf H}\\ {\bf 882.58}\\ {\bf 0.25\times 0.15\times 0.1}\\ P\bar{\bf I}\\ {\bf 2}\\ {\bf 10.962(1)}\\ {\bf 14.131(2)}\\ {\bf 14.834(2)}\\ {\bf 101.011(2)}\\ {\bf 109.568(2)}\\ {\bf 94.234(2)}\\ {\bf 2101.6(4)}\\ {\bf 1.40}\\ {\bf 0.81}\\ h: -{\bf 14}\ to\ {\bf 14} \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å] d(calcd.) [gcm ⁻³] μ (Mo- K_{α}) [mm ⁻¹] hkl range	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35 \text{ to } 33\\ k: -26 \text{ to } 19 \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\overline{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.243(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \\ k: -21 \text{ to } 21 \\ \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline C_{40}H_{38}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \\ 11.399(1) \\ 11.133(1) \\ 35.651(3) \\ 90 \\ 97.978(2) \\ 90 \\ 4480.6(6) \\ 1.46 \\ 0.92 \\ h: -15 \text{ to } 6 \\ k: -14 \text{ to } 13 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 1071.68 \\ 0.1 \times 0.05 \times 0.05 \\ PI \\ 2 \\ 10.897(1) \\ 15.635(2) \\ 17.131(2) \\ 66.431(2) \\ 74.600(2) \\ 80.263(3) \\ 2572.6(5) \\ 1.38 \\ 0.81 \\ h: -14 \text{ to } 14 \\ k: -20 \text{ to } 20 \end{array}$	$\begin{array}{c} 5 \\ & \mathbf{C}_{24}\mathbf{H}_{30}\mathbf{N}_{12}\mathbf{OS}_{6}\mathbf{Zn}_{4} \cdot \\ \mathbf{6H}_{2}\mathbf{O} \\ & 1064.56 \\ & 0.2 \times 0.1 \times 0.1 \\ & \mathbf{R3} \\ & 6 \\ & 15.993(2) \\ & 15.993(2) \\ & 15.993(2) \\ & 33.715(5) \\ & 90 \\ & 90 \\ & 120 \\ & 7468.0(15) \\ & 1.40 \\ & 2.20 \\ & h: -19 \text{ to } 19 \\ & k: -19 \text{ to } 19 \end{array}$	$\begin{array}{c} \textbf{8} \\ \hline \\ C_{32}H_{34}BClN_6O_7S_2Zn\cdot\\ 2C_2H_5OH\\ 882.58\\ 0.25\times0.15\times0.1\\ P\bar{1}\\ 2\\ 10.962(1)\\ 14.131(2)\\ 14.834(2)\\ 101.011(2)\\ 109.568(2)\\ 94.234(2)\\ 2101.6(4)\\ 1.40\\ 0.81\\ h: -14 \text{ to } 14\\ k: -18 \text{ to } 18 \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å] d(calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2\times0.25\times0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35\ to\ 33\\ k: -26\ to\ 19\\ l: -27\ to\ 27\\ \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\overline{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.243(1) \\ 105.253(2) \\ 101.468(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \\ h: -12 \text{ to } 13 \\ h: -21 \text{ to } 21 \\ l: -22 \text{ to } 22 \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \\ 11.399(1) \\ 11.133(1) \\ 35.651(3) \\ 90 \\ 97.978(2) \\ 90 \\ 4480.6(6) \\ 1.46 \\ 0.92 \\ h: -15 \text{ to } 6 \\ k: -14 \text{ to } 13 \\ l: -47 \text{ to } 44 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^{\bullet}\\ CH_2Cl_2\\ 1071.68\\ 0.1\times 0.05\times 0.05\\ P\overline{l}\\ 2\\ 10.897(1)\\ 15.635(2)\\ 17.131(2)\\ 66.431(2)\\ 74.600(2)\\ 80.263(3)\\ 2572.6(5)\\ 1.38\\ 0.81\\ h: -14\ to\ 14\\ k: -20\ to\ 20\\ l: -22\ to\ 22 \end{array}$		$\begin{array}{c} \textbf{8} \\ \hline \\ C_{32}H_{34}BClN_6O_7S_2Zn \\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \\ 0.81 \\ h: -14 \text{ to } 14 \\ h: -14 \text{ to } 14 \\ h: -18 \text{ to } 18 \\ l: -19 \text{ to } 19 \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å ³] d(calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35\ to\ 33\\ k: -26\ to\ 19\\ l:\ -27\ to\ 27\\ 32298\\ \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \\ k: -21 \text{ to } 21 \\ k: -21 \text{ to } 21 \\ k: -22 \text{ to } 22 \\ 23401 \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn \\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \\ 11.399(1) \\ 11.133(1) \\ 35.651(3) \\ 90 \\ 97.978(2) \\ 90 \\ 4480.6(6) \\ 1.46 \\ 0.92 \\ h: -15 \text{ to } 6 \\ k: -14 \text{ to } 13 \\ l: -47 \text{ to } 44 \\ 27857 \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^4\\ CH_2Cl_2\\ 1071.68\\ 0.1\times0.05\times0.05\\ P\bar{1}\\ 2\\ 10.897(1)\\ 15.635(2)\\ 17.131(2)\\ 66.431(2)\\ 74.600(2)\\ 80.263(3)\\ 2572.6(5)\\ 1.38\\ 0.81\\ h: -14\ to\ 14\\ k: -20\ to\ 20\\ l: -22\ to\ 22\\ 23909 \end{array}$		$\begin{array}{c} {\bf 8} \\ \hline \\ C_{32}H_{34}BClN_6O_7S_2Zn \\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \\ 0.81 \\ h: -14 \text{ to } 14 \\ h: -18 \text{ to } 18 \\ h: -19 \text{ to } 19 \\ 19203 \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å] d (calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range Measured reflections Independent reflections	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35\ to\ 33\\ k: -26\ to\ 19\\ k: -27\ to\ 27\\ 32298\\ 12428\\ \end{array}$	3f $C_{44}H_{72}B_2N_{12}S_4Zn$ 984.37 0.5 × 0.4 × 0.4 $P\overline{1}$ 2 10.576(1) 16.243(1) 16.809(1) 105.253(2) 101.468(2) 104.034(2) 2594.2(4) 1.26 0.68 h: -12 to 13 k: -21 to 21 h: -22 to 22 23401 11738	$\begin{array}{c} \textbf{4a} \\ \hline C_{40}H_{38}BClN_8O_4S_3Zn+CH_2Cl_2\\ 987.52\\ 0.15\times0.15\times0.25\\ P2_1/c\\ \textbf{4} \\ 11.399(1)\\ 11.133(1)\\ 35.651(3)\\ 90\\ 97.978(2)\\ 90\\ \textbf{4480.6(6)}\\ 1.46\\ 0.92\\ h: -15\ to\ 6\\ k:\ -14\ to\ 13\\ l:\ -47\ to\ 44\\ 27857\\ 10752\\ \end{array}$	$\begin{array}{c} \textbf{4b} \\ \hline C_{46}H_{50}BClN_8O_4S_3Zn^4\\ CH_2Cl_2\\ 1071.68\\ 0.1\times0.05\times0.05\\ P\bar{1}\\ 2\\ 10.897(1)\\ 15.635(2)\\ 17.131(2)\\ 66.431(2)\\ 74.600(2)\\ 80.263(3)\\ 2572.6(5)\\ 1.38\\ 0.81\\ h: -14 \text{ to } 14\\ k: -20 \text{ to } 20\\ l: -22 \text{ to } 22\\ 23909\\ 12173\\ \end{array}$		$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn \\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \\ 0.81 \\ h: -14 \text{ to } 14 \\ k: -18 \text{ to } 18 \\ h: -19 \text{ to } 19 \\ 19203 \\ 9895 \\ \hline \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] $\beta [°]$ $\gamma [°]$ $V [Å^3]$ $d(calcd.) [gcm^{-3}]$ $\mu(Mo-K_a) [mm^{-1}]$ hkl range Measured reflections Independent reflections Observed refl. $[I > 2\sigma(I)]$	3d $C_{64}H_{64}B_2N_{12}S_4Zn$ $8C_6H_6$ 1841.47 $0.2 \times 0.25 \times 0.35$ C2/c 4 26.507(3) 20.022(2) 20.914(3) 90 10229.9(18) 1.20 0.38 h: -35 to 33 k: -26 to 19 l: -27 to 27 32298 12428 4317 56	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.243(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \\ k: -21 \text{ to } 21 \\ h: -22 \text{ to } 22 \\ 23401 \\ 11738 \\ 6604 \\ \hline \end{array}$	$\begin{array}{c} \textbf{4a} \\ \hline \\ C_{40}H_{38}BClN_8O_4S_3Zn+\\ CH_2Cl_2 \\ 987.52 \\ 0.15 \times 0.15 \times 0.25 \\ P2_1/c \\ 4 \\ 11.399(1) \\ 11.133(1) \\ 35.651(3) \\ 90 \\ 97.978(2) \\ 90 \\ 4480.6(6) \\ 1.46 \\ 0.92 \\ h: -15 \text{ to } 6 \\ k: -14 \text{ to } 13 \\ l: -47 \text{ to } 44 \\ 27857 \\ 10752 \\ 5145 \\ \hline \end{array}$	4b C ₄₆ H ₅₀ BClN ₈ O ₄ S ₃ Zn· CH ₂ Cl ₂ 1071.68 0.1 × 0.05 × 0.05 $P\bar{1}$ 2 10.897(1) 15.635(2) 17.131(2) 66.431(2) 74.600(2) 80.263(3) 2572.6(5) 1.38 0.81 h: -14 to 14 k: -20 to 20 l: -22 to 22 23909 12173 2730 (25)	5 $C_{24}H_{30}N_{12}OS_6Zn_4$ · $6H_2O$ 1064.56 $0.2 \times 0.1 \times 0.1$ R_3^3 6 15.993(2) 15.993(2) 33.715(5) 90 120 7468.0(15) 1.40 2.20 h: -19 to $19h: -19$ to $19h: -19$ to $19h: -11$ to 411960132192134	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn \\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \\ 0.81 \\ h: -14 \text{ to } 14 \\ k: -18 \text{ to } 18 \\ h: -19 \text{ to } 19 \\ 19203 \\ 9895 \\ 5057 \\ \hline \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å ³] d(calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range Measured reflections Independent reflections Observed refl. [$I > 2\sigma(I)$] Parameters D $= 2\sigma(I)$	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35 \text{ to } 33\\ k: -26 \text{ to } 19\\ l: -27 \text{ to } 27\\ 32298\\ 12428\\ 4317\\ 596\\ 12420\\ \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\overline{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \\ k: -21 \text{ to } 21 \\ l: -22 \text{ to } 22 \\ 23401 \\ 11738 \\ 6604 \\ 576 \\ 11200 \end{array}$	4a $C_{40}H_{38}BCIN_8O_4S_3Zn$ · CH_2Cl_2 987.52 $0.15 \times 0.15 \times 0.25$ $P2_1/c$ 4 11.399(1) 11.133(1) 35.651(3) 90 97.978(2) 90 4480.6(6) 1.46 0.92 h: -15 to 6 k: -14 to 13 l: -47 to 44 27857 10752 5145 578 10752	4b C ₄₆ H ₅₀ BClN ₈ O ₄ S ₃ Zn· CH ₂ Cl ₂ 1071.68 0.1 × 0.05 × 0.05 $P\bar{1}$ 2 10.897(1) 15.635(2) 17.131(2) 66.431(2) 74.600(2) 80.263(3) 2572.6(5) 1.38 0.81 h: -14 to 14 k: -20 to 20 l: -22 to 22 23909 12173 2730 605		$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn \\ 2C_2H_5OH \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \\ 0.81 \\ h: -14 \text{ to } 14 \\ k: -18 \text{ to } 18 \\ h: -19 \text{ to } 19 \\ 19203 \\ 9895 \\ 5057 \\ 509 \\ 0005 \\ \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å ³] d(calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range Measured reflections Independent reflections Observed refl. [$I > 2\sigma(I)$] Parameters Refined reflections D	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35 \text{ to } 33\\ h: -35 \text{ to } 33\\ h: -26 \text{ to } 19\\ l: -27 \text{ to } 27\\ 32298\\ 12428\\ 4317\\ 596\\ 12428\\ 0.060\\ \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \\ k: -21 \text{ to } 21 \\ l: -22 \text{ to } 22 \\ 23401 \\ 11738 \\ 6604 \\ 576 \\ 11738 \\ 0.079 \end{array}$	4a $C_{40}H_{38}BCIN_8O_4S_3Zn$ · CH_2Cl_2 987.52 0.15 × 0.15 × 0.25 $P2_1/c$ 4 11.399(1) 11.133(1) 35.651(3) 90 97.978(2) 90 4480.6(6) 1.46 0.92 h: -15 to 6 k: -14 to 13 l: -47 to 44 27857 10752 5145 578 10752 0.077	4b C ₄₆ H ₅₀ BClN ₈ O ₄ S ₃ Zn· CH ₂ Cl ₂ 1071.68 0.1 × 0.05 × 0.05 $P\bar{1}$ 2 10.897(1) 15.635(2) 17.131(2) 66.431(2) 74.600(2) 80.263(3) 2572.6(5) 1.38 0.81 h: -14 to 14 k: -20 to 20 l: -22 to 22 23909 12173 2730 605 12173 0.105	5 $C_{24}H_{30}N_{12}OS_6Zn_4$ · $6H_2O$ 1064.56 $0.2 \times 0.1 \times 0.1$ R_3 6 15.993(2) 15.993(2) 33.715(5) 90 90 120 7468.0(15) 1.40 2.20 h: -19 to $19k: -19$ to $19k: -19$ to $19k: -19$ to $19k: -110$ to $19k: -110$ to $1101000100010001000100010001000100010$	$\begin{array}{c} \textbf{8} \\ \hline C_{32}H_{34}BClN_6O_7S_2Zn^{*}\\ 2C_2H_5OH\\ 882.58\\ 0.25 \times 0.15 \times 0.1\\ P\bar{1}\\ 2\\ 10.962(1)\\ 14.131(2)\\ 14.834(2)\\ 101.011(2)\\ 109.568(2)\\ 94.234(2)\\ 2101.6(4)\\ 1.40\\ 0.81\\ h^{*} - 14 \text{ to } 14\\ k^{*}_{:} - 18 \text{ to } 18\\ h^{*}_{:} - 19 \text{ to } 19\\ 19203\\ 9895\\ 5057\\ 509\\ 9895\\ 0.061\\ \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å] d(calcd.) [gcm ⁻³] d(calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range Measured reflections Independent reflections Observed refl. [$I > 2\sigma(I)$] Parameters Refined reflections R_1 (obsd. refl.) w^R (α ll refl.)	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35 \text{ to } 33\\ k: -26 \text{ to } 19\\ l: -27 \text{ to } 27\\ 32298\\ 12428\\ 4317\\ 596\\ 12428\\ 4317\\ 596\\ 12428\\ 0.069\\ 0.204\\ \end{array}$	$\begin{array}{c} \textbf{3f} \\ \hline \\ C_{44}H_{72}B_2N_{12}S_4Zn \\ 984.37 \\ 0.5 \times 0.4 \times 0.4 \\ P\bar{1} \\ 2 \\ 10.576(1) \\ 16.243(1) \\ 16.809(1) \\ 105.253(2) \\ 101.468(2) \\ 104.034(2) \\ 2594.2(4) \\ 1.26 \\ 0.68 \\ h: -12 \text{ to } 13 \\ k: -21 \text{ to } 21 \\ l: -22 \text{ to } 22 \\ 23401 \\ 11738 \\ 6604 \\ 576 \\ 11738 \\ 0.078 \\ 0.230 \end{array}$	4a $C_{40}H_{38}BCIN_8O_4S_3Zn^4$ CH_2Cl_2 987.52 $0.15 \times 0.15 \times 0.25$ $P2_1/c$ 4 $11.399(1)$ $11.133(1)$ $35.651(3)$ 90 $97.978(2)$ 90 $4480.6(6)$ 1.46 0.92 $h: -15$ to 6 $k: -14$ to 13 $l: -47$ to 44 27857 10752 5145 578 10752 0.077 0.264	4b C ₄₆ H ₅₀ BClN ₈ O ₄ S ₃ Zn· CH ₂ Cl ₂ 1071.68 0.1 × 0.05 × 0.05 $P\overline{l}$ 2 10.897(1) 15.635(2) 17.131(2) 66.431(2) 74.600(2) 80.263(3) 2572.6(5) 1.38 0.81 h: -14 to 14 k: -20 to 20 l: -22 to 22 23909 12173 2730 605 12173 0.105 0.387	5 $C_{24}H_{30}N_{12}OS_6Zn_4 \cdot 6H_2O$ 1064.56 $0.2 \times 0.1 \times 0.1$ R_3 6 15.993(2) 15.993(2) 33.715(5) 90 90 120 7468.0(15) 1.40 2.20 h: -19 to $19k: -19$ to $19k: -19$ to $19k: -119$ to $19k: -110$ to $11032190.0430.132$	$\begin{array}{c} {\bf 8} \\ \hline C_{32} {\rm H}_{34} {\rm BClN}_6 {\rm O}_7 {\rm S}_2 {\rm Zn} \cdot \\ {\rm 2C}_3 {\rm H}_5 {\rm OH} \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P\bar{1} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \\ 0.81 \\ h: -14 \ to \ 14 \\ k: -18 \ to \ 18 \\ h: -19 \ to \ 19 \\ 19203 \\ 9895 \\ 5057 \\ 509 \\ 9895 \\ 0.061 \\ 0.187 \\ \end{array}$
Empirical formula Molecular mass Crystal size [mm] Space group Z a [Å] b [Å] c [Å] a [°] β [°] γ [°] V [Å] d(calcd.) [gcm ⁻³] μ (Mo- K_a) [mm ⁻¹] hkl range Measured reflections Independent reflections Observed refl. [$I > 2\sigma(I)$] Parameters Refined reflections R_1 (obsd. refl.) wR_2 (all refl.) Residual electron density	$\begin{array}{c} \textbf{3d} \\ \hline \\ C_{64}H_{64}B_2N_{12}S_4Zn^*\\ 8C_6H_6\\ 1841.47\\ 0.2 \times 0.25 \times 0.35\\ C2/c\\ 4\\ 26.507(3)\\ 20.022(2)\\ 20.914(3)\\ 90\\ 112.828(2)\\ 90\\ 112.828(2)\\ 90\\ 10229.9(18)\\ 1.20\\ 0.38\\ h: -35 \text{ to } 33\\ k: -26 \text{ to } 19\\ l: -27 \text{ to } 27\\ 32298\\ 12428\\ 4317\\ 596\\ 12428\\ 4317\\ 596\\ 12428\\ 0.069\\ 0.204\\ +1 2/-0 5 \end{array}$	3f $C_{44}H_{72}B_2N_{12}S_4Zn$ 984.37 0.5 × 0.4 × 0.4 $P\overline{1}$ 2 10.576(1) 16.243(1) 16.809(1) 105.253(2) 101.468(2) 104.034(2) 2594.2(4) 1.26 0.68 h: -12 to 13 k: -21 to 21 l: -22 to 22 23401 11738 6604 576 11738 0.078 0.230 +1.2/-0.8	4a $C_{40}H_{38}BCIN_8O_4S_3Zn^4$ CH_2Cl_2 987.52 $0.15 \times 0.15 \times 0.25$ $P2_1/c$ 4 $11.399(1)$ $11.133(1)$ $35.651(3)$ 90 $97.978(2)$ 90 448 1.46 0.92 $h: -15$ to 6 $k: -14$ to 13 $l: -47$ to 44 27857 10752 5145 578 10752 0.264 $+13/-11$	4b $C_{46}H_{50}BCIN_8O_4S_3Zn$ CH_2Cl_2 1071.68 $0.1 \times 0.05 \times 0.05$ PI 2 10.897(1) 15.635(2) 17.131(2) 66.431(2) 74.600(2) 80.263(3) 2572.6(5) 1.38 0.81 h: -14 to $14k: -20$ to $20l: -22$ to $2223909121732730605121730.1050.387+11/-0.8$	5 $C_{24}H_{30}N_{12}OS_6Zn_4 \cdot 6H_2O$ 1064.56 $0.2 \times 0.1 \times 0.1$ R_3 6 15.993(2) 15.993(2) 33.715(5) 90 90 120 7468.0(15) 1.40 2.20 h: -19 to $19k: -19$ to $19k: -19$ to $19k: -19$ to $19k: -110$ to $19k: -110$ to $19k: -110$ to $1103219213410632190.0430.132+13/-0.4$	$\begin{array}{c} {\bf 8} \\ \hline C_{32}{\rm H}_{34}{\rm BClN}_6{\rm O}_7{\rm S}_2{\rm Zn} \cdot \\ {\rm 2C}_2{\rm H}_5{\rm OH} \\ 882.58 \\ 0.25 \times 0.15 \times 0.1 \\ P{\rm I} \\ 2 \\ 10.962(1) \\ 14.131(2) \\ 14.834(2) \\ 101.011(2) \\ 109.568(2) \\ 94.234(2) \\ 2101.6(4) \\ 1.40 \\ 0.81 \\ h: -14 \ to \ 14 \\ k: -18 \ to \ 18 \\ h: -19 \ to \ 19 \\ 19203 \\ 9895 \\ 5057 \\ 509 \\ 9895 \\ 5057 \\ 509 \\ 9895 \\ 0.061 \\ 0.187 \\ +0.8/-0.8 \\ \end{array}$

H, Me (pz)], 3.62 [dq, J = 7.0/5.2 Hz, 4 H, CH₂ (Et)], 3.82 (s, 6 H, OMe), 6.40 [s, 1 H, H (pz)], 6.94 [d, J = 2.1 Hz, 2 H, H (im)], 7.02 [d, J = 2.1 Hz, 2 H, H (im)], 7.03–7.48 (m, 11 H, Ar), 7.54–7.60 (m, 2 H, Ph) ppm. IR (KBr): $\tilde{v} = 2499$ w (BH), 1180 s, 1096 s, br. (ClO₄) cm⁻¹.

Catalytic Studies. Oxidation of 2-Propanol: 2a (33.8 mg, 0.05 mmol) and [BNA]Cl (12.5 mg, 0.05 mmol) were dissolved in [D₆]DMSO (2 mL). 2-Propanol (0.2 mL, 2.6 mmol) was added and the mixture stirred for 24 h at 100 °C in the dark. The oxidation reaction was monitored by recording the ¹H NMR signals of acetone at $\delta = 2.08$ ppm and of the benzyl group of BNAH at $\delta = 4.30$ ppm.

Reduction of *p*-Nitrobenzaldehyde: 2a (20.3 mg, 0.03 mmol), *p*-nitrobenzaldehyde (4.5 mg, 0.03 mmol) and BNAH (6.5 mg, 0.03 mmol) were dissolved in [D₆]DMSO (2 mL) and the mixture was stirred at 100 °C for 24 h. The reduction was monitored by recording the ¹H NMR signals of *p*-nitrobenzyl alcohol at δ = 4.25 ppm and of the benzyl group of BNA⁺ at δ = 5.91 ppm.

Structure Determinations:^[28] Crystals were obtained as described above. Diffraction data were recorded at 200 K (**2a**, **3c**, **3d**, **3f**, **4a**, **4b**, **5**, **8**) and at room temp. (**1c**, **1d**, **3a**, **3b**) with a Nonius CAD4 (**1d**, **3a**, **3d**, **5**) or a Bruker Smart CCD diffractometer (**1c**, **2a**, **3b**, **3c**, **3f**, **4a**, **4b**, **8**). Empirical absorption corrections were applied for **3b**, **3c**, **4a**, **4b** and **8**. The structures were solved with direct methods and refined anisotropically with the SHELX program suite.^[29] Hydrogen atoms were included with fixed distances and isotropic temperature factors 1.5-times those of their attached atoms. Parameters were refined against F^2 . The *R* values are defined as $R_1 = \Sigma [F_o - F_c]/\Sigma F_o$ and $wR_2 = \{\Sigma[w(F_o^2 - F_c^2)^2]\Sigma[w(F_o^2)^2]\}^{1/2}$. Drawings were produced with SCHAKAL.^[30] Table 1 lists the crystallographic data.

Acknowledgments

This work was supported by the Deutsche Forschungsgemeinschaft (Graduiertenkolleg "Ungepaarte Elektronen") and the Fonds der Chemischen Industrie. We are indebted to Drs. W. Deck, M Tesmer and H. Brombacher for assistance during the crystal structure determinations.

- ^[1] G. Parkin, Adv. Inorg. Chem. 1995, 42, 291-393.
- ^[2] N. Kitajima, W. B. Tolman, Prog. Inorg. Chem. 1995, 43, 419-531.
- ^[3] H. Vahrenkamp, Acc. Chem. Res. 1999, 32, 589-596.
- ^[4] G. Parkin, Chem. Commun. 2000, 1971-1985.
- ^[5] M. Garner, J. Reglinski, I. Cassidy, M. D. Spicer, A. R. Kennedy, *Chem. Commun.* **1996**, 1975–1976.

- ^[6] J. Reglinski, M. D. Spicer, M. Garner, A. R. Kennedy, J. Am. Chem. Soc. 2000, 121, 2317–2318.
- [7] I. Cassidy, M. Garner, A. R. Kennedy, G. B. S. Potts, J. Reglinski, P. A. Slavin, M. D. Spicer, *Eur. J. Inorg. Chem.* 2002, 1235–1239.
- [8] J. L. White, J. M. Tanski, D. Rabinovich, J. Chem. Soc., Dalton Trans. 2002, 2987–2991, and references cited therein.
- [9] B. M. Bridgewater, G. Parkin, *Inorg. Chem. Commun.* 2001, 4, 126–129.
- [10] M. Tesmer, M. Shu, H. Vahrenkamp, *Inorg. Chem.* 2001, 40, 4022-4029.
- ^[11] C. Kimblin, T. Hascall, G. Parkin, *Inorg. Chem.* **1997**, *36*, 5680-5681.
- [12] C. Kimblin, B. M. Bridgewater, T. Hascall, G. Parkin, J. Chem. Soc., Dalton Trans. 2000, 1267–1274.
- ^[13] C. Kimblin, B. M. Bridgewater, D. G. Churchill, T. Hascall, G. Parkin, *Inorg. Chem.* **2000**, *39*, 4240–4243.
- ^[14] J. Seebacher, M. Shu, H. Vahrenkamp, *Chem. Commun.* **2001**, 1026–1027.
- ^[15] J. Seebacher, H. Vahrenkamp, unpublished.
- ^[16] In 2002 the Cambridge Crystallographic Data File had 76 entries for zinc complexes of thiourea-derived ligands.
- [^{17]} R. Castro, J. A. Garcia-Vazquez, J. Romero, A. Sousa, Y. D. Chang, J. Zubieta, *Inorg. Chim. Acta* **1995**, *237*, 143–146.
- ^[18] H. Kayama, Y. Saito, Bull. Chem. Soc. Jpn. 1954, 27, 12-16.
- ^[19] C. Kimblin, B. M. Bridgewater, D. G. Churchill, G. Parkin, *Chem. Commun.* 1999, 2301–2302.
- ^[20] S. Ramaswamy, H. Eklund, B. V. Plapp, *Biochemistry* 1994, 33, 5230-5237.
- ^[21] R. Walz, H. Vahrenkamp, Inorg. Chim. Acta 2000, 314, 58-62.
- ^[22] Y. Zhang, H. Vahrenkamp, Inorg. Chim. Acta, in print.
- ^[23] M. Förster, R. Burth, A. K. Powell, T. Eiche, H. Vahrenkamp, *Chem. Ber.* **1993**, *126*, 2643–2648.
- ^[24] R. Alsfasser, M. Ruf, S. Trofimenko, H. Vahrenkamp, *Chem. Ber.* **1993**, *126*, 703-710.
- ^[25] J. Elguero, R. Jaquier, Bull. Soc. Chim. Fr. 1966, 2832-2845.
- ^[26] R. G. Jones, E. C. Kornfeld, K. C. McLaughlin, R. C. Anderson, J. Am. Chem. Soc. **1949**, 71, 4000–4002.
- ^[27] J. Kister, G. Assef, G. Mille, J. Metzger, *Can. J. Chem.* 1979, 57, 813–821.
- ^[28] CCDC-203392 (1c), -203393 (1d), -203394 (2a), -203395 (3a), -203396 (3b), -203397 (3c), -203398 (3d), -203399 (3f), -203400 (4a), -203401 (4b), -203402 (5) and -156428 (8) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge at www.ccdc.cam.ac.uk/conts/retrieving.html [or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; Fax: (internat.) + 44-1223/336-033; E-mail: deposit@ccdc.cam.ac.uk].
- ^[29] SHELXTL, program package of the Bruker Smart CCD diffractometer, version 5.1, **2002**.
- ^[30] E. Keller, *SCHAKAL for Windows*, Universität Freiburg, **1999**. Received February 12, 2003