This article was downloaded by: [University of Sydney] On: 04 January 2015, At: 17:02 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/uopp20

ALTERNATIVE SYNTHESIS OF N,N'-DJPHENYLTHIOUREA AND ITS ANALYSIS BY LC-MS

Yu-Chun Xiong ^a , Zhi-Min Zong ^a , Fa-Qin Jiang ^a , Xin-Hua Yuan ^a , Xiao-Hua Wang ^a , Ya-Fei Ji ^a , Li Xu ^b , Hong-Zhan Xiao ^b , Lin-Tao He ^b , Chun-Qi Li ^c & Xian-Yong Wei ^a

^a School of Chemical Engineering , China University of Mining and Technology , Xuzhou, 221008, Jiangsu, P.R. CHINA

^b Analytical Center , Beijing Institute of Microchemistry , Beijing, 100091, P.R., CHINA

^c National Institute of Advanced Industrial Science and Technology, Tsukuba, 305-8569, JAPAN E-mail: Published online: 11 Feb 2009.

To cite this article: Yu-Chun Xiong , Zhi-Min Zong , Fa-Qin Jiang , Xin-Hua Yuan , Xiao-Hua Wang , Ya-Fei Ji , Li Xu , Hong-Zhan Xiao , Lin-Tao He , Chun-Qi Li & Xian-Yong Wei (2003) ALTERNATIVE SYNTHESIS OF N,N²-DJPHENYLTHIOUREA AND ITS ANALYSIS BY LC-MS, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 35:4, 409-411, DOI: <u>10.1080/00304940309355849</u>

To link to this article: http://dx.doi.org/10.1080/00304940309355849

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

ALTERNATIVE SYNTHESIS OF N,N'-DIPHENYLTHIOUREA AND ITS ANALYSIS BY LC-MS

Submitted by (11/15/02)

Yu-Chun Xiong,[†] Zhi-Min Zong,[†] Fa-Qin Jiang,[†] Xin-Hua Yuan,[†] Xiao-Hua Wang,[†] Ya-Fei Ji,[†] Li Xu,^{††} Hong-Zhan Xiao,^{††} Lin-Tao He,^{††} Chun-Qi Li^{†††} and Xian-Yong Wei*[†]

- [†] School of Chemical Engineering China University of Mining and Technology Xuzhou 221008, Jiangsu, P.R. CHINA
- ^{††} Analytical Center, Beijing Institute of Microchemistry Beijing 100091, P.R. CHINA
- *** National Institute of Advanced Industrial Science and Technology Tsukuba 305-8569, JAPAN E-mail: wei_xian_yong@yahoo.com.cn

N,N'-Diphenylthiourea (DPTU) has been widely used during the past decades as medicine for leprosy, as a flotation agent to separate ferric sulfide from other metal sulfides, as a ligand to form N,N'-diphenylthioureido complexes with some metals such as ruthenium, osmium and iridium,¹ as a curing accelerator for caoutchouc and synthetic rubber (especially for natural latex and neoprene),^{2,3} as an extractant for isolating precious metals⁴ and as a thermal stabilizer for PVC. It also serves as an intermediate and a starting material for the preparation of diphenylguanidine, dyestuffs, curing bladder, rubber bag, tire cement, wires, cables, as a detergent for cleaning airport runways and many other commercial products. Therefore, extensive investigations have been carried out to devise an effective synthesis of DPTU and other N,N'-disubstituted thioureas.^{5,6} However, there still exist many practical limitations in current DPTU syntheses and some problems in DPTU analysis.

Ramadas *et al.*⁷ investigated the direct preparation of a series of *N*,*N'*-disubstituted thioureas from amines and carbon disulfide. Although DPTU was obtained in very high yields (98%), the need to add triethanolamine, Lac sulfur powder and a large amount of water makes the method inconvenient and less economically feasible. Ballabeni *et al.*⁸ studied the reaction of aniline with carbon disulfide at 100°C both in the presence and in the absence of catalyst. The maximum yield of DPTU was 83% with a basic catalyst, 43% with an acidic clay and 17% without any catalyst. They considered that the basic catalyst favored the elimination of H₂S and thereby significantly promoted the reaction. However, as they noted, since the catalyst activity is diminished by the H₂S generated during the reaction, higher yields of DPTU could not be achieved. Moreover, the necessity of using an autoclave makes their process inconvenient. Mohanta *et al.*⁹ recently reported an investigation of the synthesis of substituted thioureas using 1-(methyldithiocarbonyl)imidazole as a thiocarbonyl transfer reagent and obtained a high yield

of DPTU, albeit of low purity (mp. 150-152°C). In addition, their method is limited by the use of expensive 1-(methyldithiocarbonyl)imidazole or of N-methyl-N'-(methyldithiocarbonyl)imidazolium iodide as a reactant. We report herein our study on the reaction of aniline with carbon disulfide over sulfonated polystyrene.

 $PhNH_2 + CS_2 \qquad \xrightarrow{sulfonated} PhNH_2 + CS_2 \qquad \xrightarrow{sulfonated} PhNHCNHPh + H_2S$

The reaction was conducted at reflux temperature (*ca.* 60°C) under ambient pressure to afford DPTU selectively. After being cooled to room temperature, the reaction mixture deposited a large amount of crystals. To separate DPTU from the sulfonated polystyrene and to recover the catalyst, the solids, including the sulfonated polystyrene and DPTU, were extracted into carbon disulfide. DPTU can be readily separated from the extraction solution by crystallization and purified by recrystallization from carbon disulfide and the sulfonated polystyrene can be easily recovered. Carbon disulfide was used both as a reactant and as a solvent during the reaction and the extraction. Under the reaction conditions, the resulting H_2S is rapidly released from the reaction solution.

Because of its thermal lability and low volatility, DPTU cannot be analyzed by GC/MS. Only Mohanta *et al.*⁹ reported mass spectral data of DPTU, but the data are not reliable since GC/MS was used for the analysis. We analyzed DPTU using GC and GC/MS. The results show that DPTU decomposed completely to aniline and phenyl isothiocyanate (PITC). We performed LC/MS analysis of DPTU using 85% methanol- H_2O as mobile phase and obtained data (m/z: 251, 229, 136 and 94) which suggest that DPTU was only partly decomposed to aniline and PITC by the "soft ionization" of LC/MS, and thus LC/MS provides an effective approach to a less destructive analysis of DPTU.

EXPERIMENTAL SECTION

Aniline, carbon disulfide and sulfonated polystyrene (D001-FC, macroporous type, total exchange capacity $4.35 \ge \text{mmol/g}$, purchased from Shandong Dongda Chemical Industry Company) are commercially available and were used without further purification and any treatment. Mp was determined in capillaries on a domestic WRC-1B melting point apparatus and is uncorrected. ¹H NMR spectrum was recorded on a JEOL Lambda 400 spectrometer (400 MHz) with TMS as the internal standard in C₄D₈O. Chemical shifts are expressed in parts per million (δ , ppm). FTIR spectrum was obtained on a Nicolet Magna IR-560 spectrometer. LC/MS analysis was performed on a Bruker Esquire system (C18 column, 4.6 mm x 250 mm x 5 μ m; ESI source, positive ion detection, 113.4 V). Elemental analyses for C, H, N and S were carried out on a Leco CHN-2000 Elemental analyzer and a Leco SC-32 Sulfur Determinator, respectively.

Typical Procedure.- A mixture of aniline (10 mL, 0.11 mol), carbon disulfide (13 mL, 0.22 mol) and sulfonated polystyrene (1 g) in a 150 mL, round bottom flask was stirred under reflux

for 2 h. After being cooled to room temperature, the reaction mixture was transferred with the help of *ca*. 100 mL carbon disulfide into a Soxhlet extractor to separate catalyst from the resulting DPTU. After being allowed to cool to room temperature, the extract deposited a large amount of crystals, which were recrystallized from carbon disulfide to afford 22.5 g (90%) of DPTU (99.9% pure by LC) as white, flaky crystals, mp. 153.5-154°C, *lit.*⁴ 152-153°C. IR (KBr): 3211, 1596, 1551, 1494, 1450, 1342, 1241, 761, 757, 698, 486 cm⁻¹. ¹H NMR: δ 9.00 (s, 2H, S=C-NH), 7.44 (d, *J* = 8.3 Hz, 4H, ArH on 2-position), 7.27 (t, *J* = 7.7 Hz, 4H, ArH on 3-position), 7.07 (t, *J* = 7.9 Hz, 2H, ArH on 4-position). MS (*m*/z, %): 251.1 [(M+Na)⁺, 17.5], 229.1 [(M+H)⁺, 95.2], 136.1 [(M-PhNH₂+H)⁺, 100], 94.2 [(M-PhN=C=S+H)⁺, 68.0]. *Anal.* Calcd for C₁₃H₁₂N₂S: C, 68.39; H, 5.30; N, 12.27; S, 14.01

REFERENCES

- 1. S. D. Robinson, A. Sahajpal and J. W. Steed, Inorg. Chim. Acta, 306, 205 (2000).
- 2. G. Sridevi, J. P. Rao and K. K. Reddy, Synth. Commun., 19, 965 (1989).
- 3. G. L'abbé and S. Leurs, Tetrahedron, 48, 7505 (1992).
- 4. P. J. H. Seeverens, E. J. M. Klaassen and F. J. M. J. Maessen, Spectrochim. Acta Part B, 38, 727 (1983).
- 5. S. Ozaki, Chem. Rev., 72, 457 (1972).
- 6. N. Yamazaki, T. Tomioka and F. Higashi, Synthesis, 384 (1975).
- 7. K. Ramadas, N. Janarthanan and S. Velmathi, Synth. Commun., 27, 2255 (1997).
- M. Ballabeni, R. Ballini, F. Bigi, R. Maggi, M. Parrini, G. Predieri and G. Sartori, J. Org. Chem., 64, 1029 (1999).
- 9. P. K. Mohanta, S. Dhar, S. K. Samal, H. Ila and H. Junjappa, Tetrahedron, 56, 629 (2000).
