

Article

Subscriber access provided by Libraries of the | University of Hawaii at Manoa

# A Combinatorial Synthesis of Fused Tetracyclic Heterocycles Containing [1,6]Naphthyridine Derivatives under Catalyst Free Conditions

Chao Li, Xing-Ye Mu, Yu-Ling Li, Yun Liu, and Xiang-Shan Wang

ACS Comb. Sci., Just Accepted Manuscript • DOI: 10.1021/co400020w • Publication Date (Web): 12 Apr 2013

Downloaded from http://pubs.acs.org on April 19, 2013

## **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Combinatorial Science is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

## A Combinatorial Synthesis of <mark>Fused Tetracyclic Heterocycles</mark> Containing

## [1,6]Naphthyridine Derivatives under Catalyst Free Conditions

## Chao Li, Xing-Ye Mu, Yu-Ling Li, Yun Liu, Xiang-Shan Wang\*

A three-component reaction between an aromatic aldehyde, an amine, and *tert*-butyl 2,4-dioxopiperidine-1-carboxylate in EtOH at refluxing temperature gave fused tetracyclic heterocycles in high yields. The amines include 1*H*-indazol-5-amine, 1*H*-indazol-6-amine, 1*H*-indol-5-amine and 1*H*-benzo[*d*]imidazol-5-amine, giving 11-aryl-3*H*-indazolo[5,4-*b*][1,6] naphthyridine, 11-aryl-1*H*-indazolo[6,7-*b*][1,6]naphthyridine, 11-aryl-3*H*-imidazol[4',5':3,4]benzo[1,2-*b*][1,6]naphthyridine derivatives, respectively.



## A Combinatorial Synthesis of Fused Tetracyclic Heterocycles Containing

## [1,6]Naphthyridine Derivatives under Catalyst Free Conditions

Chao Li, Xing-Ye Mu, Yu-Ling Li, Yun Liu, Xiang-Shan Wang\*

School of Chemistry and Chemical Engineering, Jiangsu Key Laboratory of Green Synthetic

Chemistry for Functional Materials, Jiangsu Normal University, Xuzhou Jiangsu 221116, P. R.

China

E-mail: xswang1974@yahoo.com

Abstract—A three-component reaction between an aromatic aldehyde, an amine, and *tert*-butyl 2,4-dioxopiperidine-1-carboxylate in EtOH at refluxing temperature gave fused tetracyclic heterocycles in high yields. The amines include 1*H*-indazol-5-amine, 1*H*-indazol-6-amine, 1*H*-indol-5-amine and 1*H*-benzo[*d*]imidazol-5-amine, giving 11-aryl-3*H*-indazolo[5,4-*b*][1,6] naphthyridine, 11-aryl-1*H*-indazolo[6,7-*b*][1,6]naphthyridine, 11-aryl-3*H*-indolo[5,4-*b*][1,6]naphthyridine and 11-aryl-3*H*-imidazo[4',5':3,4]benzo[1,2-*b*][1,6]naphthyridine derivatives, respectively.

**Keywords**: [1,6]naphthyridine, 1*H*-indazol-5-amine, 1*H*-indazol-6-amine, 1*H*-indol-5-amine, 1*H*-benzo[*d*]imidazol-5-amine, *tert*-butyl 2,4-dioxopiperidine-1-carboxylate

### Introduction

Naphthyridines are an important class of heterocycles found in a variety of biologically active molecules. It is reported that they are used for the prevention and treatment of angiogenic disorders and cancers.<sup>1</sup> They are also used as various kinds of inhibitors, such as the first clinical stage inhibitor of protein kinase CK2 for the treatment of cancer, <sup>2</sup> novel DNA ligase inhibitors with anti-staphylococcal activity,<sup>3</sup> novel selective inhibitors of p38 mitogen activated protein kinase,<sup>4</sup> and the potent inhibitors of the protein kinase C isozymes.<sup>5</sup> In addition, there are many other remarkable applications reported in the literature.<sup>6</sup> Therefore, much attention has been devoted to the synthesis of these bio-active compounds.<sup>7</sup>

Indolonaphthyridine, indazolonaphthyridine and imidazobenzonaphthyridine, which contain both indole (indazole, imidazole) and naphthyridine rings, are fused tetracyclic heterocycles. These heterocyclic motifs are found in numerous alkaloid natural products, such as indolo[1,7]naphthyridine,<sup>8</sup> which is also named

*Incargranine B* (Figure 1, left). It was isolated from the whole plant of *Incarvillea mairei var. grandiflora*. Another alkaloid of *Canthin-6-one* (Figure 1, right) <sup>9</sup> containing indolonaphthyridine nucleus was obtained and identified from the branches of *Picrolemma granatensis*. Some of them have been used as a selective 5-HT2C/2B receptor antagonist,<sup>10</sup> a 5-HT2C/2B receptor antagonist with low 5-HT2A receptor affinity,<sup>11</sup> and the cardiovascular agents.<sup>12</sup> They have also been used as a inhibite of brain ischemia in rats for treating the brain disorder.<sup>13</sup>



Figure 1. The interesting structures of natural products containing indolonaphthyridine

To the best of our knowledge, only a few references exist concerning their synthesis.<sup>14</sup> Although these reported reactions have developed some useful synthetic procedures, still several limitations remained. For example, most of the procedures involve several steps, or low yields, or metal catalysts, or organic solvents. Thus, a simple, efficient, and green method to synthesize these fused tetracyclic heterocycles would be attractive.

Distinguished from classical sequential two-component synthetic procedures, multi-component reactions (MCRs) use three or more chemical starting materials for product formation. They have attracted considerable attention in organic synthesis and medicinal chemistry owing to their convergence and productivity.<sup>15</sup> As a continuation of our research devoted to the preparation of heterocycles by three-component reactions,<sup>16</sup> herein, we would like to report a convenient and efficient three-component synthesis of fused tetracyclic heterocycles. They include 3*H*-indazolo[5,4-*b*][1,6]naphthyridine, 1*H*-indazolo[6,7-*b*][1,6]naphthyridine, 3*H*-indolo[5,4-*b*][1,6]naphthyridine and 3*H*-imidazo[4',5':3,4]benzo[1,2-*b*][1,6] naphthyridine derivatives, respectively.

## **Results and Discussions**

Treatment of 4-chlorobenzaldehyde  $1\{1\}$ , *tert*-butyl 2,4-dioxopiperidine-1-carboxylate  $2\{1\}^{17}$  and 1*H*-indazol-5-amine  $3\{1\}$  in refluxing EtOH under catalyst free conditions, gave *tert*-butyl 7,8-dihydro-10-oxo-11-aryl-3*H*-indazolo[5,4-*b*][1,6] naphthyridine-9(6*H*,10*H*,11*H*)-carboxylate  $4\{1,1,1\}$  in high yield and regioselectivity, rather than *tert*-butyl 6,7-dihydro-9-oxo-10-aryl-1*H*-indazolo[5,6-*b*] [1,6]naphthyridine-8(5*H*,9*H*,10*H*)-carboxylate 4' (Scheme 1).



Scheme 1. Optimization for the synthesis of  $4\{1,1,1\}$ 

Subsequently, the reaction of 4-chlorobenzaldehyde  $1\{1\}$ ,  $2\{1\}$  and  $3\{1\}$  was used as a representative to optimize the experimental conditions. The reaction was first carried out in EtOH at room temperature for 10 h and only trace amount of product  $4\{1,1,1\}$  was observed by TLC (Table 1, entry 1). The reaction temperature is key to product yields. At reflux, EtOH gives the best results. Furthermore, it should be noted that the products dissolved completely in refluxing EtOH, and crystallized in cold EtOH, therefore, the products of  $4\{1,1,1\}$  were obtained directly by filtration without further purification. In addition, the effects of different solvents, such as CH<sub>3</sub>CN,

 benzene, DMF, and THF (Table 1, entries 5-8) were also compared, and EtOH gave

## the highest yield.

**Table 1.** Synthetic results of  $4\{1,1,1\}$  under different reaction conditions<sup>*a*</sup>

Entry	temperature (°C)	Time(h)	Solvent	Yields $(\%)^b$
1	r.t.	10	EtOH	Trace
2	50	16	EtOH	68
3	Reflux	10	EtOH	93
4	80	8	Toluene	87
5	Reflux	14	CH <sub>3</sub> CN	82
6	Reflux	16	Benzene	89
7	80	10	DMF	78
8	Reflux	14	THF	85

<sup>*a*</sup> Reagents and conditions: 4-chlorobenzaldehyde  $\mathbf{1}$ {*I*} (0.140 g, 1.0 mmol),  $\mathbf{2}$ {*I*} (0.213 g, 1.0 mmol),  $\mathbf{3}$ {*I*} (0.133 g, 1.0 mmol), solvent (10 mL). <sup>*b*</sup> Isolated yields.

Based on the optimized reaction conditions, we next explored variations in the aldehyde **1** (Scheme 1), the diversity of aldehyde  $1{I-20}$  was shown in Figure 2. Table 2 demonstrates that the reaction also tolerates significant functionalization of the aromatic aldehydes; both electron-donating (such as alkoxyl group) and electron-withdrawing (such as halide) groups can be accommodated, and substituents *ortho, meta*, or *para* to the aromatic group can all be generated. All aldehydes reacted well to provide a library of naphthyridine derivatives **4** (Table 2). Subsuquently, we investigated the applicability of amines, and found that 1H-indazol-5-amine  $3{I}$ , 1H-indazol-6-amine  $3{2}$ , 1H-indol-5-amine  $3{3}$  and 1H-benzo[d]imidazol-5-amine  $3{4}$  (Figure 3) all gave satisfactory results (Table 2). It should be noted that  $4{10,1,4}$  was *tert*-butyl 11-aryl-10-oxo-7,8,10,11-tetrahydro-1H-imidazo[4',5':3,4] benzo[1,2-b][1,6]naphthyridine-9(6H)-carboxylate rather than 3H-imidazol-5-amine  $3{4}$ 

was used as a reactant. The crystal structure of  $4\{10,1,4\}$  is shown in Figure 4. In order to prove this was not a special example; another single crystal of  $4\{12,1,4\}$  was grown and determined. It gave the same result as shown in Figure 5. All the products of 4 were characterized by <sup>1</sup>H NMR, IR and HRMS, and the data were in good agreement with the preconceived structures.







Figure 3. Diversity of amines 3{1-4}



Scheme 2. Regioselective synthesis of naphthyridines 4

Entry	Ar	Product 4	Time /h	Yields /%
1	$4-ClC_6H_4$	<b>4</b> { <i>1</i> , <i>1</i> , <i>1</i> }	10	93
2	$4-BrC_6H_4$	$4\{2,1,1\}$	14	90
3	4-MeOC <sub>6</sub> H <sub>4</sub>	$4{3,1,1}$	14	96
4	3-ClC <sub>6</sub> H <sub>4</sub>	<b>4</b> {4,1,1}	13	92
5	3-BrC <sub>6</sub> H <sub>4</sub>	4{5.1.1}	10	87
6	2-ClC <sub>6</sub> H <sub>4</sub>	4{6.1.1}	10	92
7	2-BrC <sub>6</sub> H <sub>4</sub>	$4{711}$	12	90
8	3 4 - C + C + C + C	$4{811}$	8	90
9	$3.5-(MeO)-C_{2}H_{2}$	$4{9}11$	8	87
10	$3_{-}MeOC_{-}H$		10	95
10	$C \square$	$4_{\{10,1,1\}}$	10	90
11	C <sub>6</sub> 115 Diporonul	$4\{11,1,1\}$	12	90
12		$4\{12,1,1\}$	10	93
13	$4 - C_{1}C_{6}T_{4}$	$4\{1,1,2\}$ 4(2,1,2)	12	90
14	$4-BIC_6H_4$	$4\{2,1,2\}$	12	90
15	$3-CIC_6H_4$	$4{4,1,2}$	15	82
16	$3-BrC_6H_4$	$4{5,1,2}$	16	86
17	$2-CIC_6H_4$	$4\{0,1,2\}$	10	92
18	$2\text{-BrC}_6\text{H}_4$	$4\{/,1,2\}$	12	90
19	$3,4-Cl_2C_6H_3$	<b>4</b> {8,1,2}	14	93
20	Piperonyl	<b>4</b> { <i>12</i> , <i>1</i> , <i>2</i> }	16	89
21	$4-FC_6H_4$	<b>4</b> { <i>13</i> , <i>1</i> , <i>2</i> }	12	85
22	$4-NO_2C_6H_4$	<b>4</b> { <i>14</i> , <i>1</i> , <i>2</i> }	10	86
23	$2,3-Cl_2C_6H_3$	<b>4</b> { <i>15</i> , <i>1</i> , <i>2</i> }	10	92
24	2-Thienyl	<b>4</b> { <i>20</i> , <i>1</i> , <i>2</i> }	10	90
25	$4-ClC_6H_4$	<b>4</b> { <i>1</i> , <i>1</i> , <i>3</i> }	9	89
26	$4-BrC_6H_4$	<b>4</b> {2,1,3}	12	89
27	$4-CH_3OC_6H_4$	<b>4</b> { <i>3</i> , <i>1</i> , <i>3</i> }	10	90
28	$3-ClC_6H_4$	<b>4</b> { <i>4</i> , <i>1</i> , <i>3</i> }	10	85
29	3,4-Cl <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	<b>4</b> { <i>8</i> , <i>1</i> , <i>3</i> }	10	90
30	3,5-(MeO) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	<b>4</b> { <i>9</i> , <i>1</i> , <i>3</i> }	12	87
31	3-CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	<b>4</b> { <i>10</i> , <i>1</i> , <i>3</i> }	12	86
32	4-FC <sub>6</sub> H <sub>4</sub>	4{13,1,3}	10	90
33	$2-FC_6H_4$	4{16,1,3}	6	86
34	3.4-Me <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	4{17.1.3}	12	84
35	$2.4-Cl_2C_6H_3$	4{18,1,3}	8	92
36	2-Thienvl	4{20,1,3}	9	85
37	4-ClC <sub>6</sub> H <sub>4</sub>	4{1.1.4}	12	86
38	4-BrC <sub>4</sub> H <sub>4</sub>	$4\{2,1,4\}$	10	79
39	4-CH <sub>2</sub> OC <sub>2</sub> H <sub>4</sub>	$4{314}$	13	79
40	3-ClC/H	$4{4   4}$	13	85
41	2 - C + C + C + C + C + C + C + C + C + C	$4{614}$	9	89
42	$34-Cl_{0}C_{14}$	$4{81}$	10	86
42	3, -0.206113 3, 5-(MeO) - C - H	$1_{\{0,1,7\}}$	14	90
	3-CH.OC.H	<b>T</b> (2,1,75 A510 1 A)	17	90
44	Diperonyl	$\pi_{1}, 10, 1, 4$	12	86
<del>4</del> 0	riperonyi	<b>-</b> {12,1,4}	15	00

4{19,1,4}

2-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>





<sup>a</sup> Reaction condition: 1 (1.0 mmol), 2 (0.213 g, 1.0 mmol), 3 (1.0 mmol), EtOH (10.0 mL), 80 °C.

Scheme 3. Regioselective synthesis of product  $4\{10,1,4\}$ 



**Figure 4.** Crystal structure of 4{10,1,4} with DMF solvate

X-ray diffraction analysis indicates that the 1,4-dihydropyridine rings in  $4\{10,1,4\}$ and  $4\{12,1,4\}$  are adopt a half-chair conformation. The atoms C1, C7, C9, C13 and N4 are coplanar, with C8 deviating from the defined planes by 0.297 (3) for  $4\{10,1,4\}$ and 0.285 (4) Å for  $4\{12,1,4\}$ , respectively. In  $4\{10,1,4\}$ , the adjacent pyridine ring also adopts a half-chair conformation, the atom C11 deviates from the defined plane by 0.657 (3) Å. However, it adopts a skew boat conformation in  $4\{12,1,4\}$ , with C11 and N3 deviating from the defined plane by 0.749 (6) and 0.197 (5) Å, respectively. The center 1,4-dihydropyridine ring is nearly parallel to the benzimidazole and

#### **ACS Combinatorial Science**

pyridine rings, forming the dihedral angles of 6.2 (1) and 13.8 (2)° (9.4 (1) and 9.0 (1)° for  $4\{10,1,4\}$ ), respectively. Meanwhile, it is nearly perpendicular to the benzene ring, forming a dihedral angle of 88.3 (1)° (87.5 (1)° for  $4\{10,1,4\}$ ).



**Figure 5.** Crystal structure of 4{12,1,4} with DMF solvate

According to the structures of **4** and the references,<sup>18</sup> we think the subsequent Knoevenagel condensation, Michael addition, intra-molecular nucleophilic addition and dehydration may take place in the formation of the observed products. The key step with high regioselectivity is the Michael addition reaction between 1*H*-indazol-6-amine and  $\alpha$ , $\beta$ -unsaturated ketone. The activity of the 7-position is higher than that of the 5-position in 1*H*-indazol-6-amine, because it is both the  $\alpha$ -site of an amino group and the benzyl position of a pyrazole ring at the same time. The possible mechanism is outlined in Scheme 4 (using 1*H*-indazol-6-amine as an example).



Scheme 4. The possible mechanism for the formation of 4

## Conclusion

In conclusion, we found a mild, efficient and highly regioselective method for the synthesis of fused tetracyclic heterocycles containing [1,6]naphthyridine derivatives via a three component reaction of an aromatic aldehyde, amine and *tert*-butyl 2,4-dioxopiperidine-1-carboxylate under catalyst free conditions. The procedure affords high regioselectivity while featuring mild reaction conditions, high yields and operational simplicity without catalysts.

### 4. Experimental

Melting points were determined in open capillaries and are uncorrected. IR spectra were recorded on a Tensor 27 spectrometer in KBr pellet. <sup>1</sup>H NMR spectra was obtained from a solution in DMSO- $d_6$  with Me<sub>4</sub>Si as internal standard using a Bruker-400 spectrometer. HRMS analyses were carried out using a Bruker-micro-TOF-Q-MS analyzer.

### General procedure for the synthesis of 4

A dry 25 mL flask was charged with aromatic aldehyde **1** (1.0 mmol), amine (1.0 mmol), *tert*-butyl 2,4-dioxopiperidine-1-carboxylate (0.213 g, 1.0 mmol), and EtOH (10.0 mL). The reaction mixture was stirred at refluxing temperature for 8-16 h until all the reactant amine was consumed (monitored by TLC). The products were obtained by filtration without further purification, after the reaction mixture was cooled to room temperature.

tert-Butyl 11-(4-chlorophenyl)-10-oxo-7,8,10,11-tetrahydro-3H-indazolo[5,4-b][1,6]

naphthyridine-9(6*H*)-carboxylate  $4\{1,1,1\}$ : M. p.: 221~223 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz):  $\delta_{\rm H}$  1.41 (s, 9H, 3CH<sub>3</sub>), 2.59~2.64 (m, 1H, CH), 2.75~2.82 (m, 1H, CH), 3.50~3.60 (m, 1H, CH), 3.96~4.02 (m, 1H, CH), 5.52 (s, 1H, CH), 7.07 (d, *J* = 8.4 Hz, 1H, ArH), 7.23 (d, *J* = 8.4 Hz, 2H, ArH), 7.34~7.37 (m, 3H, ArH), 8.01 (s, 1H, ArH), 9.67 (s, 1H, NH), 13.01 (s, 1H, NH). <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>, 100 MHz):  $\delta_{\rm C}$  27.8, 30.7, 35.7, 38.5, 80.8, 98.6, 109.5, 113.7, 116.7, 121.5, 127.9, 128.7, 129.5, 130.3, 146.7, 148.3, 148.4, 152.6, 162.3, 164.2. IR (KBr): *v* 3297, 3079, 2977, 2933, 2885, 1719, 1681, 1605, 1543, 1492, 1401, 1369, 1331, 1253, 1213, 1142, 1092, 1049, 949, 846, 805, 775, 738, 703 cm<sup>-1</sup>. HRMS (ESI, *m*/*z*): Calcd for C<sub>24</sub>H<sub>22</sub>N<sub>4</sub>O<sub>3</sub>Cl [M - H]<sup>-</sup> 449.1380, found 449.1384.

### Acknowledgement:

We are grateful to the National Natural Science foundation of China (20802061, 21104064), a project funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions, Qing Lan Project (10QLD008, GSFM2011003) and College Industrialization Project (JHB2012-31) of Jiangsu Province for financial support.

Supporting Information Available: Characterized data inclauding NMR, IR and HRMS of 4, copies of CIFs for  $4\{10,1,4\}$  and  $4\{12,1,4\}$  are available free of charge via the internet at http://pubs.acs.org.

#### **References and notes**

1. Hadj-Slimane, R.; Garbay, C. Pharmaceutical compositions comprising neuropilin inhibitors, and their use for the prevention and/or treatment of angiogenic disorders and cancers, Eur. Pat. Appl. **2012**, EP 2522341 A1 20121114.

2. (a) Pierre, F.; Chua, P. C.; O'Brien, S. E.; Siddiqui-Jain, A.; Bourbon, P.; Haddach, M.; Michaux, J.; Nagasawa, J.; Schwaebe, M. K.; Stefan, E. Discovery and SAR of 5-(3-chlorophenylamino)benzo[*c*][2,6]naphthyridine-8-carboxylic acid (CX-4945), the first clinical stage inhibitor of protein kinase CK2 for the treatment of cancer, *J. Med. Chem.* **2011**, *54*, 635-654. (b) Battistutta, R.; Cozza, G.; Pierre, F.; Papinutto, E.; Lolli, G.; Sarno, S.; O'Brien, S. E.; Siddiqui-Jain, A.; Haddach, M.; Anderes, K. Unprecedented selectivity and structural

determinants of a new class of protein kinase CK2 inhibitors in clinical trials for the treatment of cancer, *Biochemistry* **2011**, *50*, 8478-8488,

Surivet, J.-P.; Lange, R.; Hubschwerlen, C.; Keck, W.; Specklin, J.-L.; Ritz, D.; Bur, D.; Locher,
H.; Seiler, P.; Strasser, D. S. Structure-guided design, synthesis and biological evaluation of novel
DNA ligase inhibitors with in vitro and in vivo anti-staphylococcal activity, *Bioorg. Med. Chem. Lett.* 2012, *22*, 6705-6711.

 Lumeras, W.; Vidal, L.; Vidal, B.; Balague, C.; Orellana, A.; Maldonado, M.; Dominguez, M.; Segarra, V.; Caturla, F. 1,7-Naphthyridine 1-oxides as novel potent and selective inhibitors of p38 mitogen activated protein kinase, *J. Med. Chem.* 2011, *54*, 7899-7910.

5. van Eis, M. J.; Evenou, J.-P.; Floersheim, P.; Gaul, C.; Cowan-Jacob, S. W.; Monovich, L.; Rummel, G.; Schuler, W.; Stark, W.; Strauss, A. 2,6-Naphthyridines as potent and selective inhibitors of the novel protein kinase C isozymes, *Bioorg. Med. Chem. Lett.* **2011**, *21*, 7367-7372.

6. (a) Press, N. J.; Taylor, R. J.; Fullerton, J. D.; Tranter, P.; McCarthy, C.; Keller, T. H.; Arnold, N.; Beer, D.; Brown, L.; Cheung, R. Solubility-driven optimization of phosphodiesterase-4 inhibitors leading to a clinical candidate, J. Med. Chem. 2012, 55, 7472-7479. (b) Haynes, N.-E.; Scott, N. R.; Chen, L. C.; Janson, C. A.; Li, Jia, K.; Lukacs, C. M.; Railkar, A.; Tozzo, E.; Whittard, T.; Brown, N. F. Identification of an adamantyl azaguinolone JNK selective inhibitor, ACS Med. Chem. Lett. 2012, 3, 764-768. (c) Bauer, U.; Giordanetto, F.; Bauer, M.; O'Mahony, G.; Johansson, K. E.; Knecht, W.; Hartleib-Geschwindner, J.; Carlsson, E. T.; Enroth, C. Discovery of 4-hydroxy-1,6-naphthyridine-3-carbonitrile derivatives as novel PDE10A inhibitors, Bioorg, Med. Chem. Lett. 2012, 22, 1944-1948. (d) Sato, Yusuke; Zhang, Y.; Seino, T.; Sugimoto, T.; Nishizawa, S.; Teramae, N. Highly selective binding of naphthyridine with a trifluoromethyl group to cytosine opposite an abasic site in DNA duplexes, Org. Biomol. Chem. 2012, 10, 4003-4006. (e) Zhang, Y.; Quock, L. P.; Chung, E.; Ohgami, Y.; Quock, R. M. Involvement of a NO-cyclic GMP-PKG signaling pathway in nitrous oxide-induced antinociception in mice, Eur. J. *Pharm.* 2011, 654, 249-253. (f) Varadi, L.; Gray M.; Groundwater, P. W.; Hall, A. J; James, A. L.; Orenga, S.; Perry, J. D.; Anderson, R. J. Synthesis and evaluation of fluorogenic 2-amino-1,8-naphthyridine derivatives for the detection of bacteria, Org. Biomol. Chem. 2012, 10, 2578-2589.

7. (a) Ioannidou, H. A.; Martin, A.; Gollner, A.; Koutentis, P. A. Three-step synthesis of ethyl

#### **ACS Combinatorial Science**

canthinone-3-carboxylates from ethyl 4-bromo-6-methoxy-1,5-naphthyridine-3-carboxylate via a Pd-catalyzed Suzuki-Miyaura coupling and a Cu-catalyzed amidation reaction, J. Org. Chem. 2011, 76, 5113-5122. (b) Li, M.; Shao, P.; Wang, S.-W.; Kong, W.; Wen, L.-R. Four-component cascade heteroannulation of heterocyclic ketene aminals: Synthesis of functionalized tetrahydroimidazo[1,2-a]pyridine derivatives, J. Org. Chem. 2012, 77, 8956-8967. (c) Takahashi, Y.; Kubota, T.; Shibazaki, A.; Gonoi, T.; Fromont, J.; Kobayashi, J. Nakijinamines C-E, new heteroaromatic alkaloids from the sponge Suberites species, Org. Lett. 2011, 13, 3016-3018. (d) Mailyan, A. K.; Peregudov, A. S.; Dixneuf, P. H.; Bruneau, C.; Osipov, S. N. Cyclobutene ring-opening of bicyclo[4.2.0]octa-1,6-dienes: Access to CF<sub>3</sub>-substituted 5,6,7,8-tetrahydro-1,7-naphthyridines, J. Org. Chem. 2012, 77, 8518-8526. (e) Silva, D.; Chioua, M.; Samadi, A.; Carmo, C. M.; Jimeno, M.-L.; Mendes, E.; Rios, C. L.; Romero, A.; Villarroya, M.; Lopez, M. G. Synthesis and pharmacological assessment of diversely substituted pyrazolo[3,4-b]quinoline, and benzo[b]pvrazolo[4,3-g][1,8]naphthyridine derivatives, Eur. J. Med. Chem. 2011, 46, 4676-4681. (f) Wang, K.-Y.; Chen, C.; Liu, J.-F.; Wang, Q.; Chang, J.; Zhu, H.-J.; Li, C. Novel multifunctional organic semiconductor materials based on 4,8-substituted 1,5-naphthyridine: synthesis, single crystal structures, opto-electrical properties and quantum chemistry calculation, Org. Biomol. Chem. 2012, 10, 6693-6704. (g) Tian, C.; Jiao, X.; Liu, X.; Li, R.; Dong, L.; Liu, X.; Zhang, Z.; Xu, J.; Xu, M.; Xie, P. First total synthesis and determination of the absolute configuration of 1-N-methyl-3-methylamino-[N-butanoic acid-3-(9-methyl-8-propen-7-one)amide]-benzo[f][1,7]naphthyridine-2-one, a novel benzonaphthyridine alkaloid, Tetrahedron Lett. 2012, 53, 4892-4895. (h) Rustagi, V.; Tiwari, R.; Verma, A. K. AgI-catalyzed cascade strategy: regional re naphthyridines, thienopyridines, and quinoxalines in water, Eur. J. Org. Chem. 2012, 4590-4602. 8. Shen, Y.-H.; Su, Y.-Q.; Tian, J.-M.; Lin, S.; Li, H.-L.; Tang, J.; Zhang, W.-D. A unique indolo-[1,7]naphthyridine alkaloid from incarvillea mairei var. grandiflora (Wehrh.) grierson,

Helvetica Chimica Acta 2010, 93, 2393-2396.

9. Rodrigues Fo., E.; Fernandes, J. B.; Vieira, P. C.; Da Silva, M.; Fatima, das G. F. Canthin-6-one alkaloids from Picrolemma granatensis, *Phytochemistry* **1992**, *31*, 2499-2501.

10. Muehle, H.; Nozulak, J.; Cercus, J.; Kuesters, E.; Beutler, U.; Penn, G.; Zaugg, W. Sample for the evolution of the synthesis of a development product. Preparation of the indolonaphthyridine

derivative SDZ SER-082 in enantiomeric purity as a selective 5-HT2C/2B receptor antagonist, *Chimia* **1996**, *50*, 209-213.

Nozulak, Joachim; Kalkman, Hans O.; Floerscheim, Philipp; Hoyer, Daniel; Schoeffter, Philipp; Buerki, Hans R. (+)-cis-4,5,7*a*,8,9,10,11,11*a*-Octahydro-7*H*-10-methylindolo[1,7-*bc*]
[2,6]-naphthyridine: A 5-HT2C/2B receptor antagonist with low 5-HT2A receptor affinity, *J. Med. Chem.* 1995, *38*, 28-33.

12. Yanagisawa, H.; Shimoji, Y.; Hashimoto, T. Preparation of indolonaphthyridine derivatives as cardiovascular agents, *Jpn. Kokai Tokkyo Koho* **1993**, JP 05310738 A 19931122.

 Oomoto, T.; Tanaka, J. Indolonaphthyridinedione derivative for treatment of brain disorder, *Jpn. Kokai Tokkyo Koho* 1993, JP 05262648 A 19931012.

14. (a) Mastalarz, H.; Jasztold-Howorko, R.; Rulko, F.; Croisy, A.; Carrez, D. Synthesis and cytostatic properties of some 6H-indolo[2,3-b][1,8]naphthyridine derivatives, Arch. Pharm. 2004, 337, 434-439. (b) Rastorgueva, N. A.; Ryabova, S. Yu.; Lisitsa, E. A.; Alekseeva, L. M.; Granik, V. G. Synthesis of derivatives of a new heterocyclic system, indolo[2,3-f][1,7]naphthyridine, Russian Chemical Bulletin, 2003, 52, 2149-2156. (c) Zhang, Q.; Shi, C.; Zhang, H.-R.; Wang, K. K. Synthesis of 6H-indolo[2,3-b][1,6]naphthyridines and related compounds as the 5-aza analogues of ellipticine alkaloids, J. Org. Chem. 2000, 65, 7977-7983. (d) Gribble, G. W.; Saulnier, M. G.; Obaza-Nutaitis, J. A.; Ketcha, D. M. A versatile and efficient construction of the 6H-pyrido[4,3-b]carbazole ring system. Syntheses of the antitumor alkaloids ellipticine, 9-methoxyellipticine, and olivacine, and their analogs, J. Org. Chem. 1992, 57, 5891-5899. (e) Benson, S. C.; Li, J. H.; Snyder, John K. Indole as a dienophile in inverse electron demand Diels-Alder reactions. 3. Intramolecular reactions with 1.2.4-triazines to access the canthine skeleton, J. Org. Chem. 1992, 57, 5285-5287. (f) Da Settimo, A.; Primofiore, G.; Ferrarini, P. L.; C.; Martini, C.; Pennacchi, E.; Lucacchini, A. Synthesis of new Mori. 6H-indolo[2,3-b][1,8]naphthyridines and their specific inhibition of benzodiazepine receptor, Farmaco, Edizione Scientifica 1986, 41, 577-85. (g) Nivoliers, F.; Decormeille, A.; Godard, A.; Queguiner, G. 6H-indolo[3,2-b]naphthyridines. Aza-analogs of ellipticine. Tetrahedron Lett. 1980, 21, 4485-4486. (h) Da Settimo, A.; Primofiore, G.; Santerini, V.; Biagi, G.; D'Amico, L. One-step synthesis of 6H-indolo[2,3-b][1,8]naphthyridines. A new heterocyclic ring system, J. Org. Chem. , 42, 1725-1728.

### **ACS Combinatorial Science**

1
2
3
4
5
5
6
7
8
õ
9
10
11
12
12
13
14
15
16
17
17
18
19
20
21
21
22
23
24
25
20
26
27
28
20
29
30
31
32
33
33
34
35
36
27
31
38
39
40
11
41
42
43
44
15
40
46
47
48
10
43
50
51
52
52
53
54
55
56
57
57
58
59

60

15. (a) Vijjay, N. C.; Rajesh, A. U.; Vinod, S.; Bindu, A. R.; Sreekanth, J. S.; Lakshmi, B. Strategies for heterocyclic construction via novel multicomponent reactions based on isocyanides and nucleophilic carbenes, *Acc. Chem. Res.* **2003**, *36*, 899-907. (b) Tietze, L. F. Domino reactions in organic synthesis, *Chem. Rev.* **1996**, *96*, 115-136.

16. (a) Wang, X. S.; Sheng, J.; Lu, L.; Yang, K.; Li, Y. L. Combinatorial synthesis of 3-arylideneaminoquinazolin-4(1*H*)-one derivatives catalyzed by iodine in ionic liquids, *Acs Comb. Sci.* 2011, *13*, 196-199. (b) Wang, X. S.; Yin, M. Y.; Wang, W.; Tu, S. J. A stereoselective Povarov reaction leading to *exo*-tetrahydroindolo-[3,2-*c*]quinoline derivatives catalyzed by iodine, *Eur. J. Org. Chem.* 2012, (25), 4811-4818. (c) Wang, X. S.; Zhou, J.; Yin, M. Y.; Yang, K.; Tu, S. J. Efficient and highly selective method for the synthesis of benzo(naphtho)quinoline derivatives catalyzed by iodine, *J. Comb. Chem.* 2010, *12*, 266-269.

Orsini, P.; Maccario, A.; Colombo, N. Regioselective γ-alkylation of *tert*-butyl
2,4-dioxopiperidine-1-carboxylate, *Synthesis* 2007, (20), 3185-3190.

 (a) Wang, X. S.; Zhang, M. M.; Zeng, Z. S.; Shi, D. Q.; Tu, S. J.; Wei, X. Y.; Zong, Z. M. A Simple and clean procedure for the synthesis of polyhydroarcidine and quinoline derivatives: Reaction of Schiff base with 1,3-dicarbonyl compounds in aqueous medium, *Tetrahedron Lett*.
2005, 46, 7169-7173. (b) Shi, F.; Zhang, S.; Wu, S.-S.; Gao, Y.; Tu, S. J. A diversity-oriented synthesis of pyrazolo[4,3-f]quinoline derivatives with potential bioactivities via microwaveassisted multi-component reactions. *Molecular Diversity*, 2011, 15, 497-505.