


This article was downloaded by: [Brown University]

On: 11 January 2013, At: 20:21

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcyc20>

Stereoselective Synthesis of Substituted 2-(Z-Styrylsulfonyl)-1H-imidazoles and Benzothiazole

N. Guravaiah^a & V. Rajeswar Rao^a

^a Department of Chemistry, National Institute of Technology, Warangal, India

Version of record first published: 22 Feb 2010.

To cite this article: N. Guravaiah & V. Rajeswar Rao (2010): Stereoselective Synthesis of Substituted 2-(Z-Styrylsulfonyl)-1H-imidazoles and Benzothiazole, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 40:6, 808-813

To link to this article: <http://dx.doi.org/10.1080/00397910903009422>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

STEREOSELECTIVE SYNTHESIS OF SUBSTITUTED 2-(Z-STYRYLSULFONYL)-1H-IMIDAZOLES AND BENZOTHIAZOLE

N. Guravaiah and V. Rajeswar Rao

Department of Chemistry, National Institute of Technology, Warangal, India

A novel method has been reported involving the synthesis of z-styryl sulfides by the reaction of phenylacetylene with 2-mercapto benzimidazoles and benzthiazole using NaOH as a base in absolute ethanol. On further oxidation with H₂O₂, these resulted in the formation of the desired styryl sulfones in good yields with retention of stereochemistry.

Keywords: 2-Mercaptoimidazoles; phenylacetylene; styryl sulfides; sulfones

INTRODUCTION

The use of styrylsulfones in organic chemistry and medicinal chemistry has increased dramatically.^[1] Styrylsulfones have been employed in many synthetic methodologies as intermediates, enabling the preparation of great number of functionalized products, such as natural products and bioactive substances. The functional group is found in numerous biologically interesting compounds. These compounds include anticancer^[2–4] and inhibitors for several enzymes such as cyclooxygenase-2(COX-2).^[5] In contrast to different methods developed for Z,Z- and E,E-bis(styryl)sulfides and sulfones,^[6] less attention was focused on stereoselective synthesis of styryl sulfones. In fact, the latter were reported earlier from our laboratories by condensation of sodium *E*-styrenesulfonates with α -halocarbonyl compounds^[7] and 2-halomethyl benzimidazoles.^[8] Our sustained interest in this field led us to develop a simple and elegant methodology for a new class of stereoselective styrylsulfones through a facile route. In the literature,^[4] it has been reported that the synthesis of the title compounds requires a multistep synthesis (five steps). The first step involved in the preparation of 1H-benzo[d]imidazole-2-thiols is condensation of orthophenylene diamines with CS₂ in the presence of potassium hydroxide (KOH) in ethanol. On reaction with various phenacyl halides in the presence of K₂CO₃ in acetone, these gave 2-(1H-benzo[d]imidazol-2-ylthio)-1-phenylethanones. They were converted to 2-[(1H-benzo[d]imidazol-2-yl)sulfanyl]-1-phenyl-1-ethanols by reduction with sodium borohydride. These compounds were oxidized to get 2-[(1H-benzo[d]imidazol-2-yl)sulfonyl]-1-phenyl-1-ethanols. Ultimately, they were subjected to

Received February 13, 2009.


Address correspondence to V. Rajeswar Rao, Department of Chemistry, National Institute of Technology, Warangal, AP-506 004, India. E-mail: vramesw@yahoo.com

dehydration with methanesulfonyl chloride to afford 1-{sulfonyl}2-{(E)-2-phenyl-1-ethyl)sulfonyl} derivatives, which in turn were demesylated to 1H-benzo[d]imidazole-2-yl[(E)-2-phenyl-1-ethyl)sulfones with 33% HBr in acetic acid. Recently, it was reported that certain styrylsulfone derivatives like FRI-20 inhibit tumor cell growth and viability by inhibiting the mitogen-activated protein kinase (MAPK) signal transduction pathway. The compounds regulate the extracellular signal-related kinase (ERK) and inhibit the proliferation of breast and prostate tumor cells in a dose-dependent manner without affecting normal cell growth. The cell growth inhibitory activity of this compound is dictated by the nature and position of the functional groups. In view of the wide range of biological activities exhibited by heteryl styrylsulfones, we report the stereoselective synthesis of some heteryl(Z)-styryl sulfones. These methodologies are quite useful, but they have some limitations, such as requiring the isolation of intermediates, longer reaction times, and lower overall yields. It is thus evident that there remains a scope for the development of clean and efficient methodologies for the preparation of the title compounds.


Herein, we report the synthesis of Z-styrylsulfones in a two-step reaction (Schemes 1 and 2). The first step involves the nucleophilic addition of the substituted 2-mercapto benzimidazoles, 2-mercapto benzthiazoles, and 4,5 diphenyl-2-mercapto imidazoles to phenyl acetylene in the presence of a base in ethanol. The reflux period varies from 12 to 16 h. The progress of the reaction was monitored by thin-layer chromatography (TLC). After completion of the reaction, the mixture was cooled to room temperature, poured into crushed ice, and neutralized with conc. HCl to give 3-(1-phenyl-2-(Z-styrylthio)-1H-imidazol-4-yl)-2H-chromen-2-ones. Oxidation of these sulfides with hydrogen peroxides in acetic acid resulted in the formation of Z-styrylsulfones. The compounds were purified by column chromatography and fully characterized by elemental analysis, IR, ^1H NMR, ^{13}C NMR, and mass spectrometry.

EXPERIMENTAL

All reagents and solvents were obtained from commercial sources and used without further purification unless otherwise stated. Reactions were monitored by



Scheme 1. R = OCH_3 , X = NH; R = OCF_2H , X = NH; R = H, X = NH; R = H, X = s.



Downloaded by [Brown University] at 20:21 11 January 2013

Downloaded by [Brown University] at 20:21 11 January 2013

Downloaded by [Brown University] at 20:21 11 January 2013

Downloaded by [Brown University] at 20:21 11 January 2013

Spectral Data of the Compounds

5-Methoxy-2-(styrylthio)-1H-benzo[d]imidazole. Yield: 85%, mp 134–136°C. IR (KBr, γ_{\max} cm⁻¹): 1594 (C=C), 1625 (C=N), 3403 (NH). ¹H NMR (CDCl₃ + DMSO-d₆, δ ppm): 3.68 (s, 3H OCH₃), 6.61 (d, 1H, J = 12 Hz, styryl), 6.71 (d, 1H, J = 2 Hz, Ar-H), 6.74 (s, 1H, Ar-H), 7.02 (d, 1H, J = 12 Hz, styryl), 7.16 (m, 1H, Ar-H), 7.26–7.33 (m, 5H, Ar-H). EI-MS 282 (M⁺). ¹³C NMR (CDCl₃ + DMSO-d₆, δ ppm) 54.9, 110.6, 119.5, 126.7, 127.3, 127.6, 127.8, 135.2, 146.1, 155.2. Anal. calcd. for C₁₆H₁₄N₂OS: C, 68.06; H, 5.00; N, 9.92. Found: C, 68.00; H, 4.94; N, 9.95%.

5-(Difluoromethoxy)-2-(styrylthio)-1H-benzo[d]imidazole. Yield 88%, mp 108–110°C. IR (KBr, γ_{\max} cm⁻¹): 1596 (C=C), 1624 (C=N), 3426 (NH). ¹H NMR (CDCl₃, δ ppm): 6.85 (d, 1H, J = 12 Hz, styryl), 7.27 (d, 1H, J = 12 Hz, styryl), 7.31–7.39 (m, 7H, Ar-H), 7.44 (t, 1H, OCF₂H), 7.53 (d, 1H, Ar-H, J = 6 Hz), 9.72 (s, 1H, NH), EI-MS 319 (M + H)⁺. Anal. calcd. for C₁₆H₁₂F₂N₂OS: C, 60.37; H, 3.80; N, 8.80. Found: C, 60.33; H, 3.85; N, 8.85%.

2-(Styrylthio)-1H-benzo[d]imidazole. Yield 92%, mp 130–132°C. IR (KBr, γ_{\max} cm⁻¹): 1597 (C=C), 1616 (C=N), 3394 (NH). ¹H NMR (CDCl₃, δ ppm): 6.75 (d, 1H, J = 12 Hz, styryl), 6.85 (d, 1H, J = 12 Hz, styryl), 7.10 (d, 1H, J = 8 Hz Ar-H), 7.25–7.45 (m, 8H, Ar-H). Anal. calcd. for C₁₅H₁₂N₂S: C, 71.40; H, 4.79; N, 11.10. Found: C, 71.45; H, 4.83; N, 11.00%.

2-(Styrylthio)benzo[d]thiazole. Yield 80%, mp 114–116°C. IR (KBr, γ_{\max} cm⁻¹): 1598 (C=C), 1616 (C=N), 3348 (NH). ¹H NMR (CDCl₃, δ ppm): 6.65 (d, 1H, J = 12 Hz, styryl), 6.85 (d, 1H, J = 12 Hz, styryl), 7.20–7.49 (m, 9H, Ar-H). Anal. calcd. for C₁₅H₁₁NS₂: C, 66.88; H, 4.12; N, 5.20. Found: C, 66.84; H, 4.10; N, 5.24%.

4,5-Diphenyl-2-(styrylthio)-1H-imidazole. Yield 78%, mp 138–140°C. IR (KBr, γ_{\max} cm⁻¹): 1595 (C=C), 1623 (C=N), 3448 (NH). ¹H NMR (CDCl₃, δ ppm): 6.76 (d, 1H, J = 12 Hz, styryl), 7.01 (d, 1H, J = 12 Hz, styryl), 7.28–7.51 (m, 15H, Ar-H), 12.95 (s, 1H, NH). ¹³C NMR (CDCl₃, δ ppm) 122.7, 127.2, 127.4, 127.6, 127.7, 128.2, 128.4, 128.6, 135.9, 138.7. EI-MS 355 (M + H)⁺. Anal. calcd. for C₂₃H₁₈N₂S: C, 77.93; H, 5.12; N, 7.90. Found: C, 77.90; H, 5.10; N, 7.86%.

General Procedure for the Synthesis of Z-Styrylsulfones

To a solution containing Z-styrylsulfides (15 mmol) in 20 mL of glacial acetic acid, 30% hydrogen peroxide (8 mL) was added dropwise at room temperature. After the addition was complete, the reaction mixture was allowed to stir at room temperature for 24 h. The mixture was then poured into ice-cold water and stirred for 10 min. The solid was separated, filtered, and dried to give the required compound. All the other compounds were prepared similarly.

Spectral Data

5-Methoxy-2-(styrylsulfonyl)-1*H*-benzo[*d*]imidazole. Yield 90%, mp 220–222°C. IR (KBr, $\gamma_{\max} \text{ cm}^{-1}$): 1155, 1356(SO₂), 1508 (C=C), 1601 (C=N), 3434 (NH). ¹H NMR (CDCl₃, δ ppm): 3.90 (s, 3H OCH₃), 7.05 (d, 1H, *J* = 12 Hz, styryl), 7.35 (d, 1H, *J* = 12 Hz, styryl), 7.60–7.81 (m, 8H, Ar-H). EI-MS 313 (M-H)⁺. Anal. calcd. for C₁₆H₁₄N₂O₃S: C, 61.13; H, 4.49; N, 8.91. Found: C, 61.00; H, 4.38; N, 8.96%.

5-(Difluoromethoxy)-2-(styrylsulfonyl)-1*H*-benzo[*d*]imidazole. Yield 89%, mp 198–200°C. IR (KBr, $\gamma_{\max} \text{ cm}^{-1}$): 1148, 1341 (SO₂), 1594 (C=C), 1626 (C=N), 3588 (NH). ¹H NMR (CDCl₃, δ ppm): 7.0 (d, 1H, *J* = 12 Hz, styryl), 7.18 (d, 1H, *J* = 12 Hz, styryl), 7.20–7.40 (m, 9H, Ar-H). Anal. calcd. for C₁₆H₁₂F₂N₂O₃S: C, 54.85; H, 3.45; N, 8.00. Found: C, 54.80; H, 3.41; N, 8.10%.

2-(Styrylsulfonyl)-1*H*-benzo[*d*]imidazole. Yield 92%, mp 213–215°C. IR (KBr, $\gamma_{\max} \text{ cm}^{-1}$): 1119, 1344 (SO₂), 1618 (C=N), 1606 (C=C), 3588 (NH). ¹H NMR (CDCl₃, δ ppm): 7.15 (d, 1H, *J* = 12 Hz, styryl), 7.45 (d, 1H, *J* = 12 Hz, styryl), 7.60–7.80 (m, 9H, Ar-H). Anal. calcd. for C₁₅H₁₂N₂O₂S: C, 63.36; H, 4.25; N, 9.85. Found: C, 63.39; H, 4.28; N, 9.88%.

2-(Styrylsulfonyl)benzo[*d*]thiazole. Yield 88%, mp 208–210°C. IR (KBr, $\gamma_{\max} \text{ cm}^{-1}$): 1141, 1315 (SO₂), 1626 (C=N), 1606 (C=C), 3392 (NH). ¹H NMR (CDCl₃, δ ppm): 7.10 (d, 1H, *J* = 12 Hz, styryl), 7.30 (d, 1H, *J* = 12 Hz, styryl), 7.4 (d, 1H, *J* = 8 Hz, Ar-H), 7.50–7.70 (m, 8H, Ar-H). Anal. calcd. for C₁₅H₁₁NO₂S₂: C, 59.78; H, 3.68; N, 4.65. Found: C, 59.81; H, 3.70; N, 4.69%.

4,5-Diphenyl-2-(styrylsulfonyl)-1*H*-imidazole. Yield 85%, mp 228–230°C. IR (KBr, $\gamma_{\max} \text{ cm}^{-1}$): 1173, 1395 (SO₂), 1509 (C=C), 1601 (C=N), 3430 (NH). ¹H NMR (CDCl₃, δ ppm): 7.15 (d, 1H, *J* = 12 Hz, styryl), 7.25 (d, 1H, *J* = 12 Hz, styryl), 7.30–7.45 (m, 15H, Ar-H). EI-MS 387 (M + H)⁺. Anal. calcd. for C₂₃H₁₈N₂SO₂: C, 71.48; H, 4.69; N, 7.25. Found: C, 71.50; H, 4.72; N, 7.12%.

CONCLUSIONS

In conclusion, we have developed a simple, inexpensive, and efficient stereoselective synthesis of styrylsulfones without using any catalyst. The study of the biological activity of these new styrylsulfones is in progress and will be published elsewhere.

ACKNOWLEDGMENT

This work was supported by grants from the Council of Scientific Industrial Research (CSIR), New Delhi, for Project No. 01 (2062) 06/EMR-II.

REFERENCES

1. Sharma, V. M.; Adi Seshu, K. V.; Chandra Sekhar, V.; Madan, S.; Vishnu, B.; Babu, P. A.; Vamsee Krishna, C.; Sreenu, J.; Ravi Krishna, V.; Venkateswarlu, A.; Rajagopal, S.;

- Ajaykumar, R.; Kumar, T. S. Synthesis and biological evaluation of [4-(2-phenylethenesulfonylmethyl)phenyl]-quinazolin-4-yl-amines as orally active anti-cancer agents. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 67–71.
- Ramana Reddy, M. V.; Muralidhar Reddy, M.; Stephen, C. C.; Venkat, R. P.; Iqbal, N. M.; Kang, A. D.; Premkumar Reddy, E. Design, synthesis, and biological evaluation of (*E*)-styrylbenzylsulfones as novel anticancer agents. *J. Med. Chem.* **2008**, *51*, 86–100.
 - (a) Premkumar Reddy, E.; Ramana Reddy, M. V. Styryl sulfones anticancer agents. US Patent 6548553, April 15, 2003; (b) Premkumar Reddy, E.; Ramana Reddy, M. V. Styryl sulfones anticancer agents. European Patent 1027330, September 12, 2004; (c) Premkumar Reddy, E.; Ramana Reddy, M. V. Styryl sulfones anticancer agents. US Patent 6359013, March 19, 2002; (d) Premkumar Reddy, E.; Ramana Reddy, M. V. Z-Styryl sulfone anticancer agents. U.S. Patent, 6201154, March 13, 2001.
 - Manohar Sharma, V.; Aravind Babu, P.; Chandrasekhar, V.; Vamseekrishna, C.; Sreenu, J.; Venkata Adishesu, K.; Ravikrishna, V.; Vishnu, B.; Venkateswarlu, A.; Rajgopal, S.; Ajaya Kumar, R.; Sravankumar, T.; Prem Kumar, M.; Ramanujam, R.; Gopalakrishnan, B.; Akash, K.; Vijay, U. V.; Srinivas Rao, M. New styryl sulfones as anticancer agents. *Eur. J. Med. chem.* **2003**, *38*, 811–824.
 - Ramana Reddy, M. V.; Muralidhar Reddy, M.; Venkat, R. P.; Padmavathi, V.; Rengasamy, B.; Stanley, C. B.; Premkumar Reddy, E. Design, synthesis, and biological evaluation of (*E*)- and (*Z*)-styryl-2-acetoxyphenyl sulfides and sulfones as cyclooxygenase-2 inhibitors. *J. Bioorg. Med. Chem.* **2005**, *13*, 1715–1723.
 - Takikawa, Y.; Shimada, K.; Matsumoto, H.; Tanabe, H.; Takizawa, S. A convenient preparation of *z*-styryl, *z,z*-, and *e,z*-distyryl sulfides in liquid ammonia. *Chem. Lett.* **1983**, 1351–1354; (b) Ramana Reddy, M. V.; Manjubhashini, A. B.; Reddy, S.; Ramana Reddy, P. V.; Bhaskar Reddy, D. A novel and convenient synthesis of E,E-bis(styryl) sulfones and E,E-1,4-diaryl-1,3-butadienes. *Syn. Commun.* **1991**, *21*, 1589–1596; (c) Kamigata, N.; Ozaki, J.; Kobayashi, M. Reaction of alkenesulfonyl chlorides with olefins catalyzed by a ruthenium(II) complex: A novel method for synthesis of (E,E)-1,4-diaryl-1,3-butadienes. *J. Org. Chem.* **1985**, *50*, 5045–5050.
 - Guravaih, N.; Rajeswar Rao, V. Facile polyethylene glycol (PEG-400)-promoted synthesis of some new heteryl (*E*)-styrylsulfones. *J. Chem. Res.* **2009**, *4*, 237–239.
 - Guravaih, N.; Rajeswar Rao, V. An economical and efficient one pot synthesis of (*E*)-styrylsulfone. *J. Chem. Res.* (in press).
 - Bhatt, M. V.; Iyer, B. H.; Guna, P. C. 2-Thiol-4:5-diphenylimidazole derivatives. *Curr. Sci.* **1948**, *17*, 184.