

Article

Electronic Structures and Reactivity Profiles of Aryl Nitrenoid-Bridged Dicopper Complexes

Kurtis M. Carsch, James T. Lukens, Ida M. DiMucci, Diana A. Iovan,
Shao-Liang Zheng, Kyle M. Lancaster, and Theodore A. Betley

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.9b09616 • Publication Date (Web): 09 Jan 2020

Downloaded from pubs.acs.org on January 9, 2020

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Electronic Structures and Reactivity Profiles of Aryl Nitrenoid–Bridged Dicopper Complexes

Kurtis M. Carsch,[†] James T. Lukens,[‡] Ida M. DiMucci,[‡] Diana A. Iovan,[†] Shao-Liang Zheng,[†] Kyle M. Lancaster,^{‡*} Theodore A. Betley^{†*}

[†]Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street, Cambridge MA 02138, United States

[‡]Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca NY 14853, United States

ABSTRACT: Dicopper complexes templated by dinucleating, pacman dipyrin ligand scaffolds (^{Mes}dmx, ^{tBu}dmx: dimethylxanthine-bridged, cofacial bis-dipyrin) were synthesized by deprotonation/metalation with mesitylcopper (CuMes; Mes: mesityl) or by transmetalation with cuprous precursors from the corresponding deprotonated ligand. Neutral imide complexes (^Rdmx)Cu₂(μ²–NAr) (R: Mes, ^tBu; Ar: 4-MeOC₆H₄, 3,5-(F₃C)₂C₆H₃) were synthesized by treatment of the corresponding dicuprous complexes with aryl azides. While one-electron reduction of (^{Mes}dmx)Cu₂(μ²–N(C₆H₄OMe)) with potassium graphite initiates an intramolecular, benzylic C–H amination at room temperature, chemical reduction of (^{tBu}dmx)Cu₂(μ²–NAr) leads to isolable [(^{tBu}dmx)Cu₂(μ²–NAr)]^{0/–} product salts. The electronic structures of the thermally robust [(^{tBu}dmx)Cu₂(μ²–NAr)]^{0/–} complexes were assessed by variable-temperature electron paramagnetic resonance spectroscopy, X-ray absorption spectroscopy (Cu L_{2,3}/K-edge, N K-edge), optical spectroscopy, and DFT/CASSCF calculations. These data indicate that the formally Class IIIA mixed valence complexes of the type [(^Rdmx)Cu₂(μ²–NAr)][–], featuring significant NAr-localized spin following reduction from electronic population of the [Cu₂(μ²–NAr)] π* manifold and contrasting previous methods for engendering iminyl character through chemical oxidation. The reactivity of the isolable imido and iminyl complexes are examined for prototypical radical-promoted reactivity (e.g., nitrene transfer, H-atom abstraction), where the divergent reactivity is rationalized by the relative degree of N-radical character afforded from different aryl substituents.

1.0 Introduction

The incorporation of heteroatom functionality into unactivated C–H bonds remains a paramount challenge in both fine chemical synthesis and the conversion of hydrocarbon feedstocks into commodity chemicals.^{1–4} Dicopper complexes featuring bridging imido or oxo moieties have been implicated in the amination and hydroxylation of aliphatic C–H bonds, albeit by two distinct mechanisms. The dicopper imido complexes reported by Warren and coworkers are proposed to be in equilibrium with a mononuclear copper complex and a terminal copper nitrenoid species, to which the latter is attributed for the observed C–H bond amination reactivity (Figure 1).^{5–8} Steric profile tuning of the ancillary ligands for both β-diketiminate and dipyrin platforms allowed for observation of reductively coupled terminal nitrenoid complexes⁸ as well as the isolation of authentic terminal copper nitrene complexes,⁹ respectively. The amination reactivity can be contrasted with the partial hydroxylation of methane to methanol by zeolite Cu-ZSM-5,¹⁰ ascribed to a bent, dicupric oxide core (Figure 1).¹¹ The bent geometry of the zeolite-confined Cu₂(μ²–O) core has been proposed to endow oxyl-character on the oxenoid fragment in the transition state, facilitating low-barrier H-atom abstraction and radical recombination yielding methanol.¹² We were thus interested in synthesizing dicopper coordination complexes featuring bridging moieties to establish rigorous electronic structure to reactivity correlations. More specifically, we seek to probe how redox changes to a [Cu₂(μ²–E)]ⁿ core can induce spin accumulation at the bridgehead ligand, promote radical reactivity across the Cu–E bond vector, and achieve substrate functionalization.

To correlate the electronic structure with the reactivity profile of bimetallic transition metal complexes, we have prepared a dinucleating, dimethylxanthene-tethered, bis(dipyrin) ligand scaffold which proximally orients two metals cofacially to facilitate multi-electron chemistry. These architectures are well suited to stabilizing bridged transition metal species as evidenced by the isolation of bridging ferrous oxo and hydroxo complexes.¹³ Furthermore, we hypothesize a semi-rigid spacer minimizes entropic penalties and promotes a dinuclear arrangement for the exploration of dinuclear bridging complexes across distinct redox states. These design principles are prevalent in tethered diiron porphyrin complexes,^{14–16} competent for catalytic hydroxylation of exogenous substrates upon photolysis. In addition, the dipyrin scaffold limits access to the primary coordination sphere, rendering the metal ions high-spin and electrophilic.^{17–19}

In this study, we examine nitrene group transfer to a dicopper core. The electronic structure and reactivity profile of the [Cu₂(μ²–NR)] core is assessed as a function of redox state. Notably, we observe that one-electron reduction of the neutral [Cu₂(μ²–NR)] complex induces spin accumulation at the bridging nitrenoid fragment (Figure 1), leading to a diagnostic iminyl signature via N K-edge spectroscopy that is indistinguishable from subvalent iminyl agents generated from oxidation of the metal imido precursors. The reactivity profiles of both the neutral and anionic [Cu₂(μ²–NR)] complexes are assessed in the contexts of nitrene transfer to nucleophiles and propensity for H-atom abstraction, with divergent pathways as a function of the aryl substituent attributable to varying degrees of radical character at the nitrenoid bridgehead.





Figure 1. Dicopper mediated C–H bond functionalization: (top) monomer-dimer imido equilibrium for alkane amination (Warren); (middle) $[\text{Cu}_2(\mu^2\text{-O})]$ polarization for methane hydroxylation (Solomon); (bottom) radical accumulation on imido bridge for $[\text{Cu}_2(\mu^2\text{-NR})]^-$ (this work).

2. Results

2.1 Synthesis of Dinuclear Dicopper Imido Complexes

Synthesis of the mesityl-flanked, cofacial bis(dipyrrin) ligand featuring a dimethylxanthene spacer ($^{\text{Mes}}\text{dmx})\text{H}_2$ was modified from the procedure to prepare the reported ($^{\text{tBu}}\text{dmx})\text{H}_2$ analogue (see Supporting Information for experimental details).¹³ Acid-catalyzed condensation of 2-mesityl-1H-pyrrole²⁰ with dimethylxanthene bis(aldehyde),²¹ followed by oxidation of the bis(dipyrrromethane) with excess 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ, 3.0 equiv.) afforded ($^{\text{Mes}}\text{dmx})\text{H}_2$ in excellent yield (79%) on multi-gram scales as a red-orange powder. Under an atmosphere of nitrogen, ($^{\text{Mes}}\text{dmx})\text{H}_2$ was deprotonated with mesitylcopper²² in a thawing benzene/acetonitrile solvent mixture (20:1, 10 equiv. acetonitrile) to afford ($^{\text{Mes}}\text{dmx})\text{Cu}_2(\text{NCMe})_2$ (**1**) as a red-pink powder (Scheme 1). Trituration of the crude reaction mixture with cold acetonitrile precipitated **1** in analytically pure form (82%). Single crystals of **1** suitable for X-ray diffraction were prepared by layering acetonitrile with a concentrated diethyl ether solution of **1** at -35°C overnight. The solid-state structure displays a large copper-copper distance (6.3800(7) Å) and non-cofacial arrangement ($28.7(1)^\circ$ distortion with respect to the dipyrrin planes), suggesting no intramolecular communication between individual Cu-dipyrrin units (Figure S-72).


Scheme 1



Heating a benzene solution of **1** with a minor excess (1.2 equiv.) of 4-methoxyphenyl azide (4-MeOC₆H₄N₃) or 3,5-bis(trifluoromethyl)phenyl azide (3,5-(F₃C)₂C₆H₃N₃) (45 °C, 16 h) afforded the dicopper imido complexes ($^{\text{Mes}}\text{dmx})\text{Cu}_2(\mu^2\text{-N}(\text{C}_6\text{H}_4\text{OMe}))$ (**2**) and ($^{\text{Mes}}\text{dmx})\text{Cu}_2(\mu^2\text{-N}(3,5\text{-(F}_3\text{C)}_2\text{C}_6\text{H}_3))$ (**3**), respectively (Scheme 2). Both reactions were accompanied by gradual effervescence with a marked color change from red-pink (**1**) to deep violet (**2**) or purple-pink (**3**). Trituration of imido complexes **2** and **3** with cold acetonitrile allowed for isolation of analytically pure products after filtration in excellent yields (**2**: 92%, **3**: 79%). The diamagnetic ¹H NMR spectra, satisfactory elemental analyses, and featureless perpendicular-mode EPR spectra indicate an overall singlet spin ground state and establish bulk purity for **2** and **3** (Figure S-7, S-9).

The solid-state structures of **2** and **3** (Figure 2a, S-75) determined by single crystal X-ray diffraction reveal contractions of the copper-copper distances (**2**: 2.822(1) Å, **3**: 2.844(1) Å) as compared to **1** (6.3800(7) Å). These bond metrics are comparable to the geometric sum of the Van der Waals radii (approx. 1.40 Å)²³ and nearly 0.10 Å shorter than reported dinuclear copper imido complexes, (Table 1).^{5-7,24} This geometric contraction results in a more cofacial arrangement of the dipyrrin units (**2**: $6.7(1)^\circ$ torsion, **3**: $7.0(1)^\circ$ torsion) and a relatively acute copper-imido-copper bond angle (**2**: $102.3(1)^\circ$, **3**: $103.1(2)^\circ$). The displacement of the copper centers with respect to the dipyrrin plane (**2**: 0.879(6) Å, 0.854(5) Å; **3**: 0.848(3) Å, 0.848(3) Å) compared to **1** (0.242(4) Å, 0.407(3) Å) reflects the capability of the tethered bis(dipyrrin) ligand platform to support diverse intermetallic distances.

Scheme 2



We also explored how modifications of the dipyrrin flanking units influence the resulting $[\text{Cu}_2]$ core properties. In contrast to metalation of the ($^{\text{Mes}}\text{dmx})\text{H}_2$ platform, metalation of ($^{\text{tBu}}\text{dmx})\text{H}_2$ in the presence of coordinating solvents with mesitylcopper did not furnish the desired bis(cuprous) synthon. Consequently, an alternative transmetalation protocol was pursued. Exposure of ($^{\text{tBu}}\text{dmx})\text{H}_2$ in benzene to potassium bis(trimethylsilyl)amide (2.1 equiv.) rapidly precipitated an insoluble orange powder, which was isolated by filtration and rinsed with boiling toluene to furnish the putative dianionic $[(^{\text{tBu}}\text{dmx})\text{K}_2]$ (**4**). Due to the insolubility of **4** in organic media, **4** was not structurally characterized and used without further purification. Treatment of **4** as a suspension in tetrahydrofuran with excess copper bromide dimethylsulfide (2.2 equiv.) followed by addition of excess aryl azide (1.8 equiv.; 4-MeOC₆H₄N₃ or 3,5-(F₃C)₂C₆H₃N₃; 75 °C, 16 h) afforded the corresponding imido complexes ($^{\text{tBu}}\text{dmx})\text{Cu}_2(\mu^2\text{-N}(\text{C}_6\text{H}_4\text{OMe}))$ (**5**) and ($^{\text{tBu}}\text{dmx})\text{Cu}_2(\mu^2\text{-N}(3,5\text{-(F}_3\text{C)}_2\text{C}_6\text{H}_3))$ (**6**), respectively, as deep purple solids (Scheme 3). Both products were isolated by filtration in cold acetonitrile and recrystallized from a saturated acetonitrile/diethyl ether solution at -35°C in moderate yield (**5**: 65 %, **6**: 49 %). Single crystal X-ray diffraction reveals short copper-copper distances (**5**: 2.837(1) Å, **6**: 2.875(1) Å) akin to those of **2** and **3** (Figure 2b, S-76), (Table 1).

Table 1. Selected Bond Lengths (Å) and Angles (°) for isostructural complexes **2**, **3**, **5**–**9**.

	2	3 [†]	5 [‡]	6 [‡]	7	8	9
Cu1–Cu2	2.822(1)	2.844(1)	2.837(1)	2.875(1)	2.9031(7)	2.9448 (7)	2.8401(5)
Cu1–Ndipyrin	1.949(4)	1.923(2)	1.942(7)	1.934(8)	1.972(3)	2.058(2)	1.973(2)
Cu1–Ndipyrin	1.930(4)	1.937(2)	1.958(7)	1.929(7)	2.039(3)	1.985(2)	2.043(2)
Cu2–Ndipyrin	1.925(4)	1.923(2)	1.929(7)	1.931(8)	2.058(3)	2.028(2)	1.999(2)
Cu2–Ndipyrin	1.962(3)	1.937(2)	1.977(6)	1.963(7)	1.958(2)	2.004(2)	2.006(2)
Cu1–N_{Ar}	1.822(4)	1.814(4)	1.822(7)	1.830(8)	1.852(3)	1.869(2)	1.860(2)
Cu2–N_{Ar}	1.802(3)	1.814(4)	1.821(7)	1.818(7)	1.848(3)	1.860(2)	1.853(2)
Cu1–N_{Ar}–Cu2	103.2(2)	103.3(2)	102.3(3)	104.0(4)	103.4(1)	104.3(1)	99.8(1)
N_{Ar}–C_{ipso}	1.375(6)	1.406(4)	1.383(10)	1.386(12)	1.318(4)	1.351(3)	1.342(3)
C_{ipso}–C_{ortho}	1.406(7)	1.404(4)	1.402(11)	1.410(12)	1.438(5)	1.438(4)	1.435(4)
C_{ipso}–C_{ortho}	1.424(7)	1.404(4)	1.391(12)	1.404(13)	1.428(5)	1.417(3)	1.420(4)

[†]Cu1 and Cu2 are symmetry-equivalent. [‡]Two molecules are present in the asymmetric unit.

Scheme 3






Figure 2. Solid state molecular structures of (a) (Mes-dmx)Cu₂(μ²-N(C₆H₄OMe)) (**2**), (b) (tBu-dmx)Cu₂(μ²-N(3,5-(F₃C)₂C₆H₃)) (**6**) at 50 % ellipsoid probability. Color scheme: Cu (cobalt blue), F (yellow-green), N (blue), O (red). Solvent molecules, solid-state disorder, and hydrogen atoms are omitted for clarity.

2.3 Redox Chemistry of [Cu₂(μ²-NR)]

To assess the redox properties of the dicopper imido complexes, we collected cyclic voltammetry (CV) on imido complexes **2**, **3**, **5**, and **6** in tetrahydrofuran (0.01 mM sub-

strate, 0.2 M [ⁿBu₄N][PF₆]). One quasi-reversible reduction was observed at moderate potentials for both *p*-OMe substituted imidos (**2**: −1.32 V, **5**: −1.18 V vs. [Cp₂Fe]⁺⁰, Figure S–61), indicating the [Cu₂(μ²-NAr)] core could be chemically reduced. Imido complexes **3** and **6** similarly reveal quasi-reversible reductive processes at milder potentials consistent with the imido electron withdrawing substituents (**3**: −0.95 V, **6**: −0.70 V vs. [Cp₂Fe]⁺⁰, Figure S–61). Accordingly, treatment of **2**, **5**, or **6** in thawing tetrahydrofuran with stoichiometric potassium graphite (KC₈) in the presence of Cryptand 222 (C₂₂₂, 1.2 equiv.) precipitated insoluble graphite and furnished a rapid color change from deep purple to brown-pink (**2**, **5**) or brown-orange (**6**) (Scheme 4). For the reduction product of **2**, an isolated yield could not be obtained due to the thermal instability of the product (*vide infra*); nonetheless, rapid addition of stoichiometric silver trifluoromethanesulfonate (AgOTf) *in situ* following KC₈ addition at low temperature afforded quantitative recovery of **2** with deposition of metallic silver, suggesting the single-electron reduction proceeds in quantitative yield to a new paramagnetic species. Single-crystals of **7** suitable for X-ray diffraction were obtained by layering the crude reduction reaction of **2** in tetrahydrofuran with a thawing benzene/hexanes mixture at −35 °C, ensuring all manipulations were conducted at cryogenic temperatures (see Supporting Information for experimental details). Identification of the brown-pink crystals by single crystal X-ray diffraction revealed the mono-anionic bimetallic copper imido complex, [K(C₂₂₂)][(^{Mes}dmx)Cu₂(μ²-N(C₆H₄OMe))] (**7**), (Figure 3a). Similarly, bulk recrystallization of the reduction of **5** and **6** with stoichiometric KC₈ in the presence of C₂₂₂ from a tetrahydrofuran/diethyl ether solution layering at −35 °C afforded large crystals of [K(C₂₂₂)][(^{tBu}dmx)Cu₂(μ²-NAr)] (Ar: 4-MeOC₆H₄, **8**; 3,5-(F₃C)₂C₆H₃, **9**). Unlike thermally sensitive **7**, anions **8** and **9** persist in solution at room temperature over several days.

Scheme 4



Single crystal X-ray diffraction confirmed the composition of anionic imido complexes **7–9**, representative structures of which are provided in Figure 3 for **7** and **9**. All three anionic complexes display elongation of the Cu–N_{imido} bonds (average, Å, **2**: 1.812(4), **7**: 1.850(3); **5**: 1.822(7), **8**: 1.865(2); **6**: 1.824(8), **9**: 1.857(2), all relevant bond lengths are provided in Table 1). Complexes **7** and **9** display elongated Cu–Cu bonds (**7**: 2.9031(7) Å; **9**: 2.8401(5) Å) whereas **8** reveals a Cu–Cu bond contraction (**8**: 2.9448(7) Å). The reduction similarity proceeds with elongation of the dipyrin-copper bond lengths (average, Å, **2**: 1.942(4), **7**: 2.007(3); **5**: 1.952(8), **8**: 2.019(2); **6**: 1.939(8), **9**: 2.005(2)), in accord with more electron-rich Cu ion. Further bond metrics analysis reveals a contraction of the N_{imido}–C_{isop} bond (average, Å, **2**: 1.375(6); **7**: 1.318(4); **5**: 1.383(10); **8**: 1.351(3); **6**: 1.386(12); **9**: 1.342(3)) and elongation of the C_{isop}–C_{ortho} bond (average/Å, **2**: 1.415(7), **7**: 1.433(5); **5**: 1.397(12), **8**: 1.427(3); **6**: 1.407(13), **9**: 1.428(4)).




Figure 3. Solid state molecular structures of (a) [K(C₂₂₂)][(^{Mes}dmx)Cu₂(μ²-N(C₆H₄OMe))] (**7**) at 35 % ellipsoid probability and (b) [K(C₂₂₂)][(^{tBu}dmx)Cu₂(μ²-N(3,5-(F₃C)₂C₆H₃))] (**9**) at 50 % ellipsoid probability. Color scheme: Cu (cobalt blue), F (yellow-green), N (blue), O (red), K (pink). Solvent molecules, solid-state disorder, and hydrogen atoms are omitted for clarity.

X-band electron paramagnetic resonance (EPR) spectroscopy on a thawing 2-methyltetrahydrofuran solution of **7** displays a seven-line pattern, which was appropriately modeled with predominant isotropic copper hyperfine coupling (⁶³Cu A_{iso} = 87.4 MHz) and a minor contribution from nitrogen hyperfine coupling (¹⁴N A_{iso} = 17.8 MHz) (Figure 4a). At lower temperatures (4 K), the hyperfine coupling becomes anisotropic with copper hyperfine coupling values ⁶³Cu A_x = 9.3 MHz, A_y = 43.9 MHz, A_z = 9.7 MHz and ¹⁴N A_x = 86.6 MHz, A_y = 125.6 MHz, A_z = 130.3 MHz (Figure 4b). Akin to **7**, EPR spectra of both **8** and **9** are indicative of an S = 1/2 spin ground state which display seven-line patterns at room temperature in 2-methyltetrahydrofuran (Figure S–21 for **8**, S–24 for **9**). The large anisotropic nitrogen hyperfine coupling resolved at low temperatures (Figure S–22 for **8**, S–25 for **9**) is consistent with a significant contribution of the N 2p_x orbital to the now singly-occupied redox active molecular orbital (*vide infra*). Absorptions of near-infrared wavelengths (1000–2000 nm) were

detected for both the neutral (**5**, **6**) and anionic (**8**, **9**). This largely precludes assignment of any of these NIR bands as intervalence charge transfer (Figure S–60). This assertion is supported by TDDFT calculations, as no bands in the NIR or visible region present as intervalence charge transfer bands (IVCT; Figure S–72, Table S–1). Moreover, variable temperature UV–vis absorption measurements on **8** in 2-methyltetrahydrofuran indicate no substantial electronic rearrangements as a function of temperature (Figure S–59).




Figure 4. (a) Solution 2-methyltetrahydrofuran (9.843 GHz) EPR spectrum of **7** obtained at 298 K expanded to show an isotropic S = 1/2 signal (red) simulated (black) with the following parameters: g_{iso} = 2.033, σ_{gx} = 0.0277, σ_{gy} = 0.0039, σ_{gz} = 0.0051; ⁶³Cu A_{iso} = 87.4 MHz; ¹⁴N A_{iso} = 17.8 MHz. (b) Frozen 2-methyltetrahydrofuran (9.378 GHz) EPR spectrum **7** obtained at 4 K expanded to show an anisotropic S = 1/2 signal (red) simulated (black) with the following parameters: g_x = 1.979, g_y = 2.032, g_z = 2.074; σ_{gx} = 0.0080, σ_{gy} = 0.0014, σ_{gz} = 0.0098; ⁶³Cu A_x = 9.3 MHz, A_y = 43.9 MHz, A_z = 9.7 MHz; ¹⁴N A_x = 86.6 MHz, A_y = 125.6 MHz, A_z = 130.3 MHz.

2.4 X-ray Absorption Spectroscopy

Multi-edge X-ray absorption spectroscopy (XAS) comprising Cu K-edge, Cu L_{2,3}-edge, and N K-edge measurements were carried out to probe and differentiate the electronic structures of neutral **5**, **6** and anionic **8**, **9**. Cuprous (^{tBu}L)Cu(NCMe) (**10**) and cupric [(^{tBu}L)CuCl]₂ (**11**) complexes were prepared and measured as reference dipyrin-supported copper complexes. Previously, Lancaster and coworkers illustrated that high energy Cu K-edge pre-edge features (~8981 ± 0.5 eV), typically assigned as Cu 1s → 3d excitations, can instead be indicative of a ligand-based acceptor orbital with a more reduced Cu center in lieu of a high-valent Cu^{III} oxidation state assignment.²⁵ In the present instance, the presence of pre-edge features around ~8979 eV ± 0.5 eV traditionally assigned to a Cu^{II} center mirrors the case of TEMPO[•] coordinated to Cu^{II} dihalides, in which again the acceptor orbital is dominated by ligand character.²⁶ Moreover, pre-edge transitions at ~8980.5 eV have been observed in the Cu K-edge XAS spectrum of Cu^I species with low lying ligand π* orbitals, such as in Cu-bipyridine species.^{27,28} In summary, these data demonstrate Cu K-edge XAS pre-edge peak energies are not altogether diagnostic of physical oxidation states in these often-covalent systems; nonetheless, they remain useful to discuss to test the veracity of electronic structure calculations.

In scrutinizing the Cu K-edge XAS spectra of all species, a single pre-edge feature is observed at 8978.7 eV in the spectrum obtained for **11**, the authentic cupric reference. This feature is assigned as a Cu 1s → 3d transition consistent with the

expected Cu^{II} oxidation state assignment (Figure 5a). Similar pre-edge features are found in the Cu K-edge spectra of **5** and **6** at 8979.1 eV, although they appear at higher energy within the region conventionally ascribed to Cu^{II}. These features represent transitions into the LUMO of these complexes, which comprise an anti-bonding interaction between the N 2p_x of the imide and the weakly π -bonding Cu–Cu MO formed by the in-phase combination of Cu 3d_{x₂-y₂}. In contrast, the corresponding one-electron reduced anions **8** and **9** exhibit XANES spectra that are devoid of resolved pre-edge peaks (Figure 5b). These spectra have rising edges shifted to slightly lower energy, consistent with redox participation by Cu. Higher energy rising edge transitions observed at *ca.* 8983 eV gain intensity upon reduction from the neutral dicopper imido species. These intense rising edge features have been previously observed in Cu K-edge XAS spectra and are assigned primarily as Cu 1s \rightarrow 4p transitions that can contain contributions from ligand-to-metal charge transfer (LMCT) shakedown transitions.²⁹ Due to the dipole-allowed nature of an $s \rightarrow p$ ($\Delta l = +1$) transition, larger intensities are observed compared to the relative weak, lower energy Cu 1s \rightarrow 3d and Cu 1s \rightarrow L π^* features. Moreover, the increase in the intensity of these Cu 1s \rightarrow 4p rising-edge features is suggestive of greater Cu^I character. These spectra are consistent with diminished Cu character in low-lying acceptor orbitals upon reduction of **5** and **6** to yield **8** and **9**, respectively.

Cu L_{2,3}-edges provides entry for a more quantitative analysis of the composition of frontier orbitals. This analysis is possible because ligand orbital admixture into frontier Cu 3d-based orbitals modulates the allowedness and thus, intensity, of Cu 2p \rightarrow 3d transitions. An inverse relationship between the ligand orbital admixture coefficient and the summed L₃ (Cu 2p_{3/2} \rightarrow Cu 3d) and L₂ (Cu 2p_{1/2} \rightarrow Cu 3d) intensities is observed.³⁰ The L₃ mainline features occur at 931.3 eV (**4**) and 931.0 eV (**5**) while the L₃-edge of **11** occurs at 930.7 eV. Analysis of the summed L₃ + L₂ integrated area reveals 20 \pm 1 % and 24 \pm 1 % Cu 3d character (per Cu) in the LUMO orbitals of **5** and **6**, respectively (Figure 6b). These areas translate to 40% Cu 3d in the LUMO of **5** and 48% Cu 3d in the LUMO of **6**. Accounting for the two indistinguishable copper centers, these experimental values accord well with those calculated for the total Cu character (*vide infra*) at 40.0 % and 47.0 % for **5** and **6** respectively, representing a covalent ground-state electronic structure in which significant mixing of ligand character is expected. The Cu L_{2,3}-edge spectra of the reduced species show evidence of significant photo-damage, thus preventing a quantitative analysis of Cu character in the LUMO. Nonetheless, upon comparison of the first (and thus, least-damaged) scans of **8** and **9** (Figure 6b), the L₃ feature is significantly diminished in intensity as compared to **5** and **6**, and a pronounced satellite feature around 934.8 eV (iso-energetic with that observed in **10**) is apparent. The weakness of the absorbance features for **8** and **9** implicates a ligand-dominated SOMO and suggests that both anionic **8** and **9** contain reduced copper centers relative to the unambiguous cupric sample **11**.




Figure 5. Cu K-edge XAS spectra of (a) $[(^t\text{Bu}^{\text{dmx}})\text{Cu}_2(\mu^2\text{-NAr})]^n$ for $n = 0$, Ar: 3,5-(CF₃)₂-C₆H₃ (**6**, red); $n = -1$, Ar: 3,5-(F₃C)₂C₆H₃ (**9**, black); and mononuclear $(^t\text{Bu}^{\text{L}})\text{Cu}^{\text{I}}(\text{NCMe})$ (**10**, purple); $(^t\text{Bu}^{\text{L}})\text{Cu}^{\text{I}}\text{Cl}$ (**11**, gold); (b) $n = 0$, Ar: 4-MeOC₆H₄ (**5**, blue); Ar: 3,5-(F₃C)₂C₆H₃ (**6**, red); $n = -1$, Ar: 4-MeOC₆H₄ (**8**, green); Ar: 3,5-(F₃C)₂C₆H₃ (**9**, black).




Figure 6. Cu L_{2,3}-edge XAS spectra of (a) $[(t\text{Bu-dmx})\text{Cu}_2(\mu^2\text{-NAr})]^n$ for $n = 0$, Ar: 3,5-(CF₃)₂-C₆H₃ (**6**, red); $n = -1$, Ar: 3,5-(F₃C)₂C₆H₃ (**9**, black); and mononuclear $(t\text{BuL})\text{Cu}^{\text{I}}(\text{NCMe})$ (**10**, purple); $(t\text{BuL})\text{Cu}^{\text{II}}\text{Cl}$ (**11**, gold); (b) $n = 0$, Ar: 4-MeOC₆H₄ (**5**, blue); Ar: 3,5-(F₃C)₂C₆H₃ (**6**, red); $n = -1$, Ar: 4-MeOC₆H₄ (**8**, green); Ar: 3,5-(F₃C)₂C₆H₃ (**9**, black).

N K-edge XAS was obtained to assess the redox participation of the aryl nitrenoid moiety. Multiple features assigned as transitions from N 1s to antibonding MOs comprised of N and aryl character from the dipyrin ligand are seen along the rising edge of the N K-edge for all species, consistent with our prior measurements of mononuclear dipyrin-supported Fe imide/imine complexes³¹ as well as for analogous dipyrin-supported nickel complexes³² (Figure 7).

Additional, weak features at *ca.* 395-396 eV have been previously assigned as excitations into subvalent NAr-based MOs given the aforementioned Fe/Ni imine³¹,³² and copper nitrene⁹ precedent. These assignments are supported by electronic structure calculations (*vide infra*). In **5** and **6**, these transitions are observed at 396.0 eV and are assigned as transitions into a MO comprised of both Cu 3d_{x²-y²} and NAr 2p character. Upon one-electron reduction, these transitions are shifted to lower energy at 395.2 eV, and the intensities are diminished.

The diminished intensity of the 395.2 eV features in **8** and **9** implicates an MO containing NAr 2p_x character participating in the **5/8** and **6/9** redox couples.




Figure 7. N K-edge XAS spectra of (a) $[(t\text{Bu-dmx})\text{Cu}_2(\mu^2\text{-NAr})]^n$ for $n = 0$, Ar: 3,5-(CF₃)₂-C₆H₃ (**6**, red); $n = -1$, Ar: 3,5-(F₃C)₂C₆H₃ (**9**, black); and mononuclear $(t\text{BuL})\text{Cu}^{\text{I}}(\text{NCMe})$ (**10**, purple); $(t\text{BuL})\text{Cu}^{\text{II}}\text{Cl}$ (**11**, gold); (b) $n = 0$, Ar: 4-MeOC₆H₄ (**5**, blue); Ar: 3,5-(F₃C)₂C₆H₃ (**6**, red); $n = -1$, Ar: 4-MeOC₆H₄ (**8**, green); Ar: 3,5-(F₃C)₂C₆H₃ (**9**, black).


2.5 Nitrenoid reactivity

Transition metal bound imido complexes facilitate a range of nitrene transfer chemistry.^{33,34} Electron-rich metal imide complexes have been observed to undergo transfer to nucleophiles (e.g., tertiary phosphines, isocyanides, CO), whereas electrophilic imide complexes can exhibit olefin aziridination or the amination of C–H bonds.³⁵⁻⁴⁰ To establish the effect of the dinuclear cuprous proximal arrangement afforded by the dimethylxanthene spacer, **10** was treated with substoichiometric aryl azides (0.5 equiv) at elevated temperatures (60 °C, C₆D₆) to promote acetonitrile displacement, re-


sulting in quantitative formation of azoarene and accompanied by formation of dimeric cuprous species ($^t\text{Bu}^-\text{L})_2\text{Cu}_2$ (**18**) as the major copper-containing species. Complex **18** displays substantial pyrrole torsion with respect to each pyrrole plane ($27.0(1)^\circ$, $30.9(1)^\circ$) to accommodate an exceptionally short intermetallic distance ($2.443(2)$ Å, Figure S–88). Complex **18** was similarly competent for azoarene formation from aryl azides at elevated temperatures. The absence of spectroscopically observable intermediates prior to azoarene formation corroborates the paramount functionality of the dimethylxanthene spacer in stabilizing $[(^R\text{dmx})\text{Cu}_2(\mu^2\text{-NAr})]$ complexes against rapid azoarene formation.

The neutral $(^R\text{dmx})\text{Cu}_2(\mu^2\text{-NAr})$ complexes (**2**, **3**, **5**, **6**) all undergo nitrene transfer upon exposure to nucleophiles. For example, treatment of the neutral complexes with excess tertiary phosphine (PMe_3 , **2** or **3**; PPh_3 , **5** or **6**) afforded the corresponding phosphinimide products with concomitant formation of the phosphine ligated $[(^R\text{dmx})\text{Cu}_2(\text{PR}_3)_2]$ products, $(^{\text{Mes}}\text{dmx})\text{Cu}_2(\text{PMe}_3)_2$ (**12**), $(^t\text{Bu}^-\text{dmx})\text{Cu}_2(\text{PPh}_3)_2$ (**13**), in good yields (Scheme 5). The phosphine-ligated products were verified by independent synthesis and characterized by single crystal X-ray diffraction (Figures S–47, S–58). The electron-deficient imide complexes $(^R\text{dmx})\text{Cu}_2(\mu^2\text{-N}(3,5\text{-}(\text{F}_3\text{C})_2\text{C}_6\text{H}_3))$ (**3**, **6**) reacted rapidly with excess *tert*-butyl isocyanide (CN^tBu) to afford the corresponding carbodiimide product. In contrast, the reaction of electron-rich imidos $(^R\text{dmx})\text{Cu}_2(\mu^2\text{-N}(\text{C}_6\text{H}_4\text{OMe}))$ (**2**, **5**) with CN^tBu yielded the resulting azoarene product (Scheme 6). The copper-containing species from these reactions was identified as $(^R\text{dmx})\text{Cu}_2(\text{CN}^t\text{Bu})_2$ (**R**: Mes, **14**; ^tBu , **15**) and confirmed by independent synthesis and single crystal X-ray diffraction. The difference in reactivity profiles can be attributed to varying radical-localization at the nitrogen bridgehead (*vide supra*).

Scheme 5




Scheme 6



While imido complexes **2**, **3**, **5**, and **6** were thermally stable (120°C , 24 h), azoarene generation from the imido complexes was promoted by treatment of the imide complexes with pyridine, 4-(dimethylamino)pyridine (DMAP), or cyclohexene. For example, **2** decays as a first-order process in the presence of excess DMAP (9 equiv.) with heating (65°C , 10 h) to afford $(^{\text{Mes}}\text{dmx})\text{Cu}_2(\text{dmap})_2$ (**16**) and the corresponding azoarene in excellent yield (87 %) by both ^{19}F and ^1H NMR with no spectroscopically observable intermediates. In the presence of weak H-atom donors such as diphenylhydrazine, treatment of

2 or **3** with DMAP affords the corresponding aniline as well as **16** under more mild conditions (35°C , 6 h).

Scheme 7



None of the $(^R\text{dmx})\text{Cu}_2(\mu^2\text{-NAr})$ complexes (**2**, **3**, **5**, **6**) exhibited propensity towards H-atom abstraction at room temperature, even upon exposure to weak H-atom sources (e.g., 2-hydroxy-2-azaadamantane; $\text{BDE}_{\text{O-H}} = 69$ kcal mol⁻¹). However, the anion **7** was observed to intramolecularly aminate at room temperature. Allowing solutions of **7** to stand at room temperature over several minutes generated several new diamagnetic species by ^1H NMR, with one species identified as a di(cuprous) product in which intramolecular amination of the proximal mesityl C–H bond has occurred (**17**, Scheme 7). Single crystals of the reaction product **17** were grown by vapor diffusion of diethyl ether into a concentrated solution of **17** in tetrahydrofuran. The molecular structure obtained by crystallography, while poor, was sufficient to establish the connectivity (Figures S–41). This reaction proceeds through formal loss of a single H-atom, which may contribute to the presence of multiple unidentified species in crude ^1H NMR spectrum. Attempts to mitigate ligand functionalization by introduction of external weak C–H or O–H bonds to **7** were unsuccessful.

In contrast to the reactivity exhibited by the $(^R\text{dmx})\text{Cu}_2(\mu^2\text{-NAr})$ complexes (**2**, **3**, **5**, **6**) complexes and the anion **7**, the more thermally robust imido anions $[(^t\text{Bu}^-\text{L})\text{Cu}_2(\mu^2\text{-NAr})]^-$ **8** and **9** were unreactive toward triphenylphosphine or weak C–H bonds and H-atom donors. Though containing comparable % N 2p covalency values (**5/8**: 15 %/22 % vs. **6/9**: 17%/25%), formation of the corresponding iminyl upon reduction ultimately results in most of the spin density of the frontier RAMO being delocalized over the aryl ring of the NAr substituent.

3. Discussion

3.1 Synthesis and structure.

The pacman bis(dipyrrin) dicopper complexes are well suited to undergo nitrene capture from aryl azides. While the dicuprous precursors do not exhibit substantial intercupper interactions (e.g., **13**, $d_{\text{Cu-Cu}}$: > 6.20 Å), the two-electron oxidized imido complexes have shorter interactions ($d_{\text{Cu-Cu}}$: $2.82 - 2.87$ Å) enforced by the bridging imido motif. The Cu–Cu separation is shortened compared to the bridging aryl imido complexes reported by Warren ($2.911(1)$ Å)⁵ but longer than the isoelectronic dicopper carbene complexes ($2.464(1)$ Å).⁴¹ The Cu–N_{imide} distances are consistent with reported μ^2 -imido ($1.794(5) - 1.808(5)$ Å)^{5–8} and μ^3 -copper imidos ($1.842(5) - 1.893(6)$ Å).^{42,43} Upon reduction, each of the crystallographically characterized complexes **7–9** exhibit expansion of the $[\text{Cu}_2\text{N}]^{1+}$ core, consistent with increased electron richness. The near-identical coordination environment about each copper ion in **7–9** contrasts the apparent valence-localized oxo-bridged $\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}$ complex previously isolated within this framework.¹³ Each of the reduced nitrenoid complexes also features a contracted N_{imido}–C_{ipso} bond (**7**, $1.318(4)$ Å; **8**, $1.351(3)$ Å; **9**,

1.342(3) Å) which approximate as C–N double bonds akin to those found in pyridine (Table 1). Thus, the nitrenoid units take on more of a ketamide resonance form similar to those previously reported for the bimolecularly coupled $[(^{\text{EMind}}\text{L})\text{Cu}]_2(\text{N}(3,5\text{-(MeO)}_2\text{C}_6\text{H}_3)_2)_2$ ($\text{N}_{\text{imido}}\text{--C}_{\text{ipso}}$ bond lengths: 1.257(5) Å, 1.276(6) Å).⁹ Furthermore, redox non-innocence behavior of the aryl fragment is apparent upon inspection with elongation of the $\text{C}_{\text{ipso}}\text{--C}_{\text{ortho}}$ bond upon reduction. Such arene distortions have been observed in previously isolated dipyrin-supported quintet iron iminyl complexes,^{31,44,45} doublet nickel iminyl complexes,³² and triplet copper nitrene complexes,⁹ suggesting a delocalized $^2[\text{NAr}]$ vacancy is potentially engendered upon one-electron reduction. Such a transformation is unusual, as the reduction of **2** proceeds with oxidation of the most electronegative atom of the $[\text{Cu}_2(\mu^2\text{--NAr})]$ core.

3.2 Calculated X-ray Absorption Spectra

Electronic structure calculations were carried out to facilitate discussion of the XAS data and to interrogate the nature of the redox event linking the neutral and anionic dicopper nitrenoid complexes. The electronic structures that emerge, at least for the neutral species **5** and **6**, are not altogether straightforward. Considering each dipyrin-Cu center in isolation, the highest Cu 3d-based orbital in each Cu-dipyrin fragment is predicted to be $3d_{x^2-y^2}$ (Figure 8). This affords a basis for two linear combinations representing weakly interacting $\text{Cu}_2 \pi$ and π^* orbitals. Introduction of the NAr $2p_x$ perpendicular to the Ar plane produces a 3-center MO framework representative of a 3-center-4-electron bond. This simple picture predicts that the redox-active molecular orbital (RAMO) is best described as $\psi^* \approx (\text{Cu}_2 \pi - \text{NAr } 2p_x)$, an overall π^* MO, in complete accord with observations from XAS. The Cu orbitals are depressed compared to those of the nitrenoid motif, resulting in an inverted ligand field with NAr $2p_x$ character dominant in the RAMO. We note that this is an idealized picture that assumes C_{2v} symmetry and neglects Cu–Cu interactions—in fact the two LCu inner coordination spheres do not perfectly overlap as would be required for this picture to be exactly operative; each complex exhibits significantly non-zero dihedral angles defined by the two Cu and two dipyrin *meso*-C atoms. This deviation from ideal symmetry becomes apparent in scrutinizing orbitals derived from electronic structure calculations (*vide infra*).




Figure 8. Simplified qualitative frontier molecular orbital framework in idealized C_{2v} symmetry showcasing $\text{Cu}_2\text{--NAr } \pi$ bonding in **5/6**, **8/9**.

Scalar relativistic single point DFT calculations employing the B3LYP hybrid density function with the CP(PPP) basis set on Cu and the scalar-relativistically recontracted ZORA-def2-TZVP(-f) basis set on all other atoms were carried out on all compounds for which XAS data were obtained. For neutral **5** and **6**, broken symmetry (BS) singlet diradical (BS 1,1) solutions were found to have lower energy relative to corresponding conventional unrestricted Kohn-Sham (UKS) solutions. For **8** and **9**, no BS solutions were obtained. Rather, standard doublet ($S = 1/2$) solutions emerged.

The aforementioned single point DFT solutions were used as starting points for time-dependent DFT (TDDFT) calculations of Cu and N K-edge XAS. TDDFT calculations are now well-established as a means to produce simulated XAS pre-edge data with high fidelity to experiment after proper calibration.^{27,46–48} The present compounds do not present an exception to this trend, however the underlying electronic structure interpretation requires extended explanation.

Upon application of a linear energy correction produced via correlation of TDDFT-calculated with experimental pre-edge peak energies for all compounds under discussion, the Cu K pre-edge and rising edge peak energies accord well with experiment (Figure S-61, S-63). These calculations were then used to assign acceptor MOs underlying given transitions. For **10**, the observed 8982.4 eV transition is predicted at 8982.1 eV as a $\text{Cu } 1s \rightarrow 4p$ excitation, where the acceptor MO exhibits some acetonitrile C–N π^* character. These calculations predict a weak $\text{Cu } 1s \rightarrow \text{LUMO}$ feature at 8980.6 eV (dipyrin π^* orbital) that is not resolved experimentally. The calculated spectrum of **11** reproduces the 8987.7 eV $\text{Cu } 1s \rightarrow 3d$ feature while also predicting an unresolved $\text{Cu } 1s \rightarrow \text{L } \pi^*$ transition at 8983.4 eV. The 8979.0 eV $\text{Cu } 1s \rightarrow \text{RAMO}$ transitions are reproduced for **5/6**, and upon inspection of the calculated spectra for **8/9** weak $\text{Cu } 1s \rightarrow \text{RAMO}$ transitions are observed at slightly higher energy relative to their neutral counterparts (8979.5 and 8979.9 eV respectively). In addition, **8** and **9** display weak pre-edge features at 8981.1 eV corresponding to $\text{Cu } 1s \rightarrow \text{L } \pi^*$ transitions, shifted to lower energy compared to the 8982.9 eV features predicted for the neutral complexes. The diminished intensity of the $\text{Cu } 1s \rightarrow 3d$ transitions and observed energy shifts supported the contention that **8** and **9** possess RAMOs with diminishing copper character and N-localized spin. Fewer acceptor holes are available in NAr, and participation of NAr in the redox is supported by the energetic red shift, indicating a more electron-rich NAr fragment.

Agreement between experiment and theory is maintained upon returning to the N K-edge XAS spectra (Figure S-64, S-65). Examining the spectrum of **10**, the 398.0 eV feature in the experimental N K-edge is predicted at 397.4 eV as a transition into the dipyrin-based LUMO, while the 398.9 eV feature is predicted as a combination of transitions into higher energy MOs of Cu 4p and acetonitrile π^* parentage. For **11**, the observed 397.2 eV and 398.0 eV lower-energy shoulders that represent transitions into the predominantly Cu 3d SOMO, containing mixing with Cl 3p and dipyrin N 2p, and lower-lying dipyrin π^* orbitals, are predicted at 396.3 eV and 397.8 eV. Lastly, examining the N K-edge XAS spectra of the di(copper) species, the 396.0 eV feature of **5** is predicted at 396.3 eV as a transition into the RAMO containing 47% Cu and 30% NAr (15% N 2p). The lower energy 395.2 eV feature

of **8** is once again reproduced, predicted at 395.6 eV as a transition into the RAMO that contains 34 % Cu and 44 % NAr (17 % N $2p$). Similar results were obtained for **6/9** (Figure S-64, S-65).

Unfortunately, quantification⁴⁸ of N $2p$ covalency from the given spectra was unreliable due to photodamage and hence rapid reduction in the intensity of the 396.0 eV and 395.2 eV features that is seen upon carrying out multiple scans at the same sample location. However, returning to the Cu $L_{2,3}$ -edge data, the ca. 40% and 47% overall Cu 3d participation in the LUMOs of **5** and **6**, respectively, accord with the 40% Cu 3d and 47% Cu 3d predicted by DFT. Together with anisotropic ^{14}N hyperfine coupling manifesting in the frozen-solution EPR data obtained for **6** and **9** as well as the well-resolved NAr features in the corresponding N K-edge XAS spectra, these support the contention that there is a high degree of NAr $2p$ character in the RAMO of the **5/8** and **6/9** couples.

3.3 *Ab Initio* Calculations

While the TDDFT calculations of the XAS data exhibit high fidelity to experiment, a paradoxical picture emerges when more deeply probing the electronic structure of neutral complexes **5** and **6**, for which BS(1,1) solutions are encountered via DFT. Examination of the acceptor MOs used to generate the TDDFT-calculated spectra following a quasi-restricted orbital (QRO) transformation reveals that they are $[\text{Cu}_2 \pi - \text{NAr } 2p] \pi^*$, as predicted qualitatively (Figure S-69). However, these are the canonical orbitals generated during the BS calculation. Inspection of the *unrestricted corresponding orbitals* (UCOs), which reflect the *magnetic* orbitals of the singlet diradical and are mandatory for TDDFT analysis⁴⁹ presents a problem (Figure S-69). These occupied spin orbitals are predicted to have diminished overlap ($S = 0.74$ and $S = 0.56$ for **5** and **6**, respectively), and overall are antiferromagnetically coupled with J -values of -2631 cm^{-1} and -1802 cm^{-1} for **5** and **6**, respectively in accord with the experimentally-established diamagnetic natures of these complexes. However, these *occupied* UCOs are effectively of the same nature as the *unoccupied* canonical MOs involved in the XAS pre-edge excitations.

This paradox plausibly arises due to the multiconfigurational nature of the ground state electronic structures of **5** and **6**. *Ab initio*, complete active space self-consistent field (CASSCF) calculations were carried out on these complexes were carried out these complexes using CAS(10,9) active spaces.⁵⁰ These active spaces were chosen to balance flexibility with computational expense. Truncated structural models **5'** and **6'** were generated by removing $t\text{Bu}$ substituents from the pyrrole rings and optimizing H-atom positions for similar reasons of economy. Hybrid DFT-calculated orbitals were used as starting points for the CASSCF calculations, with orbitals included in the active space comprising those involved in the $\text{Cu}_2\text{-NAr } \pi$ interaction as well as some dipyrin MOs. These calculations were carried out using the ZORA-def2-TZVP(-f) basis on Cu and N, with ZORA-def2-SVP on all other atoms. For all cases, state averaging was used to facilitate convergence, with five singlet and five triplet roots calculated. The strongly contracted N-electron valence perturbation theory 2 (NEVPT2) correction was applied to CASSCF state energies to account for dynamical electron correlation. The calculated singlet-triplet gaps after NEVPT2 correction are 1341.3 and 808.6 cm^{-1} for **5'** and **6'** in line with experimental magnetic behavior. A molecular orbital diagram reflecting the leading ground state configurations for **6'** resulting from the


CASSCF calculation is given in Figure 9. The analogous diagram for **5'** is qualitatively similar and is included in the supporting information (Figure S-68a). The ground state singlet comprises two leading configurations with weights of ca. 50% and 40%. These configurations reflect either double occupation of the $\text{Cu}_2 \pi^*$ MO or the $(\text{Cu}_2 \pi - \text{NAr } 2p_x) \pi^*$ RAMO, respectively. These configurations can be effectively be averaged to yield the BS picture.

A qualitative frontier MO diagram for **9** is given in Figure 9b for comparison to **6**. The diagram for **8** is similar and is included as figure S-68b. These single-configurational doublet states possess an unpaired electron in the $(\text{Cu}_2 \pi - \text{NAr } 2p_x) \pi^*$ RAMO in accord with predictions and the observed decreased intensities of the N K pre-edge peaks upon in the monoanions relative to the neutral analogues.

3.4 Electronic structure of the bridged imidos

All of the dicopper imido complexes (**2**, **3**, **5**, and **6**) feature highly multiconfigurational singlet ground states (Figure 9). The singlet configurations emerging from these calculations are consistent with the diamagnetic electronic configurations observed at room temperature as assessed by multinuclear NMR and a lack of any discernible EPR signal. Via CASSCF analysis, we assess the most consistent electronic description for all these complexes comprise singlet ground states whose two major configurations are defined by (a) double population of $(\text{Cu-Cu}) \pi^*$ with no contribution from the imido unit (48%); and (b) double population of a $[(\text{Cu}_2 \pi - (\text{NAr})) \pi^*]$ interaction contributing 40%. Configuration 1 illustrates how the cupric character in the dicopper imidos is diminished, consistent with the XAS data, (Figure 9a).

The formally mixed valence anionic imidos (**7-9**) possess doublet ground states exhibiting well-defined EPR signals between 4–295 K (Figures S-21-22, 24-25). Solution-phase EPR spectra display isotropic hyperfine coupling (i.e., electronic contribution from the $2s$ orbital), dominated by Cu character with minimal N contribution and consistent with spectra observed for fully delocalized, Class IIIA mixed valence dicopper complexes⁵¹⁻⁵³ in the Robin-Day classification.⁵⁴ This observation is rationalized by considering the nitrogen contribution to the RAMO arises from the $2p_x$ component, leading to an anisotropic component only apparent at lower temperatures.⁵⁵ In the frozen solution EPR spectrum, a high degree of unpaired electron localization to N $2p_x$ is apparent, contributing to the hyperfine coupling alongside the copper centers. The temperature-dependence of the EPR spectra are superficially inconsistent with a Class IIIA mixed valence assignment and more reminiscent of Class II species,⁵⁶⁻⁵⁹ but are accounted for when considering that the temperature dependence arises not from changes in electronic structure but from the participation of anisotropic hyperfine contributions.⁵⁵ While complexes **7-9** do feature optical transitions in the near IR region (Figure S-59), their corresponding neutral precursors also feature prominent transitions at similar energies. Indeed, both $(t\text{Bu-dmx})\text{Cu}_2(\mu^2\text{-N}(\text{C}_6\text{H}_4\text{OMe}))$ (**5**) and its anionic analogue **8** feature absorptions at the same energy ($\sim 1300 \text{ nm}$) with near equal intensity ($\epsilon \sim 500 \text{ M}^{-1}\text{cm}^{-1}$), whereas the NIR band for $[(t\text{Bu-dmx})\text{Cu}_2(\mu^2\text{-N}(3,5\text{-}(\text{F}_3\text{C})_2\text{C}_6\text{H}_3))]^-$ (**9**) does shift to lower energy (1750 nm) than its neutral precursor (**6**, 1450 nm).



The clearest picture of the electronic structure for the anionic imidos arises from the combined Cu ($L_{2,3}$, K-edges) and N (K-edge) XAS spectra. While neutral imidos **5** and **6** feature prominent pre-edge absorptions in the Cu K-edge spectra, the anionic imidos (**8**, **9**) lack clear experimental pre-edge features (Figure 5, inset) similar to cuprous standard (^{tBu}L)Cu^I(NCMe) (**10**). The weak pre-edge features observed in TDDFT calculations can be attributed to addition of an electron to the Cu and N based LUMO in **5/6**. Both neutral **5** and **6** exhibit intense L_3 main line features at energies (~930 eV) consistent with mononuclear analogues (^{tBu}L)Cu^{II}Cl, (**11**), while the L_3 feature is significantly diminished in the reduced nitrenoids with prominent satellites at higher energy (935 eV) (Figure 6b) akin to the mononuclear cuprous analogue (^{tBu}L)Cu^I(NCMe) (**10**). Lastly, the N K-edge reveals a systematic shift of the pre-edge absorptions as the singly-occupied molecular orbital becomes more localized on the (NAr) fragment, suggesting increased iminyl radical character exists for the anions **8** and **9**. In addition, the decreased intensity of the N K-edge pre-edge feature is indicative of population of the N

The foregoing data indicate the reduced $[\text{Cu}_2(\mu^2\text{-NAr})]$ complexes largely exhibit iminyl character. The copper centers in the anionic imido complexes exhibit diminished Cu character in acceptor orbitals based on the Cu $L_{2,3}$ -edge data and Lowdin spin density analysis. These data together are typical of bridged Cu complexes exhibiting highly covalent cores with substantial ligand participation in the electroactive orbitals.⁶³⁻⁶⁷ The RAMO of **8** and **9** display comparable Cu character to other mixed valence Cu species such as Cu_A (44 % Cu),⁶⁷ $[(\text{L}^{\text{Prdaco}}\text{S}(\text{Cu})_2)^+]$ (38 % Cu),⁶⁷ and $[(\text{I}^{\text{Bup}}\text{PNP}(\text{Cu})_2)^+]$ (24 % Cu).⁶³ Nonetheless, we note **8** and **9** are lower coordinate (tri-coordinate instead of tetra-coordinate) and contain strictly hard nitrogen atom in the primary coordination sphere in lieu of polarizable sulfur and phosphorous. Indeed, the N K-edge data reveal common features between the reduced imidos **8** and **9** with other *N*-centered, radical species.^{9,31,32} Whereas the previously reported metal-supported iminyl structures exhibit bond-length distortions within the iminyl fragment consistent with hole delocalization, complexes **8** and **9** do not exhibit substantial distortions aside from enhanced N-C_{ipso} π -bonding upon NAr reduction. The electronic description is corroborated with the calculated electronic structure which reveals a highly (NAr)-centered RAMO (Figure 9b). Insight into the increasing cuprous character can be gleaned from comparing the frontier orbitals of **6** and **9**. While chemical reduction of **6** occurs by population of the LUMO, the ground state of imido **6** is multiconfigurational with nearly equal contributions from (Cu-Cu) π^* (213) and $[(\text{Cu}_2 \pi - (\text{NAr})) \pi^*]$ (214) singlet configurations. The frontier orbital configuration of **9** leads to double population of (Cu-Cu) π^* (which was only partially occupied in **6**), and partial population of $[(\text{Cu}_2 \pi - (\text{NAr})) \pi^*]$ (Figure 9b) – effectively reducing the overall Cu₂-N bond order by 1.5. Thus, the bridging ligand radical character was quite remarkably achieved through reduction, not oxidation, of the $[\text{Cu}_2(\mu^2\text{-NAr})]$ core.

3.5 Reactivity of the imido complexes

The neutral imidos (**2**, **3**, **5**, **6**) exhibit reaction chemistry reminiscent of the electrophilic imido complexes. The nitrene motif can be transferred to incoming nucleophiles (e.g., tertiary phosphines, alkyl isocyanides), but the aryl-substituent on the imide influences the reaction trajectory. The more electrophilic $[\text{Cu}_2(\mu^2\text{-N}(3,5\text{-}(\text{F}_3\text{C})_2\text{C}_6\text{H}_3))]$ imido complexes **3** and **6** undergo clean nitrene transfer to nucleophilic reagents (PR_3 , CN^tBu), while the $[\text{Cu}_2(\mu^2\text{-N}(\text{C}_6\text{H}_3\text{OMe}))]$ imido complexes **2** and **5** produce azoarene from nitrene coupling upon reaction with $^t\text{BuNC}$. Several simple ligands (e.g., pyridine, DMAP) also can induce azoarene formation at elevated temperatures, which do not require direct attack of substrate on the imido fragment. Indeed, each of the nitrene transfer reactions observed can occur from substrate attack on $[\text{Cu}_2(\mu^2\text{-NAr})]$ core, potentially activating the core via ligation (e.g., $[\text{Cu}_2(\mu^2\text{-NAr})(\text{L})]$, L = substrate) or inducing Cu-N cleavage to form a transient terminal imido complex. The activated dicopper core can then undergo rapid bimolecular coupling or undergo free nitrene expulsion. However, the complete selectivity of nitrene coupling to yield azoarene argues against free nitrene expulsion without the detection of free nitrene products (e.g., aziridine, C-H amination, nitrene aryl-ring expansion). The observed difference in reactivity between electron rich and electron poor imido complexes may arise from the relative stabilities of the triplet and singlet nitrene states on the transient $[\text{Cu}(\text{L})\text{Cu}(\text{NAr})]$ (L = PR_3 , CN^tBu) formed. The Cu-bound $^3[\text{N}(\text{C}_6\text{H}_4\text{OMe})]$ would favor reacting with triplet configuration as opposed to added singlet reactants. The absence of intramolecular benzylic amination, intermolecular H-atom abstraction, or ring expansion of the nitrene aryl ring argue against dissociation of free nitrene in solution. Nonetheless, the release of singlet nitrene cannot be fully discounted, noting that whereas $^1[\text{N}(3,5\text{-}(\text{F}_3\text{C})_2\text{C}_6\text{H}_3)]$ may have an appreciable singlet lifetime in solution, intersystem crossing from $^1[\text{N}(4\text{-MeOC}_6\text{H}_3)]$ to a triplet ground state is exceptionally fast,⁶⁸ perhaps explaining the discrepancies in reactivity upon CN^tBu treatment.

Aside from the intramolecular amination exhibited by thermally unstable **7**, the reduced imido complexes show no propensity for nitrene transfer to substrate or proclivity towards H-atom abstraction chemistry. These results suggest the presence of radical character at the nitrogen bridgehead of the imide motif enables amination of proximal ligand, benzylic C-H bonds at room temperature. Despite the presence of *bona fide* radical character at the bridgehead, **7** fails to perform intermolecular amination chemistry. This observation may be attributed to the relatively reducing potential of **7**, rendering the resulting N-H bond excessively weak. Moreover, the inability of the ligand to support a zero-valent copper center renders two-electron reduction chemistry from **7** unlikely. The dipyrin ligand substituents (Mes in **7**, ^tBu in **8** and **9**) may sterically shield the iminyl fragment from incoming substrate, especially given the SOMO character is localized on the N_{px} orbital normal to the $[\text{Cu}_2(\mu^2\text{-NAr})]$ plane. Thus, the amination exhibited by **7** may result from the optimally oriented, mesityl benzylic methyl substituents, while the C-H bonds of the ^tBu groups of **8** and **9** are too strong to be activated by the (^2NAr) and sterically restrict access to the $[\text{Cu}_2(\mu^2\text{-NAr})]$ core. Furthermore, the anionic complexes **8** and **9** are less electrophilic and show diminished reactivity observed for the neutral analogues.

4. Conclusions

Neutral imido complexes of the type $(^R\text{dmx})\text{Cu}_2(\mu^2\text{-NAr})$ (R : Mes, ^tBu ; Ar : 4-MeOC₆H₄, 3,5-(F₃C)₂C₆H₃) were synthesized by treatment of the corresponding dicuprous precursors with aryl azides. Chemical reduction of the imido complexes afforded dicopper iminyl products as determined by EPR, Cu/N XAS, crystallography, and computations. The iminyl ligand is distinct from previously reported subvalent nitrogen motifs which arise from radical delocalization following nitride,^{69,70} imide,^{31,44,45,71} or amide⁷²⁻⁷⁵ oxidation.^{72,76} The iminyl complexes reported herein show modest disruption of the aryl units, arising from population of the NAr lowest-lying π^* orbital. The iminyl density is predominantly localized on the N $2p_x$ orbital, normal to the $[\text{Cu}_2(\mu^2\text{-NAr})]$ plane, optimally located to aminate the benzylic C-H bonds of the ($^{\text{Mes}}\text{dmx}$) ligand but remain sterically occluded in the ($^t\text{Bu}\text{dmx}$) framework and are thus robust. Despite the presence of *bona fide* radical character on the bridging iminyl, the $[(^t\text{Bu}\text{dmx})\text{Cu}_2(\mu^2\text{-NAr})]^-$ complexes are resistant to nitrene transfer and H-atom abstraction reactivity. Thus, while we were successful in generating dicopper complexes with substantive radical character accumulation at the bridgehead ligand, these complexes were found to be remarkably stable. Combined, our results contribute understanding of electronic structure, and identifying the precise localization of frontier orbital radical density, can aid in understanding group transfer reactions such as C-H bond functionalization and nitrene insertion.

ASSOCIATED CONTENT

The Supporting Information is available free of charge on the ACS Publications website, including synthesis of **1-18**, spectroscopic characterization, computational details, CheckCIF results, solid-state molecular structures, and supplementary spectra.

. AUTHOR INFORMATION

Corresponding Author

*kml236@cornell.edu

*betley@chemistry.harvard.edu

ORCID

Kurtis M. Carsch: 0000-0003-4432-7518

Ida M. DiMucci: 0000-0003-3440-7856

Diana A. Iovan: 0000-0001-9889-7183

Shao-Liang Zheng: 0000-0002-6432-9943

Kyle M. Lancaster: 0000-0001-7296-128X

Theodore A. Betley: 0000-0001-5946-9629

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

We thank E. Johnson (Harvard) for helpful discussions and the initial synthesis of ($^{\text{Mes}}\text{dmx}$)H₂. We thank S. Sproules (Glasgow) for helpful guidance with EPR interpretation. T.A.B. gratefully acknowledges support by grants from NIH (GM-115815), the Dreyfus Foundation in the form of a Teacher-Scholar Award, and Harvard University. K.M.C. acknowledges the Fannie & John Hertz Foundation and the National Science Foundation for financial support of this research. XAS data were obtained at SSRL, which is supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences under Contract No. DE-AC02-76SF00515. The SSRL Structural Molecular Biology Program is supported by the Department of Energy's Office of Biological and Environmental Research, and by NIH/HIGMS (including P41GM103393). ChemMatCARS Sector 15 is principally supported by the National Science Foundation/Department of

Energy under grant number NSF/CHE-0822838. Use of the Advanced Photon Source was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract No. DE-AC02-06CH11357. Collection of low-temperature UV/Vis spectra were obtained at the Center for Nanoscale Systems (CNS, Harvard University), a member of the National Nanotechnology Coordinated Infrastructure Network (NNCI), which is supported by the National Science Foundation under NSF award no. 1541959.

References

- (1) Labinger, J. A.; Bercaw, J. E., Understanding and Exploiting C–H Bond Activation. *Nature* **2002**, *417*, 507.
- (2) Godula, K.; Sames, D., C–H Bond Functionalization in Complex Organic Synthesis. *Science* **2006**, *312*, 67.
- (3) Bergman, R. G., C–H Activation. *Nature* **2007**, *446*, 391.
- (4) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K., C–H Bond Functionalization: Emerging Synthetic Tools for Natural Products and Pharmaceuticals. *Angew. Chem. Int. Ed.* **2012**, *51*, 8960.
- (5) Badiei, Y. M.; Krishnaswamy, A.; Melzer, M. M.; Warren, T. H., Transient Terminal Cu–Nitrene Intermediates from Discrete Dicopper Nitrenes. *J. Am. Chem. Soc.* **2006**, *128*, 15056.
- (6) Badiei, Y. M.; Dinescu, A.; Dai, X.; Palomino, R. M.; Heinemann, F. W.; Cundari, T. R.; Warren, T. H., Copper–Nitrene Complexes in Catalytic C–H Amination. *Angew. Chem. Int. Ed.* **2008**, *47*, 9961.
- (7) Aguila, M. J. B.; Badiei, Y. M.; Warren, T. H., Mechanistic Insights into C–H Amination Via Dicopper Nitrenes. *J. Am. Chem. Soc.* **2013**, *135*, 9399.
- (8) Bakhoda, A.; Jiang, Q.; Bertke, J. A.; Cundari, T. R.; Warren, T. H., Elusive Terminal Copper Arylnitrene Intermediates. *Angew. Chem. Int. Ed.* **2017**, *56*, 6426.
- (9) Carsch, K. M.; DiMucci, I. M.; Iovan, D. A.; Li, A.; Zheng, S.-L.; Titus, C. J.; Lee, S. J.; Irwin, K. D.; Nordlund, D.; Lancaster, K. M.; Betley, T. A., Synthesis of a Copper-Supported Triplet Nitrene Complex Pertinent to Copper-Catalyzed Amination. *Science* **2019**, *365*, 1138.
- (10) Groothaert, M. H.; Smeets, P. J.; Sels, B. F.; Jacobs, P. A.; Schoonheydt, R. A., Selective Oxidation of Methane by the Bis(μ -oxo)Dicopper Core Stabilized on ZSM-5 and Mordenite Zeolites. *J. Am. Chem. Soc.* **2005**, *127*, 1394.
- (11) Vanelderen, P.; Hadt, R. G.; Smeets, P. J.; Solomon, E. I.; Schoonheydt, R. A.; Sels, B. F., Cu-ZSM-5: A Biomimetic Inorganic Model for Methane Oxidation. *J. Catal.* **2011**, *284*, 157.
- (12) Woertink, J. S.; Smeets, P. J.; Groothaert, M. H.; Vance, M. A.; Sels, B. F.; Schoonheydt, R. A.; Solomon, E. I., A $[\text{Cu}_2\text{O}]^{2+}$ Core in Cu-ZSM-5, the Active Site in the Oxidation of Methane to Methanol. *Proc. Natl. Acad. Sci. U.S.A.* **2009**, *106*, 18908.
- (13) Johnson, E. J.; Kleinlein, C.; Musgrave, R. A.; Betley, T. A., Diiron Oxo Reactivity in a Weak-Field Environment. *Chem. Sci.* **2019**, *10*, 6304.
- (14) Rosenthal, J.; Pistorio, B. J.; Chng, L. L.; Nocera, D. G., Aerobic Catalytic Photooxidation of Olefins by an Electron-Deficient Pacman Bisiron(III) μ -oxo Porphyrin. *J. Org. Chem.* **2005**, *70*, 1885.
- (15) Rosenthal, J.; Luckett, T. D.; Hodgkiss, J. M.; Nocera, D. G., Photocatalytic Oxidation of Hydrocarbons by a Bis-Iron(III)- μ -oxo Pacman Porphyrin Using O_2 and Visible Light. *J. Am. Chem. Soc.* **2006**, *128*, 6546.
- (16) Deng, Y.; Chang, C. J.; Nocera, D. G., Direct Observation of the “Pac-Man” Effect from Dibenzofuran-Bridged Cofacial Bisporphyrins. *J. Am. Chem. Soc.* **2000**, *122*, 410.
- (17) Wilding, M. J. T.; Iovan, D. A.; Betley, T. A., High-Spin Iron Imido Complexes Competent for C–H Bond Amination. *J. Am. Chem. Soc.* **2017**, *139*, 12043.
- (18) Kleinlein, C.; Zheng, S.-L.; Betley, T. A., Ground State and Excited State Tuning in Ferric Dipyrin Complexes Promoted by Ancillary Ligand Exchange. *Inorg. Chem.* **2017**, *56*, 5892.
- (19) King, E. R.; Sazama, G. T.; Betley, T. A., Co(III) Imidos Exhibiting Spin Crossover and C–H Bond Activation. *J. Am. Chem. Soc.* **2012**, *134*, 17858.
- (20) Rieth, R. D.; Mankad, N. P.; Calimano, E.; Sadighi, J. P., Palladium-Catalyzed Cross-Coupling of Pyrrole Anions with Aryl Chlorides, Bromides, and Iodides. *Org. Lett.* **2004**, *6*, 3981.
- (21) Chang, C. J.; Deng, Y.; Heyduk, A. F.; Chang, C. K.; Nocera, D. G., Xanthene-Bridged Cofacial Bisporphyrins. *Inorg. Chem.* **2000**, *39*, 959.
- (22) Meyer, E. M.; Gambarotta, S.; Floriani, C.; Chiesi-Villa, A.; Guastini, C., Polynuclear Aryl Derivatives of Group 11 Metals. Synthesis, Solid State–Solution Structural Relationship, and Reactivity with Phosphines. *Organometallics* **1989**, *8*, 1067.
- (23) Pauling, L. *The Nature of the Chemical Bond*; Cornell University Press Ithaca, NY, 1960; Vol. 260.
- (24) Laitar, D. S. Synthetic and catalytic studies of Group 11 N-heterocyclic carbene complexes. Ph.D. Dissertation, Massachusetts Institute of Technology, Cambridge, **2006**.
- (25) Walroth, R. C.; Lukens, J. T.; MacMillan, S. N.; Finkelstein, K. D.; Lancaster, K. M., Spectroscopic Evidence for a $3d^{10}$ Ground State Electronic Configuration and Ligand Field Inversion in $[\text{Cu}(\text{CF}_3)_4]^{1-}$. *J. Am. Chem. Soc.* **2016**, *138*, 1922.
- (26) Walroth, R. C.; Miles, K. C.; Lukens, J. T.; MacMillan, S. N.; Stahl, S. S.; Lancaster, K. M., Electronic Structural Analysis of Copper(II)–TEMPO/ABNO Complexes Provides Evidence for Copper(I)–oxoammonium Character. *J. Am. Chem. Soc.* **2017**, *139*, 13507.
- (27) Walroth, R. C.; Uebler, J. W. H.; Lancaster, K. M., Probing Cui in Homogeneous Catalysis Using High-Energy-Resolution Fluorescence-Detected X-Ray Absorption Spectroscopy. *Chem. Comm.* **2015**, *51*, 9864.
- (28) Tomson, N. C.; Williams, K. D.; Dai, X.; Sproules, S.; DeBeer, S.; Warren, T. H.; Wieghardt, K., Re-Evaluating the Cu K Pre-Edge Xas Transition in Complexes with Covalent Metal–Ligand Interactions. *Chem. Sci.* **2015**, *6*, 2474.
- (29) Kau, L. S.; Spira-Solomon, D. J.; Penner-Hahn, J. E.; Hodgson, K. O.; Solomon, E. I., X-Ray Absorption Edge Determination of the Oxidation State and Coordination Number of Copper. Application to the Type 3 Site in *Rhus vernicifera* Laccase and Its Reaction with Oxygen. *J. Am. Chem. Soc.* **1987**, *109*, 6433.
- (30) George, S. J.; Lowery, M. D.; Solomon, E. I.; Cramer, S. P., Copper L-Edge Spectral Studies: A Direct Experimental Probe of the Ground-State Covalency in the Blue Copper Site in Plastocyanin. *J. Am. Chem. Soc.* **1993**, *115*, 2968.
- (31) Wilding, M. J. T.; Iovan, D. A.; Wrobel, A. T.; Lukens, J. T.; MacMillan, S. N.; Lancaster, K. M.; Betley, T. A., Direct Comparison of C–H Bond Amination Efficacy through Manipulation of Nitrogen-Valence Centered Redox: Imido Versus Iminyl. *J. Am. Chem. Soc.* **2017**, *139*, 14757.
- (32) Dong, Y.; Lukens, J.; Clarke, R. M.; Zheng, S.-L.; Lancaster, K.; Betley, T., Synthesis, Characterization and C–H Amination Reactivity of Nickel Iminyl Complexes. *Chem. Sci.* **2020**, *Advance Article*, doi: 10.1039/C9SC04879K.
- (33) Wigley, D. E., Organoimido Complexes of the Transition Metals. In *Prog. Inorg. Chem.*; Karlin, K. D., Ed.; Wiley, **2007**, *42*, 239–482.
- (34) Nugent, W. A.; Haymore, B. L., Transition Metal Complexes Containing Organoimido (NR) and Related Ligands. *Coord. Chem. Rev.* **1980**, *31*, 123.
- (35) Wiese, S.; McAfee, J. L.; Pahls, D. R.; McMullin, C. L.; Cundari, T. R.; Warren, T. H., C–H Functionalization Reactivity of a Nickel–Imide. *J. Am. Chem. Soc.* **2012**, *134*, 10114.
- (36) Laskowski, C. A.; Miller, A. J. M.; Hillhouse, G. L.; Cundari, T. R., A Two-Coordinate Nickel Imido Complex That Effects C–H Amination. *J. Am. Chem. Soc.* **2011**, *133*, 771.
- (37) Berry, J. F., Terminal Nitrido and Imido Complexes of the Late Transition Metals. *Comment Inorg. Chem.* **2009**, *30*, 28.

- (38) Desnoyer, A. N.; Love, J. A., Recent Advances in Well-Defined, Late Transition Metal Complexes That Make and/or Break C–N, C–O and C–S Bonds. *Chem. Soc. Rev.* **2017**, *46*, 197.
- (39) Kogut, E.; Wiencko, H. L.; Zhang, L.; Cordeau, D. E.; Warren, T. H., A Terminal Ni(III)–Imide with Diverse Reactivity Pathways. *J. Am. Chem. Soc.* **2005**, *127*, 11248.
- (40) Baek, Y.; Betley, T. A., Catalytic C–H Amination Mediated by Dipyrin Cobalt Imidos. *J. Am. Chem. Soc.* **2019**, *141*, 7797.
- (41) Dai, X.; Warren, T. H., Discrete Bridging and Terminal Copper Carbenes in Copper-Catalyzed Cyclopropanation. *J. Am. Chem. Soc.* **2004**, *126*, 10085.
- (42) Decker, A.; Fenske, D.; Maczek, K., New Imido-Bridged Transition Metal Clusters: $[(C_5H_5)_4Ti_4(NSnMe_3)_4]$, $[Co_{11}(PPh_3)_3(NPh)_{12}]$, $[Ni_{11}Br_6(N^tBu)_8]$, and $[Li(thf)_4]_4[Cu_{24}(NPh)_{14}]$. *Angew. Chem. Int. Ed.* **1996**, *35*, 2863.
- (43) Reiß, P.; Fenske, D., Neue Amido- Und Imidoverbrückte Kupferkomplexe – Synthesen Und Strukturen Von $[Li(OEt_2)_2][Cu(NPh_2)_3]$, $[ClCuN(SnMe_3)_3]$, $[CuN(SnMe_3)_2]_4$, $[Cu_{16}(NH_2^tBu)_{12}Cl_{16}]$, $[CuNH^tBu]_8$, $[Li(dme)_3][Cu_6(NHMe)_3(NMe)_2]$, $[PPh_3(C_6H_4)CuNHMe]$, $[Li(dme)][Cu(NHMe)(NPh)]_2$ Und $[Li(dme)_3][Li(dme)_2][Cu_{12}(NPh)_8]$. *Z. Anorg. Allg. Chem.* **2000**, *626*, 1317.
- (44) Iovan, D. A.; Betley, T. A., Characterization of Iron-Imido Species Relevant for N-Group Transfer Chemistry. *J. Am. Chem. Soc.* **2016**, *138*, 1983.
- (45) King, E. R.; Hennessy, E. T.; Betley, T. A., Catalytic C–H Bond Amination from High-Spin Iron Imido Complexes. *J. Am. Chem. Soc.* **2011**, *133*, 4917.
- (46) DeBeer George, S.; Neese, F., Calibration of Scalar Relativistic Density Functional Theory for the Calculation of Sulfur K-Edge X-Ray Absorption Spectra. *Inorg. Chem.* **2010**, *49*, 1849.
- (47) DeBeer George, S.; Petrenko, T.; Neese, F., Time-Dependent Density Functional Calculations of Ligand K-Edge X-Ray Absorption Spectra. *Inorg. Chim. Acta* **2008**, *361*, 965.
- (48) Lukens, J. T.; DiMucci, I. M.; Kurogi, T.; Mindiola, D. J.; Lancaster, K. M., Scrutinizing Metal–Ligand Covalency and Redox Non-Innocence Via Nitrogen K-Edge X-Ray Absorption Spectroscopy. *Chem. Sci.* **2019**, *10*, 5044.
- (49) Neese, F., Definition of Corresponding Orbitals and the Diradical Character in Broken Symmetry Dft Calculations on Spin Coupled Systems. *J. Phys. Chem. Solids* **2004**, *65*, 781.
- (50) CAS(11,9) calculations were carried out on the monoanions, although these were found to have effectively single-configurational ground states, so results of hybrid DFT single point calculations were used to describe these complexes.
- (51) LeCloux, D. D.; Davydov, R.; Lippard, S. J., Synthesis and Characterization of Spin-Delocalized Carboxylate-Bridged Cu(I)–Cu(II) Mixed-Valence Complexes Having Only Oxygen Donor Ligands. *J. Am. Chem. Soc.* **1998**, *120*, 6810.
- (52) Barr, M. E.; Smith, P. H.; Antholine, W. E.; Spencer, B., Crystallographic, Spectroscopic and Theoretical Studies of an Electron-Deiocalized Cu(1.5)–Cu(1.5) Complex. *J. Chem. Soc. Chem. Comm.* **1993**, 1649.
- (53) Gupta, R.; Zhang, Z. H.; Powell, D.; Hendrich, M. P.; Borovik, A., Synthesis and Characterization of Completely Delocalized Mixed-Valent Dicopper Complexes. *Inorg. Chem.* **2002**, *41*, 5100.
- (54) Robin, M. B.; Day, P., Mixed Valence Chemistry - A Survey and Classification. In *Adv. Inorg. Chem. Rad.*; Emeléus, H. J.; Sharpe, A. G., Eds.; Elsevier, **1968**, *10*, 247–422.
- (55) Goodman, B.; Raynor, J., Electron Spin Resonance of Transition Metal Complexes. In *Adv. Inorg. Chem. Rad.*; Emeléus, H. J.; Sharpe, A. G., Eds.; Elsevier, **1970**, *13*, 135–362.
- (56) Demadis, K. D.; Hartshorn, C. M.; Meyer, T. J., The Localized-to-Delocalized Transition in Mixed-Valence Chemistry. *Chem. Rev.* **2001**, *101*, 2655.
- (57) Gagne, R. R.; Koval, C. A.; Smith, T. J.; Cimolino, M. C., Binuclear Complexes of Macrocyclic Ligands. Electrochemical and Spectral Properties of Homobinuclear $Cu^{II}Cu^{II}$, $Cu^{II}Cu^I$, and Cu^ICu^I Species Including an Estimated Intramolecular Electron Transfer Rate. *J. Am. Chem. Soc.* **1979**, *101*, 4571.
- (58) Long, R. C.; Hendrickson, D. N., Intramolecular Electron Transfer in a Series of Mixed-Valence Copper(II)-Copper(I) Complexes. *J. Am. Chem. Soc.* **1983**, *105*, 1513.
- (59) Ziegler, M. S.; Levine, D. S.; Lakshmi, K.; Tilley, T. D., Aryl Group Transfer from Tetraarylborato Anions to an Electrophilic Dicopper(I) Center and Mixed-Valence μ -Aryl Dicopper(I, II) Complexes. *J. Am. Chem. Soc.* **2016**, *138*, 6484.
- (60) Brunold, T. C.; Gamelin, D. R.; Solomon, E. I., Excited-State Exchange Coupling in Bent Mn(III)–O–Mn(III) Complexes: Dominance of the π/σ Superexchange Pathway and Its Possible Contributions to the Reactivities of Binuclear Metalloproteins. *J. Am. Chem. Soc.* **2000**, *122*, 8511.
- (61) Brunschwig, B. S.; Creutz, C.; Sutin, N., Optical Transitions of Symmetrical Mixed-Valence Systems in the Class II–III Transition Regime. *Chem. Soc. Rev.* **2002**, *31*, 168.
- (62) Solomon, E. I.; Heppner, D. E.; Johnston, E. M.; Ginsbach, J. W.; Cirera, J.; Qayyum, M.; Kieber-Emmons, M. T.; Kjaergaard, C. H.; Hadt, R. G.; Tian, L., Copper Active Sites in Biology. *Chem. Rev.* **2014**, *114*, 3659.
- (63) Harkins, S. B.; Mankad, N. P.; Miller, A. J. M.; Szilagyi, R. K.; Peters, J. C., Probing the Electronic Structures of $[Cu_2(\mu-XR_2)]^{n+}$ Diamond Cores as a Function of the Bridging X Atom (X = N or P) and Charge (n = 0, 1, 2). *J. Am. Chem. Soc.* **2008**, *130*, 3478.
- (64) Mankad, N. P.; Harkins, S. B.; Antholine, W. E.; Peters, J. C., Multifrequency EPR Studies of $[Cu^{1.5}Cu^{1.5}]^+$ for $Cu_2(\mu-NR_2)_2$ and $Cu_2(\mu-NR_2)_2$ Diamond Cores. *Inorg. Chem.* **2009**, *48*, 7026.
- (65) Harkins, S. B.; Peters, J. C., Amido-Bridged Cu_2N_2 Diamond Cores That Minimize Structural Reorganization and Facilitate Reversible Redox Behavior between a Cu^ICu^I and a Class III Delocalized $Cu^{1.5}Cu^{1.5}$ Species. *J. Am. Chem. Soc.* **2004**, *126*, 2885.
- (66) DeBeer George, S.; Metz, M.; Szilagyi, R. K.; Wang, H.; Cramer, S. P.; Lu, Y.; Tolman, W. B.; Hedman, B.; Hodgson, K. O.; Solomon, E. I., A Quantitative Description of the Ground-State Wave Function of Cu_A by X-Ray Absorption Spectroscopy: Comparison to Plastocyanin and Relevance to Electron Transfer. *J. Am. Chem. Soc.* **2001**, *123*, 5757.
- (67) Houser, R. P.; Young, V. G.; Tolman, W. B., A Thiolate-Bridged, Fully Delocalized Mixed-Valence Dicopper(I,II) Complex That Models the Cu_A Biological Electron-Transfer Site. *J. Am. Chem. Soc.* **1996**, *118*, 2101.
- (68) Takaoka, A.; Moret, M.-E.; Peters, J. C., A Ru(I) Metalloradical That Catalyzes Nitrene Coupling to Azoarenes from Arylazides. *J. Am. Chem. Soc.* **2012**, *134*, 6695.
- (69) Gloaguen, Y.; Rebreyend, C.; Lutz, M.; Kumar, P.; Huber, M.; van der Vlugt, J. I.; Schneider, S.; de Bruin, B., An Isolated Nitridyl Radical-Bridged $\{Rh(N)Rh\}$ Complex. *Angew. Chem. Int. Ed.* **2014**, *53*, 6814.
- (70) Tran, B. L.; Washington, M. P.; Henckel, D. A.; Gao, X.; Park, H.; Pink, M.; Mindiola, D. J., A Four Coordinate Parent Imidevia a Titanium Nitridyl. *Chem. Comm.* **2012**, *48*, 1529.
- (71) Lyaskovskyy, V.; Suarez, A. I. O.; Lu, H.; Jiang, H.; Zhang, X. P.; de Bruin, B., Mechanism of Cobalt(II) Porphyrin-Catalyzed C–H Amination with Organic Azides: Radical Nature and H-Atom Abstraction Ability of the Key Cobalt(III)–Nitrene Intermediates. *J. Am. Chem. Soc.* **2011**, *133*, 12264.
- (72) Mankad, N. P.; Antholine, W. E.; Szilagyi, R. K.; Peters, J. C., Three-Coordinate Copper(I) Amido and Aminyl Radical Complexes. *J. Am. Chem. Soc.* **2009**, *131*, 3878.
- (73) Büttner, T.; Geier, J.; Frison, G.; Harmer, J.; Calle, C.; Schweiger, A.; Schönberg, H.; Grützmacher, H., A Stable Aminyl Radical Metal Complex. *Science* **2005**, *307*, 235.

(74) Adhikari, D.; Mossin, S.; Basuli, F.; Huffman, J. C.; Szilagyi, R. K.; Meyer, K.; Mindiola, D. J., Structural, Spectroscopic, and Theoretical Elucidation of a Redox-Active Pincer-Type Ancillary Applied in Catalysis. *J. Am. Chem. Soc.* **2008**, *130*, 3676.

(75) Rodríguez-Lugo, R. E.; de Bruin, B.; Trincado, M.; Grützmacher, H., A Stable Aminyl Radical Coordinated to Cobalt. *Chem. Eur. J.* **2017**, *23*, 6795.

(76) Suarez, A. I. O.; Lyaskovskyy, V.; Reek, J. N. H.; van der Vlugt, J. I.; de Bruin, B., Complexes with Nitrogen-Centered

Radical Ligands: Classification, Spectroscopic Features, Reactivity, and Catalytic Applications. *Angew. Chem. Int. Ed.* **2013**, *52*, 12510.

Table of Contents Graphic

