



Direct formation of cyclobutenylphosphonates from 1-alkynylphosphonates and $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgCl}/2\text{CuCl}$

Yulia Sinelnikove^{a,b}, Abraham Rubinstein^b, Morris Srebnik^{a,*}, Abed Al Aziz Al Quntar^{a,c,*}

^a Department of Medicinal Chemistry and Natural Products, School of Pharmacy, Hebrew University, Jerusalem 91120, Israel

^b Department of Pharmaceutics, School of Pharmacy, Hebrew University, Jerusalem 91120, Israel

^c Department of Material Engineering, Faculty of Engineering, Al Quds University, Abu Deis, Palestinian Authority

ARTICLE INFO

Article history:

Received 6 November 2008

Revised 22 November 2008

Accepted 25 November 2008

Available online 30 November 2008

Keywords:

Alkynylphosphonates

Cyclization

Cyclobutenylphosphonates

Diethylzirconocene

Copper chloride

ABSTRACT

Zirconacycles **2** prepared from 1-alkynylphosphonates **1**, zirconocene dichloride, and 2 equiv of EtMgCl are smoothly converted into cyclobutenylphosphonates **3** when treated with two equiv of CuCl in 65–81% isolated yield. The reaction is specific and general only for zirconacyclopentenyl phosphonates.

© 2008 Elsevier Ltd. All rights reserved.

Cyclobutenes are a versatile group of compounds that are both pharmacologically active¹ and are also valuable synthetic intermediates that undergo many useful transformations² involving additions to the double bond (bromination,³ hydrogenation⁴ to the important cyclobutanes,⁵ epoxidations,^{6,7} haloselenylation,⁶ hydroboration,⁸ bromohydroxylation,⁶ haloselenylation⁹) polymerization,¹⁰ ring opening to dienes,¹¹ and ring contraction¹² to name a growing list of reactions. Accordingly, they have been prepared by various methods.¹³ Organometallic routes include stannylcupration/epoxides¹⁴ and ruthenium-catalysis/propargylic alcohols.¹⁵ Various zirconium-mediated syntheses¹⁶ of cyclobutenes consist of reaction of alkynyl halides with EtMgBr,¹⁷ cross-coupling of terminal alkynes and vinyl bromides,¹⁸ reaction of γ,γ -dialkoxyallylic zirconium species with acrylamide,¹⁹ migratory insertion of an isocyanide into 1-zirconacyclopent-3-ynes,²⁰ cyclodimerization of alkynes,²¹ by double carbonylation,²² coupling of heteroaryl-substituted alkynes,²³ and coupling of 1,4-dicuprio-1,3-dienes,²⁴ diiodination of zirconacyclopentenenes followed by treatment with *n*-BuLi,²⁵ and from symmetrical 1,3-diynes via zirconacyclopentenenes.²⁶ The zirconium-mediated methods are tedious at best. Vinyl phosphonates are important chemically²⁷ and have been

shown to possess pharmacological activity.²⁸ Therefore, in our opinion, juxtaposition of the cyclobutenyl and phosphonate groups should lead to interesting compounds. In fact, cyclobutenylphosphonates would represent an essentially unexplored class of compounds. Thus, Seyferth first reported the synthesis of only the parent compound in 1971 by copper-induced decomposition of α -diazo cyclopropylmethylphosphonate in 71% yield; but it could not be purified.²⁹ In a related procedure, α -diazo β -ketophosphonates were converted into cyclobutenone phosphonates.³⁰ The parent compound was also prepared by treating a polymer-supported cyclic phenylsulfonylmethylphosphonate with triethylamine.³¹ Darling and Subramanian prepared two fused cyclobutenes by elimination of the corresponding β -aminophosphonates.³²

In this Letter, we describe the preparation of cyclobutenylphosphonates from $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgCl}/2\text{CuCl}$. We had previously reported the synthesis of the zirconacyclopentenyl phosphonates **2** by the addition of alkynylphosphonates using the reagent Cp_2ZrEt_2 prepared according to Takahashi and Negishi.³³ Various novel phosphonates were prepared by addition of aldehydes, allyl bromide, and propargylbromide to the zirconacyclopentenyl phosphonates **2**.³⁴ However, when **2b**, obtained from 1-hexynylphosphonate and 2EtMgCl, was treated with two equiv of CuCl, refluxed for 36 h, and worked up, cyclobutenylphosphonate **3b** was obtained as the sole product in 78% yield (Eq. 1, Table 1).³⁵

* Corresponding authors. Tel.: +972 2 675 7301; fax: +972 2 675 8201 (A.A.A.Q.).

E-mail addresses: morris.srebnik@ekmd.huji.ac.il (M. Srebnik), abedalaziz@ekmd.huji.ac.il, abedalaziz@eng.alquds.edu (Abed Al Aziz Al Quntar).


Table 1
Diethyl 2-alkylcyclobut-1-enylphosphonate **3** from diethyl alkynylphosphonate

Entry	R	³¹ P (ppm)	Yield (isolated) ^a (%)
3a	<i>n</i> -C ₃ H ₇	10.70	75
3b	<i>n</i> -C ₄ H ₉	10.75	78
3c	<i>n</i> -C ₅ H ₁₁	10.67	73
3d	<i>n</i> -C ₈ H ₁₇	11.01	81
3e	<i>n</i> -C ₁₂ H ₂₅	12.78	76
3f	Phenyl	10.29	70
3g	PhCH ₂ OCH ₂ CH ₂	10.35	65
3h	(PhCH ₂ O)CHCH ₂ CH ₂ CH ₂ CH ₃	9.57	65


^a After silical gel chromatography.

Table 3


Attempted cyclization of disubstituted alkynes other than phosphonates

			
Entry	R ¹	R ²	Yield ^a
a	<i>n</i> -Bu	SiMe ₃	<2%
b	<i>n</i> -Bu	Sn(<i>n</i> -Bu) ₃	<2%
c	Ph	Ph	7%
d	Ethyl	Ethyl	5%

^a Estimated by GCMS.



This is very different from another reagent system we developed consisting of Cp₂ZrCl₂/2EtMgBr/2AlCl₃ which provided methylenecyclopropylphosphonates from 1-alkynylphosphonates (Eq. 2).³⁶ Thus, the products of cyclization obtainable from zirconacyclopentenylphosphonates are very sensitive to the metal in the transmetalation step.



After addition of 2 equiv of CuCl to zirconacyclopentenyl phosphonates **2** at room temperature, the reaction was refluxed for 36 h. It was quenched with dilute aqueous HCl. The reaction can be followed by ³¹P NMR and GC/MS. The cyclobut-1-enylphosphonate products **3** were extracted with ethyl acetate and isolated by silica gel column chromatography in good yield (65–81%). This efficient inter-molecular cyclization reaction is tolerant to alkyl (**3a–e**), phenyl (**3f**), and benzyloxy (**3g–h**), groups. The addition of two equivalents of anhydrous CuCl was essential in order to reach the maximum yield.

The reaction did not proceed either under CuCl catalytic conditions or at room temperature. Other copper-containing compounds were investigated, and the yields were varied. Thiophene-2-carbonyloxy copper(I) (Table 2, entry b) afforded the highest yields

after copper chloride and copper iodide (entry c) gave the lowest yields. On the other hand, when other metal chlorides such as TiCl₄, CoCl₂, SnCl₂, SbCl₃, ZnCl₂, CeCl₃, and BaCl₂ were used (Table 2, g–n), only the ethylated product was obtained, and no trace of cyclobutenylphosphonates **3** were detected. The reaction has limitations. Bulky substituted alkynylphosphonates, that is *t*-butylethynylphosphonate, did not afford the corresponding zirconacyclopentenylphosphonate, **2**, but rather the zirconacyclopentenylphosphonate which gave *cis*-*t*-butylvinylphosphonate after hydrolysis. Alkylhalide side chains are also not compatible. Thus, diethyl 5-chloro-1-pentynylphosphonate could not be converted to the corresponding cyclobutenylphosphonate. An intractable reaction mixture was obtained presumably due to various copper-induced coupling reactions.

We briefly investigated other disubstituted alkynes (Table 3). None gave satisfactory results. The reason for this is unclear but not surprising. In the past, we have noticed that the phosphono group enhances the stability of zirconacycles. For instance, zirconacyclopentenylphosphonates do not dimerize and do not require stabilization by adding phosphine ligands.³⁷ Be that as it may, the present method of preparing 2-substituted cyclobutenylphosphonates is general, conceptually simple, and is far superior to any methodology, whether organometallic or other, reported to date.

Acknowledgments

The study was supported by a research grant from the Hebrew University. Morris Srebnik and Abraham Rubinstein are affiliated with the David R. Bloom Center of Pharmacy.

References and notes

- (a) Chao, J.; Taveras, A. G.; Aki, C.; Dwyer, M.; Yu, Y.; Purakkatt, B.; Rindgen, D.; Jakway, J.; Hipkin, W.; Fosetta, J.; Fan, X.; Lundell, D.; Fine, J.; Minniccozi, M.; Phillips, J.; Merritt, J. R. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 3778–3783; (b) Gonsiorek, W.; Fan, X. D.; Hesk, D.; Fosetta, J.; Qiu, H. C.; Jakway, J.; Billah, M.; Dwyer, M.; Chao, J. H.; Deno, G.; Taveras, A.; Lundell, D. J.; Hipkin, R. W. *J. Pharmacol. Exp. Ther.* **2007**, *322*, 477–485.
- Gauvry, N.; Huet, F. *J. Org. Chem.* **2001**, *66*, 583–588.
- Brune, H. A.; Rottele, H.; Horlbeck, G. Z. *Naturforsch., B* **1972**, *B27*, 505–510.
- Tufariello, J. J.; Milowsky, A. S.; Alnuri, M.; Goldstein, S. *Tetrahedron Lett.* **1987**, *28*, 267–270.
- Ortuno, R. M.; Moglioni, A. G.; Moltrasio, G. Y. *Curr. Org. Chem.* **2005**, *9*, 237–259.
- Mevellec, L.; Evers, M.; Huet, F. *Tetrahedron* **1996**, *52*, 15103–15116.
- Freccero, M.; Gandolfi, R.; Sarzi-Amade, M. *Tetrahedron* **1999**, *55*, 11309–11330; Leyhane, A. J.; Snapper, M. L. *Org. Lett.* **2006**, *8*, 5183–5186.
- (a) Marsac, Y.; Nourry, A.; Legoup, S.; Pipelier, M.; Dubreuil, D.; Aubertin, A. M.; Bourgougnon, N.; Benhida, R.; Huet, F. *Tetrahedron* **2005**, *61*, 7607–7612; (b) Knapp, K. M.; Goldfuss, B.; Knochel, P. *Chem. Eur. J.* **2003**, *9*, 5259–5265;

Table 2
Effect of various metal complexes on cyclization of zirconacyclopentenyl phosphonates **2**

Entry	Metal complex	Yield of 3d (%)
a	CuCl	81
b	C ₄ H ₃ SC(O)OCu ^a	70
c	CuI	8
d	Cu(II)triflate	10
e	CuCN	15
f	CuIP(OCH ₃) ₃	45
g	TiCl ₄	0 ^b
h	CoCl ₂	0 ^b
i	SnCl ₂	0 ^b
j	SbCl ₃	0 ^b
k	ZnCl ₂	0 ^b
l	CeCl ₃	0 ^b
m	BaCl ₂	0 ^b
n	Mo(CO) ₆	0 ^b

^a Thiophene-2-carbonyloxy copper(I).

^b Only ethylated product obtained.

- Bassindale, M. J.; Hamley, P.; Harrity, J. P. A. *Tetrahedron Lett.* **2001**, 42, 9055–9057.
9. Lescop, C.; Mevellec, L.; Huet, F. *J. Org. Chem.* **2001**, 66, 4187–4193.
10. (a) Kitayama, T.; Kawauchi, T.; Nakamura, M.; Sufi, B.; Padias, A. B.; Hall, H. K. *Polymer* **2004**, 45, 5085–5093; (b) Chen, X. P.; Sufi, B. A.; Padias, A. B.; Hall, H. K. *Macromolecules* **2002**, 35, 4277–4281; (c) Perrott, M. G.; Novak, B. M. *Macromolecules* **1996**, 29, 1817–1823; (d) Wu, Z.; Grubbs, R. H. *Macromolecules* **1995**, 28, 3502–3508; (e) Casey, P. K.; Mathias, L. J. *Polym. Commun.* **1991**, 32, 27–29; (f) Hall, H. K.; Padias, A. B. *J. Polym. Sci., A1* **2003**, 41, 625–635.
11. (a) Ingham, S.; Turner, R. W.; Wallace, T. W. *Chem. Commun.* **1985**, 1664–1666; (b) Binns, F.; Hayes, R.; Ingham, S.; Saengchantara, S. T.; Turner, R. W.; Wallace, T. W. *Tetrahedron* **1992**, 48, 515–530; (c) Hayes, R.; Ingham, S.; Saengchantara, S. T.; Wallace, T. W. *Tetrahedron Lett.* **1991**, 32, 2953–2954; (d) Binns, F.; Hayes, R.; Hodgetts, K. J.; Saengchantara, S. T.; Wallace, T. W.; Wallis, C. J. *Tetrahedron* **1996**, 52, 3631–3658; (e) Hodgetts, K. J.; Saengchantara, S. T.; Wallis, C. J.; Wallace, T. W. *Tetrahedron Lett.* **1993**, 34, 6321–6324; (f) GourdelMartin, M. E.; Huet, F. *Tetrahedron Lett.* **1996**, 37, 7745–7748; (g) Youcef, R. A.; Boucheron, C.; Guillaume, S.; Legoupy, S.; Dubreuil, D.; Huet, F. *Synthesis* **2006**, 633–636.
12. Mevellec, L.; Huet, F. *Tetrahedron Lett.* **1995**, 36, 7441–7444.
13. (a) de Meijere, A. Ed. Carbocyclic Four-Membered Ring Compounds; Thieme: Stuttgart, **1997**; 17f, E17e–E17f.; (b) Jiang, X. F.; Ma, S. M. *Tetrahedron* **2007**, 63, 7589–7595; (c) Murakami, M.; Matsuda, T. In *Modern Allene Chemistry*; Krause, N., Hashmi, A. S. K., Eds.; Wiley-VCH: Weinheim, 2004; pp 727–815; (d) Ma, S. M. *Chem. Rev.* **2005**, 105, 2829–2871.
14. Barbero, A.; Cuadrado, P.; Garcia, C.; Rincon, J. A.; Pulido, F. J. *J. Org. Chem.* **1998**, 63, 7531–7533.
15. Le Paih, J.; Derien, S.; Bruneau, C.; Demerseman, B.; Toupet, L.; Dixneuf, P. H. *Angew. Chem., Int. Ed.* **2001**, 40, 2912–2915.
16. Hartley, R. C.; Caldwell, S. T. *J. Chem. Soc., Perkin Trans. 1* **2000**, 477–501.
17. Kasai, K.; Liu, Y. H.; Hara, R.; Takahashi, T. *Chem. Commun.* **1998**, 1989–1990.
18. Barluenga, J.; Rodriguez, F.; Alvarez-Rodrigo, L.; Fananas, F. J. *Chem. Eur. J.* **2004**, 10, 101–108.
19. Ito, H.; Sato, A.; Taguchi, T. *Tetrahedron Lett.* **1999**, 40, 3217–3220.
20. Bach, M. A.; Beweries, T.; Burlakov, V. V.; Arndt, P.; Baumann, W.; Spannenberg, A.; Rosenthal, U. *Organometallics* **2007**, 26, 4592–4597.
21. Liu, Y. H.; Liu, M. N.; Song, Z. Q. *J. Am. Chem. Soc.* **2005**, 127, 3662–3663.
22. Mito, S.; Takahashi, T. *Chem. Commun.* **2005**, 2495–2496.
23. Song, Z. Q.; Li, Y. X.; Liu, M.; Cong, L. Q.; Liu, Y. H. *Organometallics* **2006**, 25, 5035–5044.
24. Chen, C.; Xi, C.; Lai, C.; Wang, R.; Hong, X. *Eur. J. Org. Chem.* **2004**, 647–650.
25. Negishi, E.; Liu, F.; Choueiry, D.; Mohamud, M. M. *J. Org. Chem.* **1996**, 61, 8325–8328.
26. Liu, Y.; Sun, W.-H.; Nakajima, K.; Takahashi, T. *Chem. Commun.* **1998**, 1133–1134.
27. (a) Maffei, M. *Curr. Org. Synth.* **2004**, 1, 355–375; Minami, T.; Okauchi, T.; Kouno, R. *Synthesis* **2001**, 349–357; (b) Kouno, R.; Tsubota, T.; Okauchi, T.; Minami, T. *J. Org. Chem.* **2000**, 65, 4326–4332.
28. (a) Quntar, A. A.; Gallily, R.; Katzavian, G.; Srebnik, M. *Eur. J. Pharmacol.* **2007**, 556, 9–13; (b) Al-Quntar, A. A.; Baum, O.; Reich, R.; Srebnik, M. *Arch. Pharm.* **2004**, 337, 76–80.
29. Marmor, R. S.; Seyferth, D. J. *J. Org. Chem.* **1971**, 36, 128–136.
30. Andriamiadanarivo, R.; Chantegrel, B.; Deshayes, C.; Doutheau, A. *Tetrahedron Lett.* **1993**, 34, 7923–7924.
31. Liu, X.-L.; Sheng, S.-R.; Zhou, W.; Wang, Q.-Y.; Zhang, X.-L.; Gong, B. *Synth. Commun.* **2007**, 37, 119–127.
32. Darling, S. D.; Subramanian, N. *Tetrahedron Lett.* **1975**, 3279–3282.
33. Takahashi, T.; Seki, T.; Nitto, Y.; Saburi, M.; Rousset, C. J.; Negishi, E. *J. Am. Chem. Soc.* **1991**, 113, 6266–6268.
34. Ben-Valid, S.; Quntar, A. A.; Srebnik, M. *J. Org. Chem.* **2005**, 70, 3554–3559.
35. *Procedure for the synthesis of 3b*: To 0.306 g (1.05 mmol) of zirconocene dichloride dissolved in 6 ml of dry THF at -78°C was added 1.05 ml of 2 M EtMgBr (2.1 mmol) dropwise in a 25 ml round-bottomed flask. After stirring for 5 min at -78°C , 1 mmol of alkynylphosphonate was added and the reaction was gradually warmed to 25°C , and stirred for 2 h. Then, 0.20 g (2 mmol) of CuCl was added under N_2 . After stirring at reflux for 36 h, the reaction was quenched with dilute aqueous HCl. The oily product was extracted with Et_2O (2×10 ml), separated on silica gel column (70% petroleum ether: 30% ethyl acetate). ^1H NMR (300 MHz): δ 0.91 (t, 3H, $J_{\text{HH}} = 4.2$ Hz), 1.31 (t, 6H, $J_{\text{HH}} = 4.2$ Hz), 1.28–1.38 (overlap m, 2H), 1.43 (m, 2H), 2.35 (br t, 2H, $J_{\text{HH}} = 4.8$ Hz), 2.54 (br s, 4H), 4.04–4.10 (m, 4H). ^{31}P NMR (121.4 MHz): δ 10.75; ^{13}C NMR (75.5 MHz): δ 13.81, 16.39 (d, $^3J_{\text{PC}} = 3.7$ Hz), 22.52, 27.18 (d, $^2J_{\text{PC}} = 5.1$ Hz), 29.15, 30.41, 31.04 (d, $^3J_{\text{PC}} = 19.3$ Hz), 61.31 (d, $^2J_{\text{PC}} = 3.7$ Hz), 127.19 (d, $^1J_{\text{PC}} = 109.1$ Hz), 168.53 (d, $^2J_{\text{PC}} = 4.2$ Hz); MS(EI): m/z (%) 246 (6.5), 231 (8.8), 217 (48.0), 204 (83.4), 189 (28.2), 176 (24.7), 161 (82.2), 148 (62.0), 129 (8.9), 107 (56.1), 93 (50.8), 79 (100), 65 (62.7), 53 (20.1); Anal. Calcd for $\text{C}_{12}\text{H}_{23}\text{O}_3\text{P}$: C, 58.52; H, 9.41; P, 12.58. Found: C, 58.69; H, 9.47; P, 12.44.
36. Quntar, A. A.; Srebnik, M. *J. Org. Chem.* **2006**, 71, 730–733.
37. Quntar, A. A.; Srebnik, M. *Org. Lett.* **2001**, 3, 1379–1381.