This article was downloaded by: [University of Louisville] On: 07 October 2013, At: 14:19 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lncn19

Triarylmethyl Substituted 4,5-Dicyanoimidazoles as Activators for Rp-Diastereoselective Synthesis of TpN Dinucleoside Methylphosphonates

Peter Schell^a, Wolfgang H. G. Laux^a & Joachim W. Engels^a ^a Johann Wolfgang Goethe-Universität, Institut für Organische Chemie, Marie-Curie-Str. 11, D-60439, Frankfurt/Main, Germany Published online: 04 Oct 2006.

To cite this article: Peter Schell , Wolfgang H. G. Laux & Joachim W. Engels (1999) Triarylmethyl Substituted 4,5-Dicyanoimidazoles as Activators for Rp-Diastereoselective Synthesis of TpN Dinucleoside Methylphosphonates, Nucleosides and Nucleotides, 18:6-7, 1169-1174

To link to this article: http://dx.doi.org/10.1080/07328319908044654

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

TRIARYLMETHYL SUBSTITUTED 4,5-DICYANOIMIDAZOLES AS ACTIVATORS FOR Rp-DIASTEREOSELECTIVE SYNTHESIS OF TpN DINUCLEOSIDE METHYLPHOSPHONATES

Peter Schell, Wolfgang H. G. Laux and Joachim W. Engels*

Johann Wolfgang Goethe-Universität, Institut für Organische Chemie, Marie-Curie-Str. 11, D-60439 Frankfurt/Main, Germany

ABSTRACT: 2-Triarylmethyl-4,5-dicyanoimidazoles 1-3 were synthesized and tested as activators in the methylphosphonamidite approach. TpN dinucleoside methylphosphonates generated showed diastereoselectivity of up to 8 / 1 (Rp / Sp). The influence of the different triarylmethyl substituents on diastereoselectivity is shown.

In the antisense concept oligonucleoside methylphosphonates play an important role to control gene expression. They are taken up intact by cells in culture and animals and are stable against degradation by cellular nucleases^[1,2]. In a comparative study with several modified oligonucleotides chimeric methylphosphonate-phosphordiester oligodeoxy-nucleotides were reported to present most favorable characteristics as antisense agents^[3]. This is explained by the fact that chimeric antisense oligonucleotides improve the activation of RNase H combined with higher selectivity^[4]. Due to chirality at phosphorus oligonucleoside methylphosphonates containing n methylphosphonate linkages consist of a mixture of 2ⁿ diastereomers^[5]. Because of pseudo equatorial orientation of the methyl group in the duplex, Rp-configurated oligonucleoside methylphosphonates bind better to their target strand than the corresponding Sp-configurated oligonucleoside methylphosphonates are important tools to study protein-DNA or protein-RNA interactions^[7]. Most of the methods to synthesize diastereomerically pure methylphosphonates established so far use stereouniform precursors, which are coupled in a stereoselective or stereospecific

manner, using phosphorus (V) chemistry^[5,8]. Development of these methods to oligonucleotide solid phase synthesis are only beginning^[9].

We decided to use the phosphoramidite approach for diastereoselective synthesis of methylphosphonates because it is well established and effective for solid phase synthesis. Following the accepted mechanism^[10] the amidite is activated by tetrazole, which serves as acidic and nucleophilic activator forming an azolide intermediate. It reacts with the 5'-hydroxyl group of an oligonucleoside bound to the solid support yielding a phosphite triester. The azolide intermediate is responsible for epimerisation at phosphorus^[11], as is shown by experiments with diastereomerically enriched amidites. As a conclusion stereoselection during azole catalyzed reactions of this type was assumed to be impossible.

Exact examination of the mechanism revealed the possibility to use the dynamic equilibrium of the two diastereomeric azolide intermediates for dynamic kinetic resolution^[12]. Selectivity is observed when one of the diastereomeric products is preferentially formed from the azolide intermediate. In case of the methylphosphonates this should be achieved by attachment of a substituent to the azole moiety. Due to fast epimerisation the methylphosphonamidites can be used without separation as a diastereomeric mixture.



FIG. 1 Structures of the triarylmethyl substituted 4,5-dicyanoimidazoles 1-3

In an earlier report we utilized 2-biphenyl substituted 4,5-dicyanoimidazoles as activators for the diastereoselective synthesis of dinucleoside methylphosphonates. Thus, 2-(2',4',6'-trimethylbiphenyl-2-yl)-4,5-dicyanoimidazole gave reasonable results in favor of the Rp-Isomer^[13]. This stereoselection was rationalized in the following manner.

Shielding one side of the azole moiety by the biphenyl residue should lead to an influence on the coordination sphere of phosphorus. We explained the observed stereoselectivity by an approach of the incoming nucleophile, here the second nucleoside, from the less hindered side. In order to improve stereoselectivity we envisaged to further increase the bulkiness in the 2-position of the imidazole moiety. Thus, we synthesized triarylmethyl substituted 4,5-dicyanoimidazoles 1-3 (FIG. 1).

They are easily available by starting from the appropriate triarylmethyl chlorides. Outlined as an example we describe the synthesis of 2-(triphenylmethyl)-4,5-dicyanoimidazole 1 (FIG. 2).



FIG. 2 Synthesis of 2-(triphenylmethyl)-4,5-dicyanoimidazole 1

Tritylchloride 4 is converted into its corresponding nitrile 5 by trimethylsilylcyanide (TMSCN) in dichloromethane in the presence of $TiCl_4^{[14]}$. DIBAL-H reduction of $5^{[15]}$, followed by acid catalyzed condensation with diaminomaleonitrile (DAMN) yields azomethine $6^{[16]}$. Oxidative cyclisation by *N*-chlorosuccinimide and nicotinamide as base furnishes 2-(triphenylmethyl)-4,5-dicyanoimidazole 1 in an overall yield of 62 %^[17]. 2 and 3 were synthesized analogously to the above sequence in 27 % and 34 % yield, respectively (not optimized).

In order to test the diastereomeric induction of the newly synthesized 4,5-dicyanoimidazoles 1-3 the following model reaction for the amidite coupling was studied (FIG. 3).



a) i. 1, 2 or 3, CH_2Cl_2 , room temperature; ii. TBHP; 8: B = a: T, b: A^{Bzi} , c: C^{Bzi} , d: G^{ibu} , TBDPS = *tert*-butyldiphenylsilyl, DMTr = dimethoxytrityl

FIG. 3 Synthesis of TpN dinucleoside methylphosphonates to test diastereomeric induction of 1-3

TABLE 1 Results of the coupling reactions forming 9a-d with 1, 2 and 3 as activator

Dimer		³¹ P-NMR	³¹ P-NMR	1	2	3
		Rp-Isomer [δ]	Sp-Isomer [δ]	Rp / Sp	Rp / Sp	Rp / Sp
9 a	T-T	31.73	32.58	80 / 20	-	-
9b	T-A	32.00	32.48	87 / 13	89 / 11	83 / 17
9c	T-C	32.09	32.72	79 / 21	-	-
9d	T-G	31.62	33.79	76 / 24	-	-

Accordingly, 1, 2 or 3 (4 equiv.) were used in the reaction of thymidine methylphosphonamidite 7 (1.5 equiv.) with 3'-TBDPS protected nucleosides **8a-d** (1 equiv.) in dichloromethane yielding methylphosphonite intermediates. These were oxidized at the end of the reaction with *tert*-butylhydroperoxide (TBHP) in a stereoretentive manner to give dinucleoside methylphosphonates **9a-d**. The ratio of the two diastereomers of **9a-d** was determined by integrating the corresponding signals in the ³¹P-NMR spectrum. The results are summarized in table 1. The error of this determination was estimated to be ± 2 %.

The relative ratio of Rp and Sp diastereomers reflects the influence of the activators 1-3 used. Based on our rationale, increase in bulkiness, activator 2, favours Rp configuration whereas reduction in flexibility, activator 3, lowers the selectivity compared with activator 1. To confirm our theory (*vide supra*) we modeled the two diastereomeric azole intermediates formed by the reaction of 7 and 1 using molecular mechanics^[18]. As a result the second nucleoside can attack the azole intermediate which leads to the Rp-dinucleoside methylphosphonate easier, due to less steric shielding.

Though the increase in selectivity up to 8 / 1 (Rp / Sp) is quite acceptable the general utility of the activator 2 is paid for by reduced reactivity, resulting in prolonged reaction time.

Activators 1-3 are easily accessible even on a large scale. The reached diastereoselectivities are the highest known so far. The model system used allows application to solid phase synthesis with minor changes, studies are under way.

ACKNOWLEDGMENT: This work was supported by the Deutschen Forschungsgemeinschaft (Graduiertenkolleg "Chemische und Biologische Synthese von Wirkstoffen").

REFERENCES

- a) Miller, P. S. Bio/Technology, 1991, 9, 358-362; b) Miller, P. S. Progress in Nucleic Acid Research and Molecular Biology, 1996, 52, 261-291.
- [2] Uhlmann, E.; Peyman, A. Chem. Rev., 1990, 90, 543-584.
- [3] Giles, R. V.; Spiller, D. G.; Grzybowski, J.; Clark, R. E.; Nicklin, P. and Tidd, D. T. Nucleic Acids Res., 1998, 26, 1567-1575.
- [4] Agrawal, S.; Jiang, Z.; Zhao, Q.; Shaw, D.; Sun, D.; Saxinger, C. Nucleosides & Nucleotides, 1997, 16(7-9), 927-936.
- [5] On the problem of P-substituted oligonucleotides: Lebedev, A. V.; Wickstrom, E. *Perspectives in Drug Discovery and Design*, **1996**, *4*, 17-40, cit. Lit..
- [6] a) Lesnikowski, Z. J.; Jaworska, M.; Stec, W. J. Nucleic Acids Res., 1990, 18, 2109-2115; b) Vyazovkina, E. V.; Savchenko, E. V.; Lokhov, S. G.; Engels, J. W.; Wickstrom, E.; Lebedev, A. V. *ibid.*, 1994, 22, 2404-2409; c) Reynolds, M. A.; Hogrefe, R. I.; Jaeger, J. A.; Schwartz, D. A.; Riley, T. A.; Marvin, W. B.; Daily, W. J.; Vaghefi, M. M.; Knowles, S. K.; Klem, R. E.; Arnold Jr., L. J. *ibid.*, 1996, 24, 4584-4591.
- [7] a) Smith, S. A.; McLaughlin, L. W. Biochemistry, 1997, 36, 6046-6058; b) Strauss-Soukup, J. K.; Vaghefi, M. M.; Hogrefe, R. I.; Maher III, L. J. *ibid.*, 1997, 36, 8692-8698; c) Iwai, S.; Maeda, M.; Shimada, Y.; Hori, N.; Murata, T.; Morioka, H.; Ohtsuka, E. *ibid.*, 1994, 33, 5581-5588; d) Pritchard, C. E.; Grasby, J. A.; Hamy, F.; Zacharek, A. M.; Singh, M.; Karn, J.; Gait, M.J. Nucleic Acids Res., 1994, 22, 2592-2600; e) Noble, S. A.; Fisher, E. F.; Caruthers, M. H. *ibid.*, 1984, 12, 3387-3404.
- [8] a) Lesnikowski, Z. J. Bioorganic Chemistry, 1993, 21, 127-155, cit. Lit.; b) Jaworska-Maslanka, M. M.; Kacperczyk, W.; Korczynski, D.; Lesnikowski, Z. J.

Antisense & Nucleic Acid Drug Development, 1997, 7, 23-30; c) Daily, W. J.; Schwartz, D. A.; Riley, T. A.; Arnold Jr., L. J.; Marvin, WO. B.; Scurria, M. A.; Hopkins, S. A.; Atkins, M. B.; Garcia, C. D.; Pirrung, M. C. Nucleosides & Nucleotides, 1997, 16(4), 417-432; d) Stec, W. J.; Wozniak, L. A.; Pyzowski, J.; Niewiarowski, W. Antisense & Nucleic Acid Drug Development, 1997, 7, 381-395.

- [9] a) LeBec, C.; Wickstrom, E. Tetrahedron Lett., 1994, 35, 9525-9528; b) LeBec, C.;
 Wickstrom, E. J. Org. Chem., 1996, 61, 510-513.
- [10] a) Dahl, B. H.; Nielsen, J.; Dahl, O. Nucleic Acids Res., 1987, 15(4), 1729-1743; b)
 Berner, S.; Mühlegger, K.; Seliger, H. *ibid.*, 1989, 17(3), 853-864.
- [11] Stec, W.J.; Zon, G. Tetrahedron Lett., 1984, 25(46), 5279-5282.
- [12] Noyori, R.; Tokunaga, M.; Kitamura, M. Bull. Chem. Soc. Jpn., 1995, 68, 36-56.
- [13] Schell, P.; Engels, J. W. Tetrahedron Lett., submitted for publication.
- [14] Zieger, H. E.; Wo, S. J. Org. Chem., 1994, 59, 3838-3840.
- [15] LeBel, N. A.; Post, M. E.; Whang, J. J. J. Am. Chem. Soc., 1964, 86, 3759-3767.
- [16] Begland, R. W.; Hartter, D. R.; Jones, F. N.; Sam, D. J.; Sheppard, W. A.; Webster, O. W.; Weigert, F. J. J. Org. Chem., 1974, 39(16), 2341-2350.
- [17] Moriya, O.; Minamide, H.; Urata, Y. Synthesis, 1984, 1057-1058.
- [18] Egert, E.; Beck, H.; Bolte, M.; Gemmel, E. ,,MOMO 2.00 A Molecular Modeling Program", Frankfurt University, 1997.