

Copper-catalyzed direct thiolation of azoles with aliphatic thiols†

An-Xi Zhou, Xue-Yuan Liu,* Ke Yang, Shu-Chun Zhao and Yong-Min Liang

Received 14th March 2011, Accepted 21st April 2011

DOI: 10.1039/c1ob05395g

Cu(II)-catalyzed direct thiolation of azoles with thiols is described *via* intermolecular C–S bond formation/C–H functionalization under oxidative conditions. Both aryl thiols and aliphatic thiols are used as coupling partners, and furnished the thiolation products in moderate to good yields. The reaction is compatible with a wide range of heterocycles including oxazole, thiazole, imidazole and oxadiazole.

Introduction

Transition-metal-catalyzed C–S bond formation has been a subject of intense study due to the importance of aryl sulfides, aliphatic-heteroaromatic sulfides and their derivatives in numerous biological and pharmaceutically active compounds.¹ In the past decade, the transition-metal-catalyzed cross-coupling reactions of ArX (X = Cl, Br, I, OTf and B(OH)₂) with sulfur nucleophiles, such as, aryl thiols, aliphatic thiols and diaryl disulfides, are powerful tools for the formation of C–S bonds.²

In contrast, with the increasing requirements for environmentally benign and atom economic processes, the transition-metal-catalyzed C–H functionalization has received substantial attention in recent years, as a potentially more efficient and complementary process to the conventional cross-coupling methodology. However, in this aforementioned class of reactions, much effort has been paid on C–C and C–hetero bonds. The formation of a C–S bond *via* transition-metal-catalyzed C–H activation was not realized until Yu and co-workers first reported the thiolation of 2-phenylpyridine with PhSH and MeSSMe using Cu(OAc)₂ as a catalyst under oxygen atmosphere.³ Recently, Dong and co-workers described the Pd-catalyzed direct sulfonylation of a 2-phenylpyridine C–H bond with ArSO₂Cl.⁴ Very recently, Qing described a Cu(II)-mediated reaction of an aryl C–H bond with DMSO for *ortho*-substituted methylthiolation.⁵ Meanwhile, Cheng reported CuI-catalyzed thiolation of the di- or trimethoxyl- benzene arene C–H bond with ArSSAr.⁶ Besides the aforementioned aromatic C–S bond formation, the related counterparts on heterocycles were also revealed. Doi and Batey independently illustrated Pd-catalyzed reactions to produce 2-substituted benzothiazoles through C–H functionalization/intramolecular C–S bond formation.⁷ Subsequently, Li and co-workers disclosed iron-catalyzed sulfenylation of an indole C–H bond with diaryl disulfides, with a catalytic amount of iodine supplied to promote the reaction.⁸ Additionally, Fukuzawa and co-workers demonstrated a copper-catalyzed direct

thiolation of a benzoxazole C–H bond with diaryl disulfides and aryl thiols.⁹

The direct arylation,¹⁰ alkenylation¹¹ and amination¹² of heterocycles through the transition-metal-catalyzed C–H functionalization has been developed considerably. However, only one example of the formation of a C–S bond through the analogous reaction has been reported,⁹ and the application of aliphatic thiols in the reaction of direct C–H functionalization/intermolecular C–S bond formation is still unexplored. These drawbacks have limited this reaction which prepares sulfides with biological activity or sulfides that lead to biologically active compounds. Thus, further development for direct C–H thiolation is strongly desired. Herein, we report an efficient copper-catalyzed direct thiolation of azoles with aliphatic thiols.


Results and discussion

Our initial efforts focused on the direct thiolation of benzoxazole (**1a**) (0.4 mmol) with 1-dodecanethiol (**2a**) (0.6 mmol) by using a stoichiometric amount of Cu(OAc)₂·H₂O (2.2 equiv.) in DMSO under air atmosphere; the desired product was obtained in 27% isolated yield in 8 h (Table 1, entry 1). Encouraged by this preliminary result, we further investigated the reaction parameters and found that toluene was a better reaction medium than other solvents such as DMSO, DMF, xylene and dioxane (Table 1, entries 2–5). However, the desired cross-coupling was not observed when the amount of Cu(OAc)₂·H₂O was reduced to 20 mol% (Table 1, entry 6). We hypothesized that stoichiometric amounts of copper were needed because it was consumed as a base and an oxidant. With use of 20 mol% of Cu(OAc)₂·H₂O in the presence of 2.0 equiv. Cs₂CO₃ or K₂CO₃ (Table 1, entries 7–8), the reaction under O₂ resulted in trace yield of **3a**. Thus, further reactions with catalytic amounts of Cu(OAc)₂·H₂O were conducted in the presence of metal salts as a base and an oxidant including all kinds of copper and silver salts, and these experiments showed that reactions conducted with CuO as the additive formed the cross-coupling product **3a** in excellent yield (Table 1, entry 9). When O₂ was employed (1 atm) instead of air under the optimized conditions, the yield of **3a** dropped to 50% (Table 1, entry 10). A

State Key Laboratory of Applied Organic Chemistry, Lanzhou University, Lanzhou 730000, People's Republic of China. E-mail: liuxuey@lzu.edu.cn

† Electronic supplementary information (ESI) available: ¹H NMR and ¹³C NMR spectra for all new compounds. See DOI: 10.1039/c1ob05395g

Table 1 Optimization of the Reaction Conditions^a

				
Entry	Catalyst	Additive	Solvent	Yield ^b (%)
1	Cu(OAc) ₂ ·H ₂ O	Cu(OAc) ₂ ·H ₂ O	DMSO	27
2	Cu(OAc) ₂ ·H ₂ O	Cu(OAc) ₂ ·H ₂ O	DMF	38
3	Cu(OAc) ₂ ·H ₂ O	Cu(OAc) ₂ ·H ₂ O	xylene	35
4	Cu(OAc) ₂ ·H ₂ O	Cu(OAc) ₂ ·H ₂ O	dioxane	37
5	Cu(OAc) ₂ ·H ₂ O	Cu(OAc) ₂ ·H ₂ O	toluene	48
6	Cu(OAc) ₂ ·H ₂ O	—	toluene	0
7 ^c	Cu(OAc) ₂ ·H ₂ O	Cs ₂ CO ₃	toluene	trace
8 ^c	Cu(OAc) ₂ ·H ₂ O	K ₂ CO ₃	toluene	trace
9	Cu(OAc) ₂ ·H ₂ O	CuO	toluene	86
10 ^c	Cu(OAc) ₂ ·H ₂ O	CuO	toluene	50
11	Cu(OAc) ₂ ·H ₂ O	Cu(OH) ₂ CO ₃	toluene	73
12	Cu(OAc) ₂ ·H ₂ O	Cu(OH) ₂	toluene	38
13	Cu(OAc) ₂ ·H ₂ O	Ag ₂ CO ₃	toluene	53
14	Cu(OAc) ₂ ·H ₂ O	Ag ₂ O	toluene	36
15	Cu(OAc) ₂ ·H ₂ O	AgOAc	toluene	trace
16	Cu(OAc) ₂ ·H ₂ O	AgNO ₃	toluene	0
17	Cu(OAc) ₂ ·H ₂ O	AgI	toluene	0
18	CuO	CuO	toluene	17
19 ^d	Cu(OAc) ₂ ·H ₂ O	CuO	toluene	52

^a **1a** (0.4 mmol), dodecanethiol **2a** (0.6 mmol), Cu(II) (20 mol%), additive (2.0 equiv.), dry solvent (2 mL), under air, 120 °C, 8 h. ^b Isolated yields.

^c Under O₂ atmosphere. ^d Cu(OAc)₂·H₂O (10 mol%).


comparable reaction efficiency was presented by Cu(OH)₂CO₃, while Cu(OH)₂, Ag₂CO₃ and Ag₂O showed lower reactivity (Table 1, entries 11–14). Other silver species, such as AgOAc, AgNO₃ and AgI were ineffective for this reaction (Table 1, entries 15–17). Notably, with the use of CuO instead of Cu(OAc)₂·H₂O as the catalyst, the desired product was isolated in only 17% yield under the same reaction conditions (Table 1, entry 18). Along with reducing the amount of catalyst by 10 mol%, the yield decreased to 52% (Table 1, entry 19).

Under the optimized conditions, the substrate scope towards this thiolation reaction was further investigated. A wide array of heterocycle compounds **1** were tested in the reaction with 1-dodecanethiol (**2a**), and the results are listed in Table 2. Benzoxazoles bearing substituents with diverse electronic properties, such as electron donation (methyl, methoxy groups) and slight electron deficiency (chloro derivative), all showed the better reactivity and furnished the products in moderate to good yields (Table 2, entries 2–4). However, a low yield was obtained in the case of 5-nitrobenzoxazole (Table 2, entry 5), which was deactivated by the strongly electron withdrawing nitro group. Various azoles, such as oxazoles, thiazoles and imidazoles, all smoothly reacted with **2a** to afford the desired products in good yields (Table 2, entries 6–13). Moreover, 2-phenyl-1,3,4-oxadiazole is a good coupling partner, and the desired corresponding product **3n** was obtained in moderate yield (Table 2, entry 14).

The Cu-catalyzed thiolation of azoles with various thiols was also investigated. Primary aliphatic thiols, whether short, medium or long chain, were engaged in the system, the desired corresponding products were obtained in moderate to good yields (Table 3, entries 1, 2, 6–10). Moreover, secondary aliphatic thiols, such as 2-propanethiol (**2d**) and cyclohexylmercaptan (**2e**), were used in this reaction, which produced the corresponding product

3q and **3r** in 63% and 69% yield, respectively (Table 3, entries 3 and 4). However, when 2-methylpropane-2-thiol (**2f**) was reacted with benzoxazole (**1a**), the thiolation product **3s** was isolated only in 4% yield even with a prolonged reaction time of 24 h (Table 3, entry 5). These results indicated that the steric hindrance factors of the thiols play a key role in controlling the reactivity. We were also interested in extending the direct thiolation of azoles to aryl thiols. Benzoxazole, benzothiazole and 1-methylimidazole all smoothly reacted with benzenethiol (**2k**) to afford the desired products in low to moderate yields (Table 3, entries 11–13).


A mechanism analogous to those proposed for similar copper-catalyzed processes involving thiol copper species might be applicable to this thiolation procedure.^{6,9} To investigate the reaction mechanism, the thiolations of benzoxazole with dodecyl disulfide and bis(dodecylthio)copper ((C₁₂H₂₅S)₂Cu) were studied, respectively. When dodecyl disulfide was used as the reaction partner, the product **3a** was obtained in 46% yield under the optimized conditions for 24 h (Scheme 1, eq 1). On the other hand, the reaction of benzoxazole and (C₁₂H₂₅S)₂Cu under the same conditions for 8 h produced **3a** in 81% yield (Scheme 1, eq 2). Moreover, in the absence of CuO, the reaction yield was decreased to 57%, while in the absence of both CuO and Cu(OAc)₂·H₂O, the yield was further decreased to 25% (Scheme 1, eq 3). These results indicated the importance of the formation of (C₁₂H₂₅S)₂Cu as a key intermediate, which might be obtained more quickly from the reaction of the copper salt with a thiol than from the corresponding disulfide under these reaction conditions. Therefore, a plausible mechanism is proposed. The key intermediate (RS)₂Cu would be firstly formed in the presence of a thiol and the copper salt. It reacts with benzoxazole (**1a**) to produce the corresponding ArCuSPh intermediate, which can then undergo reductive elimination to afford the product **3a**. Finally, the Cu(II) catalyst is regenerated by the oxidant.

**Scheme 1** Preliminary mechanism study.

Conclusions

In conclusion, we have disclosed an efficient C–H functionalization/intermolecular C–S bond formation process. Various

Table 2 Reactions of **2a** with various heterocycles^a

					
Entry	Heterocycle	1	Product	3	Yield ^b (%)
1		1a (R = H)		3a (R = H)	86
2		1b (R = 5-Me)		3b (R = 5-Me)	84
3		1c (R = 5-OMe)		3c (R = 5-OMe)	63
4		1d (R = 6-Cl)		3d (R = 6-Cl)	67
5		1e (R = 5-NO ₂)		3e (R = 5-NO ₂)	11
6		1f		3f	63
7		1g (R = H)		3g (R = H)	82
8		1h (R = 6-Me)		3h (R = 6-Me)	84
9		1i (R ₁ , R ₂ = H)		3i (R ₁ , R ₂ = H)	83
10		1j (R ₁ , R ₂ = Me)		3j (R ₁ , R ₂ = Me)	89
11		1k		3k	70
12		1l		3l	69
13		1m		3m	73
14		1n		3n	50

^a **1** (0.4 mmol), dodecanethiol **2** (0.6 mmol), Cu(OAc)₂ · H₂O (20 mol %), CuO (2.0 equiv), toluene (2 mL), under air, 120 °C, 8h. ^b Isolated yields.

thioethers could be efficiently obtained from this method. Aliphatic thiols as reaction partners were used for the first time in the direct thiolation of azoles. This approach is simple, general, and practical which complemented the classic methods for the rapid construction of C–S bonds.

Experimental section

General remarks

Column chromatography was carried out on silica gel. Unless noted ¹H NMR spectra were recorded at 400 MHz in CDCl₃, ¹³C NMR spectra were recorded at 100 MHz in CDCl₃ using TMS as internal standard. IR spectra were recorded on an FT-IR spectrometer and only major peaks are reported in cm⁻¹. Melting points were determined on a microscopic apparatus and

were uncorrected. All new compounds were further characterized by HRMS (high resolution mass spectra); copies of their ¹H NMR and ¹³C NMR spectra are provided. Unless otherwise noted, materials obtained from commercial suppliers were used without further purification. Toluene was dried and distilled from sodium/benzophenone.

General procedure for the preparation of **3**

Under air atmosphere, a reaction vessel was charged with heterocycle **1** (0.4 mmol), RSH **2** (0.6 mmol), Cu(OAc)₂ · H₂O (16 mg, 20 mol%), CuO (64 mg, 0.8 mmol), and PhMe (2 mL). The mixture was stirred at 120 °C and monitored by TLC. After the completion of the reaction, the residue was purified directly by short flash column chromatography on silica gel with hexane/ethyl acetate as an eluent to give the desired corresponding product.

Table 3 Reactions of heterocycles with various thiols^a

Entry	1	Thiol 2	Product	Yield ^b (%)
1	1a	2b	3o	80
2	1a	2c	3p	78
3	1a	2d	3q	63
4 ^c	1a	2e	3r	69
5 ^d	1a	2f	3s	4
6	1j	2g $\text{CH}_3(\text{CH}_2)_3\text{SH}$	3t	82
7	1j	2h $\text{CH}_3(\text{CH}_2)_5\text{SH}$	3u	81
8	1j	2i $\text{CH}_3(\text{CH}_2)_7\text{SH}$	3v	84
9	1j	2j $\text{CH}_3(\text{CH}_2)_{17}\text{SH}$	3w	68
10	1m	2h	3x	77
11 ^c	1a	2k	3y	33
12 ^e	1g	2k	3z	41
13 ^e	1n	2k	3aa	55

^a **1** (0.4 mmol), thiol **2** (0.6 mmol), $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (20 mol%), CuO (2.0 equiv.), toluene (2 mL), under air, 120 °C, 8 h. ^b Isolated yields. ^c **2e** (0.72 mmol), $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (40 mol%) were used. ^d 24 h. ^e **2k** (0.72 mmol), 10 h.

2-(Dodecylthio)benzo[d]oxazole (3a). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); white solid, mp: 23.0–24.2 °C. ¹H NMR (400 MHz, CDCl_3) δ 0.88 (t, J = 6.8 Hz, 3H), 1.25–1.31 (m, 16H), 1.43–1.50 (m, 2H), 1.78–1.86 (m, 2H), 3.30 (t, J = 7.2 Hz, 2H), 7.20–7.28 (m, 2H), 7.42 (d, J = 7.6 Hz, 1H), 7.59 (d, J = 7.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl_3) δ 165.2, 151.7, 142.0, 124.1, 123.7, 118.3, 109.7, 32.2, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.0, 28.6, 22.7, 14.1; IR (thin film, cm^{-1}) 2921, 2849, 1503, 1468, 1130, 744; HRMS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{29}\text{NOS}$ [$M + \text{H}$]⁺: 320.2043, found: 320.2047.

2-(Dodecylthio)-5-methylbenzo[d]oxazole (3b). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); white solid, mp: 29.2–31.0 °C. ¹H NMR (400 MHz, CDCl_3) δ 0.88 (t, J = 6.8 Hz, 3H), 1.26–1.31 (m, 16H), 1.43–1.50 (m, 2H), 1.77–1.85 (m, 2H), 2.43 (s, 3H), 3.29 (t, J = 7.2 Hz, 2H), 7.02 (d, J = 8.0 Hz, 2H), 7.28 (d, J = 8.0 Hz, 2H), 7.38 (s, 1H); ¹³C NMR (100 MHz, CDCl_3) δ 165.1, 150.0, 142.2, 133.9, 124.6, 118.4, 109.1, 32.2, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.0, 28.6, 22.7, 21.4, 14.1; IR (thin film, cm^{-1}) 2919, 2850, 1637, 1497, 1152, 805; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{31}\text{NOS}$ [$M + \text{H}$]⁺: 334.2199, found: 334.2191.

2-(Dodecylthio)-5-methoxybenzo[d]oxazole (3c). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); white solid, mp: 24.8–26.0 °C. ¹H NMR (400 MHz, CDCl_3) δ 0.88 (t, J = 6.8 Hz, 3H), 1.26–1.31 (m, 16H), 1.43–1.50 (m, 2H), 1.78–1.85 (m, 2H), 3.29 (t, J = 7.2 Hz, 2H), 3.83 (s, 3H), 6.81 (dd, J = 8.8, 2.4 Hz, 1H), 7.11 (d, J = 2.4 Hz, 1H), 7.29 (d, J = 9.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl_3) δ 165.8, 157.1, 146.4, 142.8, 111.6, 109.8, 102.0, 55.9, 32.3, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.0, 28.6, 22.7, 14.1; IR (thin film, cm^{-1}) 2916, 1636, 1476, 1142, 826; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{31}\text{NO}_2\text{S}$ [$M + \text{H}$]⁺: 350.2148, found: 350.2143.

6-Chloro-2-(dodecylthio)benzo[d]oxazole (3d). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); white solid, mp: 41.2–42.8 °C. ¹H NMR (400 MHz, CDCl_3) δ 0.88 (t, J = 6.8 Hz, 3H), 1.26–1.30 (m, 16H), 1.45–1.50 (m, 2H), 1.78–1.85 (m, 2H), 3.29 (t, J = 7.2 Hz, 2H), 7.24 (dd, J = 8.4, 1.6 Hz, 1H), 7.42 (d, J = 1.6 Hz, 1H), 7.47 (d, J = 8.4 Hz, 1H); ¹³C NMR (100 MHz, CDCl_3) δ 166.1, 151.9, 140.8, 129.3, 124.7, 118.6, 110.4, 32.4, 31.9, 29.7, 29.6, 29.5, 29.4, 29.3, 29.2, 29.0, 28.6, 22.7, 14.1; IR (thin film, cm^{-1}) 2915, 2850, 1504, 1467, 1214, 1140, 817; HRMS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{28}\text{ClNOS}$ [$M + \text{H}$]⁺: 354.1653, found: 354.1660.

2-(Dodecylthio)-5-nitrobenzo[d]oxazole (3e). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); mp: 58.0–59.8 °C. ¹H NMR (400 MHz, CDCl_3) δ 0.88 (t, J = 6.8 Hz, 3H), 1.26–1.35 (m, 16H), 1.45–1.52 (m, 2H), 1.82–1.89 (m, 2H), 3.34 (t, J = 7.2 Hz, 2H), 7.52 (d, J = 9.2 Hz, 1H), 8.22 (dd, J = 8.8, 1.6 Hz, 1H), 8.46 (d, J = 2.0 Hz, 1H); ¹³C NMR (100 MHz, CDCl_3) δ 169.4, 155.3, 145.2, 142.5, 119.9, 114.4, 109.7, 32.6, 31.9, 29.6, 29.5, 29.4, 29.3, 29.1, 29.0, 28.6, 22.7, 14.1; IR (thin film, cm^{-1}) 2915, 2851, 2360, 1639, 1527, 1494, 1342, 1108, 819, 736; HRMS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{28}\text{ClNOS}$ [$M + \text{H}$]⁺: 365.1893, found: 365.1896.

2-(Dodecylthio)-5-phenyloxazole (3f). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); white solid, mp: 37.8–39.0 °C. ¹H NMR (400 MHz, CDCl_3) δ 0.88 (t, J = 6.8 Hz,

3H), 1.25–1.31 (m, 16H), 1.41–1.48 (m, 2H), 1.75–1.82 (m, 2H), 3.20 (t, $J = 7.2$ Hz, 2H), 7.26–7.31 (m, 2H), 7.37–7.40 (m, 2H), 7.56–7.59 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 160.1, 152.6, 128.8, 128.1, 127.8, 123.7, 123.0, 32.6, 31.9, 29.6, 29.5, 29.4, 29.3, 29.1, 28.6, 22.7, 14.1; IR (thin film, cm^{-1}) 2920, 2849, 1633, 1476, 1165, 1121, 759, 689; HRMS (ESI) m/z : calcd for $\text{C}_{21}\text{H}_{31}\text{NOS}$ $[\text{M} + \text{H}]^+$: 346.2199, found: 346.2190.

2-(Dodecylthio)benzo[d]thiazole (3g). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.26–1.30 (m, 16H), 1.43–1.50 (m, 2H), 1.77–1.84 (m, 2H), 3.33 (t, $J = 7.2$ Hz, 2H), 7.25–7.28 (m, 1H), 7.37–7.41 (m, 1H), 7.73 (d, $J = 8.0$ Hz, 1H), 7.86 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 167.3, 153.4, 135.1, 125.9, 124.0, 121.4, 120.8, 33.6, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.1, 28.7, 22.7, 14.1; IR (thin film, cm^{-1}) 2924, 2852, 2360, 1461, 1428, 1239, 996, 755; HRMS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{29}\text{NS}_2$ $[\text{M} + \text{H}]^+$: 336.1814, found: 336.1809.

2-(Dodecylthio)-6-methylbenzo[d]thiazole (3h). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.26–1.31 (m, 16H), 1.42–1.49 (m, 2H), 1.76–1.84 (m, 2H), 2.44 (s, 3H), 3.31 (t, $J = 7.2$ Hz, 2H), 7.20 (dd, $J = 8.4, 1.2$ Hz, 1H), 7.52 (s, 1H), 7.73 (d, $J = 8.4$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.0, 151.5, 135.3, 134.1, 127.4, 120.9, 120.7, 33.6, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.1, 28.7, 22.7, 21.4, 14.1; IR (thin film, cm^{-1}) 2920, 1641, 1446, 994, 813, 728; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{31}\text{NS}_2$ $[\text{M} + \text{H}]^+$: 350.1971, found: 350.1980.

2-(Dodecylthio)thiazole (3i). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.26–1.30 (m, 16H), 1.40–1.47 (m, 2H), 1.71–1.79 (m, 2H), 3.20 (t, $J = 7.2$ Hz, 2H), 7.19 (d, $J = 3.6$ Hz, 1H), 7.65 (d, $J = 3.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.4, 142.7, 118.5, 34.6, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.1, 28.7, 22.6, 14.1; IR (thin film, cm^{-1}) 2924, 2853, 2360, 1462, 1388, 1301, 1020, 706; HRMS (ESI) m/z : calcd for $\text{C}_{15}\text{H}_{27}\text{NS}_2$ $[\text{M} + \text{H}]^+$: 286.1658, found: 286.1656.

2-(Dodecylthio)-4,5-dimethylthiazole (3j). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.26–1.32 (m, 16H), 1.38–1.43 (m, 2H), 1.68–1.75 (m, 2H), 2.28 (s, 3H), 2.29 (s, 3H), 3.08 (t, $J = 7.2$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.4, 148.4, 126.4, 34.9, 31.9, 29.6, 29.5, 29.4, 29.3, 29.0, 28.6, 22.6, 14.6, 14.1, 11.2; IR (thin film, cm^{-1}) 2924, 2853, 1642, 1560, 1460, 1420, 1297, 1110, 1020, 723; HRMS (ESI) m/z : calcd for $\text{C}_{17}\text{H}_{31}\text{NS}_2$ $[\text{M} + \text{H}]^+$: 314.1971, found: 314.1975.

2-(Dodecylthio)-1-methyl-1H-benzo[d]imidazole (3k). Silica gel column purification with hexane/ethyl acetate (4/1, v/v); white solid, mp: 32.0–33.6 °C. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.25–1.31 (m, 16H), 1.43–1.50 (m, 2H), 1.75–1.82 (m, 2H), 3.38 (t, $J = 7.2$ Hz, 2H), 3.63 (s, 3H), 7.18–7.20 (m, 3H), 7.65–7.68 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 152.5, 143.5, 136.7, 121.6, 118.1, 108.2, 32.5, 31.8, 29.8, 29.6, 29.5, 29.4, 29.3, 29.2, 29.1, 28.7, 22.6, 14.0; IR (thin film, cm^{-1}) 2922, 2851, 1462, 1441, 1362, 1275, 1230, 911, 731; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{32}\text{N}_2\text{S}$ $[\text{M} + \text{H}]^+$: 333.2359, found: 333.2362.

1-Benzyl-2-(dodecylthio)-1H-benzo[d]imidazole (3l). Silica gel column purification with hexane/ethyl acetate (4/1, v/v); white solid, mp: 47.6–49.2 °C. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.25–1.30 (m, 16H), 1.40–1.45 (m, 2H), 1.73–1.80 (m, 2H), 3.39 (t, $J = 7.2$ Hz, 2H), 5.27 (s, 2H), 7.10–7.21 (m, 5H), 7.21–7.31 (m, 3H), 7.70 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 152.5, 143.7, 136.2, 135.7, 128.8, 127.8, 126.8, 121.8, 118.2, 109.0, 47.4, 32.8, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.1, 28.7, 22.6, 14.1; IR (thin film, cm^{-1}) 2924, 1642, 1440, 731; HRMS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{36}\text{N}_2\text{S}$ $[\text{M} + \text{H}]^+$: 409.2672, found: 409.2670.

2-(Dodecylthio)-1-methyl-1H-imidazole (3m). Silica gel column purification with hexane/ethyl acetate (4/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.25–1.30 (m, 16H), 1.36–1.41 (m, 2H), 1.61–1.68 (m, 2H), 3.05 (t, $J = 7.2$ Hz, 2H), 3.61 (s, 3H), 6.91 (s, 1H), 7.05 (s, 1H), 7.38 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 142.1, 129.1, 121.9, 34.3, 33.1, 31.8, 29.7, 29.6, 29.5, 29.4, 29.3, 29.1, 28.6, 22.6, 14.0; IR (thin film, cm^{-1}) 2924, 2853, 1461, 1279, 1124, 1079, 913, 723; HRMS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{30}\text{N}_2\text{S}$ $[\text{M} + \text{H}]^+$: 283.2202, found: 283.2198.

2-(Dodecylthio)-5-phenyl-1,3,4-oxadiazole (3n). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); mp: white solid, 32.4–34.0 °C. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.26–1.31 (m, 16H), 1.43–1.50 (m, 2H), 1.80–1.87 (m, 2H), 3.29 (t, $J = 7.2$ Hz, 2H), 7.46–7.51 (m, 3H), 7.99–8.01 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.5, 164.5, 131.5, 128.9, 126.6, 123.7, 32.6, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 29.0, 28.5, 22.6, 14.1; IR (thin film, cm^{-1}) 2921, 2846, 1637, 1468, 1382, 1186, 1064; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{30}\text{N}_2\text{OS}$ $[\text{M} + \text{H}]^+$: 347.2152, found: 347.2147.

2-(Ethylthio)benzo[d]oxazole (3o). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 1.50 (t, $J = 7.2$ Hz, 3H), 3.32 (q, $J = 14.8$ Hz, 2H), 7.20–7.29 (m, 2H), 7.42–7.44 (m, 1H), 7.59–7.61 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.0, 151.7, 142.0, 124.2, 123.7, 118.3, 109.8, 26.6, 14.7; IR (thin film, cm^{-1}) 2930, 2868, 1640, 1451, 1238, 1130, 1096, 924, 743; HRMS (ESI) m/z : calcd for $\text{C}_9\text{H}_9\text{NOS}$ $[\text{M} + \text{H}]^+$: 180.0478, found: 180.0482.

2-(Propylthio)benzo[d]oxazole (3p). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 1.08 (t, $J = 7.2$ Hz, 3H), 1.86 (m, 2H), 3.29 (t, $J = 7.2$ Hz, 2H), 7.20–7.29 (m, 2H), 7.41–7.43 (m, 1H), 7.59–7.61 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.2, 151.7, 141.9, 124.1, 123.7, 118.3, 109.7, 34.1, 22.7, 13.2; IR (thin film, cm^{-1}) 2965, 2873, 1640, 1500, 1452, 1238, 1214, 1130, 1095, 805, 742; HRMS (ESI) m/z : calcd for $\text{C}_{10}\text{H}_{11}\text{NOS}$ $[\text{M} + \text{H}]^+$: 194.0634, found: 194.0639.

2-(Isopropylthio)benzo[d]oxazole (3q). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 1.53 (d, $J = 6.8$ Hz, 6H), 4.04 (m, 1H), 7.21–7.29 (m, 2H), 7.43 (d, $J = 7.6$ Hz, 1H), 7.60–7.62 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 164.7, 151.5, 142.0, 124.1, 123.8, 118.4, 109.8, 38.3, 23.3; IR (thin film, cm^{-1}) 2968, 2360, 1648, 1500, 1453, 1237, 1128, 1094, 743; HRMS (ESI) m/z : calcd for $\text{C}_{10}\text{H}_{11}\text{NOS}$ $[\text{M} + \text{H}]^+$: 194.0634, found: 194.0638.

2-(Cyclohexylthio)benzo[d]oxazole (3r). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 1.35–1.39 (m, 1H), 1.44–1.66 (m, 5H), 1.78–1.83 (m, 2H), 2.19–2.23 (m, 2H), 3.86–3.93 (m, 1H), 7.20–7.29 (m, 2H), 7.41–7.43 (m, 1H), 7.60 (d, $J = 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 164.7, 151.6, 142.0, 124.1, 123.7, 118.3, 109.8, 46.0, 33.3, 25.7, 25.5; IR (thin film, cm^{-1}) 3056, 2931, 2854, 1498, 1451, 1238, 1128, 1094, 999, 807, 743; HRMS (ESI) m/z : calcd for $\text{C}_{13}\text{H}_{15}\text{NS}$ [$\text{M} + \text{H}$] $^+$: 234.0947, found: 234.0950.

2-(tert-Butylthio)benzo[d]oxazole (3s). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 1.65 (s, 9H), 7.26–7.30 (m, 2H), 7.46–7.48 (m, 1H), 7.64–7.67 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 163.3, 151.3, 142.0, 124.3, 124.2, 119.0, 109.9, 49.7, 30.9; IR (thin film, cm^{-1}) 2964, 1500, 1452, 1238, 1120, 1089, 806, 743; HRMS (ESI) m/z : calcd for $\text{C}_{11}\text{H}_{13}\text{NOS}$ [$\text{M} + \text{H}$] $^+$: 208.0791, found: 208.0789.

2-(Butylthio)-4,5-dimethylthiazole (3t). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.93 (t, $J = 7.2$ Hz, 3H), 1.41–1.50 (m, 2H), 1.67–1.74 (m, 2H), 2.28 (s, 3H), 2.30 (s, 3H), 3.09 (t, $J = 7.2$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.3, 148.4, 126.4, 34.6, 31.3, 21.8, 14.6, 13.5, 11.2; IR (thin film, cm^{-1}) 2958, 2867, 1560, 1419, 1375, 1297, 1111, 1019; HRMS (ESI) m/z : calcd for $\text{C}_9\text{H}_{15}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 202.0719, found: 202.0724.

2-(Hexylthio)-4,5-dimethylthiazole (3u). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.27–1.33 (m, 4H), 1.39–1.46 (m, 2H), 1.68–1.75 (m, 2H), 2.28 (s, 3H), 2.29 (s, 3H), 3.09 (t, $J = 7.2$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.3, 148.3, 126.4, 34.9, 31.2, 29.2, 28.3, 22.4, 14.6, 13.9, 11.2; IR (thin film, cm^{-1}) 2925, 2856, 1639, 1419, 1375, 1296, 1111, 1020, 726; HRMS (ESI) m/z : calcd for $\text{C}_{11}\text{H}_{19}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 230.1032, found: 230.1038.

4,5-Dimethyl-2-(octylthio)thiazole (3v). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.27–1.33 (m, 8H), 1.38–1.43 (m, 2H), 2.28 (s, 3H), 2.29 (s, 3H), 3.08 (t, $J = 7.2$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.3, 148.3, 126.4, 34.9, 31.7, 29.2, 29.0, 28.6, 22.6, 14.6, 14.0, 11.2; IR (thin film, cm^{-1}) 2924, 2854, 1560, 1461, 1420, 1374, 1296, 1111, 1019, 730; HRMS (ESI) m/z : calcd for $\text{C}_{13}\text{H}_{23}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 258.1345, found: 258.1343.

4,5-Dimethyl-2-(octadecylthio)thiazole (3w). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); white solid, mp: 36.0–38.0 °C. ^1H NMR (400 MHz, CDCl_3) δ 0.88 (t, $J = 6.8$ Hz, 3H), 1.25 (brs, 28H), 1.38–1.43 (m, 2H), 1.67–1.75 (m, 2H), 2.28 (s, 3H), 2.29 (s, 3H), 3.08 (t, $J = 7.2$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.4, 148.4, 126.4, 34.9, 31.9, 29.7, 29.6, 29.5, 29.4, 29.3, 29.2, 29.1, 28.7, 22.7, 14.6, 14.1, 11.3; IR (thin film, cm^{-1}) 2919, 2581, 1560, 1465, 1419, 1020, 909, 734; HRMS (ESI) m/z : calcd for $\text{C}_{23}\text{H}_{43}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 398.2910, found: 398.2901.

2-(Hexylthio)-1-methyl-1H-imidazole (3x). Silica gel column purification with hexane/ethyl acetate (4/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 0.87 (t, $J = 6.8$ Hz, 3H), 1.26–1.32 (m, 4H), 1.37–1.44 (m, 2H), 1.61–1.69 (m, 2H), 3.05 (t, $J = 7.2$ Hz,

2H), 3.61 (s, 3H), 6.91 (s, 1H), 7.05 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 142.0, 129.1, 121.9, 34.3, 33.1, 31.2, 29.6, 28.2, 22.4, 13.9; IR (thin film, cm^{-1}) 3106, 2927, 2856, 1459, 1414, 1377, 1279, 1124, 914, 728, 685; HRMS (ESI) m/z : calcd for $\text{C}_{23}\text{H}_{43}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 199.1263, found: 199.1269.

2-(Phenylthio)benzo[d]oxazole (3y). Silica gel column purification with hexane/ethyl acetate (4/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 7.21–7.29 (m, 2H), 7.39–7.41 (m, 1H), 7.43–7.47 (m, 3H), 7.59–7.61 (m, 1H), 7.69–7.71 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 163.2, 151.9, 142.0, 134.4, 129.8, 129.6, 127.2, 124.3, 119.1, 110.0; IR (thin film, cm^{-1}) 3060, 2925, 1799, 1500, 1450, 1237, 1127, 1093, 1024, 925, 804, 743; HRMS (ESI) m/z : calcd for $\text{C}_{23}\text{H}_{43}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 228.0478, found: 228.0481.

2-(Phenylthio)benzo[d]thiazole (3z). Silica gel column purification with hexane/ethyl acetate (80/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 7.28 (t, $J = 7.2$ Hz, 1H), 7.40–7.55 (m, 4H), 7.66 (d, $J = 8.0$ Hz, 1H), 7.76 (d, $J = 6.8$ Hz, 2H), 7.90 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.6, 153.9, 135.3, 130.4, 129.9, 126.1, 124.3, 121.9, 120.7; IR (thin film, cm^{-1}) 3059, 2930, 1580, 1458, 1425, 1309, 1237, 1080, 1004, 753, 690; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{31}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 244.0249, found: 244.0252.

1-Methyl-2-(phenylthio)-1H-imidazole (3aa). Silica gel column purification with hexane/ethyl acetate (4/1, v/v); colorless liquid. ^1H NMR (400 MHz, CDCl_3) δ 3.61 (s, 3H), 7.06 (d, $J = 1.2$ Hz, 1H), 7.11–7.17 (m, 4H), 7.22–7.29 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 137.8, 134.8, 130.0, 129.1, 127.8, 126.4, 123.7, 33.7; IR (thin film, cm^{-1}) 3056, 2946, 1582, 1456, 1411, 1280, 1122, 1081, 1024, 915, 743, 693; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{31}\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 191.0637, found: 191.0640.

Note added after first publication

This article replaces the version published on 7th June 2011, which contained errors in Table 3.

Acknowledgements

We thank the National Natural Science Foundation of China (No. 21072081), and the Fundamental Research Funds for the Central Universities (lzujbky-2009-81) for financial support.

Notes and references

- (a) S. Murru, B. K. Patal, J. L. Bras and J. Muzart, *J. Org. Chem.*, 2009, **74**, 2217; (b) M. D. Berger, M. Dutia, D. Powell, B. M. Floyd, N. Torres, R. Mallon, D. Wojciechowicz, S. Kim, L. Feldberg, K. Collins and I. Chaudhary, *Bioorg. Med. Chem.*, 2008, **16**, 9202; (c) M. A. Siracusa, L. Salerno, M. N. Modica, V. Pittala, G. Romeo, M. E. Amato, M. Nowak, A. J. Bojarski, I. Mereghetti, A. Cagnotto and T. Mennini, *J. Med. Chem.*, 2008, **51**, 4529; (d) Z. Kazimierzczuk, M. Andrzejewska, J. Kaustova and V. Klimešova, *Eur. J. Med. Chem.*, 2005, **40**, 203; (e) L. Llauger, H. He, J. Kim, J. Aguirre, N. Rosen, U. Peters, P. Davies and G. Chiosis, *J. Med. Chem.*, 2005, **48**, 2892; (f) L. Liu, J. E. Stelmach, S. R. Natarajan, M.-H. Chen, S. B. Singh, C. D. Schwartz, C. E. Fitzgerald, S. J. O'Keefe, D. M. Zaller, D. M. Schmatz and J. B. Doherty, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 3979; (g) J. Dumas, D. Brittelli, J. Chen, B. Dixon, H. Hatoum-Mokdad, G. König, R. Sibley, J. Witowsky and S. Wong, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 2531; (h) S. W. Kaldor, V. J. Kalish, J. F. Davies II, B. V. Shetty, J. E. Fritz, K. Appelt, J. A. Burgess, K. M. Campanale, N. Y. Chirgadze, D. K. Clawson, B. A.

- Dressman, S. D. Hatch, D. A. Khalil, M. B. Kosa, P. P. Lubbehusen, M. A. Muesing, A. K. Patick, S. H. Reich, K. S. Su and J. H. Tatlock, *J. Med. Chem.*, 1997, **40**, 3979; (i) X.-Z. Chen, P. Xu, L. Liu, D. Zheng and P.-S. Lei, *Eur. J. Med. Chem.*, 2011, **46**, 208; (j) G. David, T. Rustam T., S. Siegfried, P. Bruno and R. Patrick, *J. Sulfur Chem.*, 2010, **31**, 493; (k) R. Paramashivappa, P. P. Kumar, P. V. S. Rao and A. S. Rao, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 657.
- 2 (a) J. F. Hartwig, *Acc. Chem. Res.*, 2008, **41**, 1534; (b) T. Kondo and T. Mitsudo, *Chem. Rev.*, 2000, **100**, 3205; (c) R. Xu, J.-P. Wan, H. Mao and Y. Pan, *J. Am. Chem. Soc.*, 2010, **132**, 15531; (d) S. V. Ley and A. W. Thomas, *Angew. Chem., Int. Ed.*, 2003, **42**, 5400; (e) K. Kunz, U. Scholz and D. Ganzer, *Synlett*, 2003, 2428.
- 3 X. Chen, X.-S. Hao, C. E. Goodhue and J.-Q. Yu, *J. Am. Chem. Soc.*, 2006, **128**, 6790.
- 4 X. Zhao, E. Dimitrijevic and V. M. Dong, *J. Am. Chem. Soc.*, 2009, **131**, 3466.
- 5 L. Chu, X. Yue and F.-L. Qing, *Org. Lett.*, 2010, **12**, 1644.
- 6 S.-H. Zhang, P.-C. Qian, M.-L. Zhang, M.-L. Hu and J. Cheng, *J. Org. Chem.*, 2010, **75**, 6732.
- 7 (a) K. Inamoto, C. Hasegawa, K. Hiroya and T. Doi, *Org. Lett.*, 2008, **10**, 5147; (b) K. Inamoto, Y. Arai, K. Hiroya and T. Doi, *Chem. Commun.*, 2008, 5529; (c) L. L. Joyce and R. A. Batey, *Org. Lett.*, 2009, **11**, 2792.
- 8 X.-L. Fang, R.-Y. Tang, P. Zhong and J.-H. Li, *Synthesis*, 2009, 4183.
- 9 S.-I. Fukuzawa, E. Shimizu, Y. Atsuumi, M. Haga and K. Ogata, *Tetrahedron Lett.*, 2009, **50**, 2374.
- 10 Recent reviews on catalytic C–H bond arylation of heterocycles, see: (a) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; (b) L. Ackermann, R. Vincente and A. R. Kapdi, *Angew. Chem., Int. Ed.*, 2009, **48**, 9792; (c) B.-J. Li, S.-D. Yang and Z.-J. Shi, *Synlett*, 2008, 949; (d) J. C. Lewis, R. G. Bergman and J. A. Ellman, *Acc. Chem. Res.*, 2008, **41**, 1013 For selected recent breakthroughs, see: (e) H. Hachiya, K. Hirano, T. Satoh and M. Miura, *Angew. Chem., Int. Ed.*, 2010, **49**, 2202; (f) J. Huang, J. Chan, Y. Chen, C. J. Borths, K. D. Baucom, R. D. Larsen and M. M. Faul, *J. Am. Chem. Soc.*, 2010, **132**, 3674; (g) C.-L. Sun, B.-J. Li and Z.-J. Shi, *Chem. Commun.*, 2010, **46**, 677; (h) L. Ackermann, A. Althammer and S. Fenner, *Angew. Chem., Int. Ed.*, 2009, **48**, 201; (i) L.-C. Campeau, D. R. Stuart, J. P. Leclerc, M. Bertrand-Laperle, E. Villemure, H.-Y. Sun, S. Lasserre, N. Guimond, M. Lecavallier and K. Fagnou, *J. Am. Chem. Soc.*, 2009, **131**, 3291; (j) M. Tobisu, I. Hyodo and N. Chatani, *J. Am. Chem. Soc.*, 2009, **131**, 12070; (k) H.-Q. Do, R. K. M. Khan and O. Daugulis, *J. Am. Chem. Soc.*, 2008, **130**, 15185; (l) D. R. Stuart and K. Fagnou, *Science*, 2007, **316**, 1172.
- 11 Selected recent examples on C–H bond alkenylations of heteroaromatics, see: (a) A. García-Rubia, B. Urones, R. G. Arrayás and J. C. Carretero, *Chem.–Eur. J.*, 2010, **16**, 9676; (b) A. García-Rubia, R. G. Arrayás and J. C. Carretero, *Angew. Chem., Int. Ed.*, 2009, **48**, 6511.
- 12 Recent reviews on catalytic C–H bond amination of heteroaromatics, see: (a) A. Armstrong and J. C. Collins, *Angew. Chem., Int. Ed.*, 2010, **49**, 2282; (b) F. Collet, R. H. Dodd and P. Dauban, *Chem. Commun.*, 2009, 5061; (c) P. Thansandote and M. Lautens, *Chem.–Eur. J.*, 2009, **15**, 5874 For selected recent breakthroughs, see: (d) M. Miyasaka, K. Hirano, T. Satoh, R. Kowalczyk, C. Bolm and M. Miura, *Org. Lett.*, 2011, **13**, 359; (e) S. Guo, B. Qian, Y. Xie, C. Xia and H. Huang, *Org. Lett.*, 2011, **13**, 522; (f) T. Kawano, K. Hirano, T. Satoh and M. Miura, *J. Am. Chem. Soc.*, 2010, **132**, 6900; (g) J. Y. Kim, S. H. Cho, J. Joseph and S. Chang, *Angew. Chem., Int. Ed.*, 2010, **49**, 9899.