

# ORGANOMETALLICS

Volume 19, Number 18, September 4, 2000

© Copyright 2000  
American Chemical Society

## Communications

### Rhodium Carbene Complexes: Highly Selective Catalysts for the Hydroformylation of Styrene Derivatives

Austin C. Chen, Li Ren, Andreas Decken, and Cathleen M. Crudden\*

Department of Chemistry, University of New Brunswick, P.O. Box 45222,  
Fredericton, New Brunswick, Canada E3B 6E2

Received May 2, 2000

*Summary:* Two new carbene–rhodium complexes are reported that are active hydroformylation catalysts, giving very high selectivities for the branched isomer (> 95:5) when vinyl arenes are used as substrates. The carbene analogue of Wilkinson's catalyst,  $[RhIMes(PPh_3)_2Cl]$  (*IMes* = 1,3-bis(2,4,6-trimethylphenyl)-imidazol-2-ylidene), and the related carbonyl complex  $[RhIMes(PPh_3)(CO)Cl]$  were both prepared, and the latter was fully characterized using spectroscopic and crystallographic means.

The hydroformylation reaction is one of the most important catalytic reactions in industry that employs homogeneous metal complexes.<sup>1</sup> Its main use is the conversion of propylene into butanal, which is then used to prepare several other products, including paint solvents and plasticizers.<sup>2</sup> The hydroformylation reaction also has applications in the synthesis of pharmaceuticals and fine chemicals. With styrene derivatives as substrates, hydroformylation followed by oxidation of the aldehyde provides a useful method for the

synthesis of 2-arylpropanoic acids such as ibuprofen.<sup>3,4</sup> Directed hydroformylation reactions have also been reported that are extremely useful for the preparation of a variety of aldehydes with high levels of stereocontrol.<sup>5,6</sup>

In all of these examples, achieving high selectivities

(3) (a) Amer, I.; Alper, H. *J. Am. Chem. Soc.* **1990**, *112*, 3674. (b) Brown, J. M.; Cook, S. J.; Khan, R. *Tetrahedron* **1986**, *42*, 5105. (c) Rieu, J.-P.; Bouchelere, A.; Cousse, H.; Mouzin, G. *Tetrahedron* **1986**, *42*, 4095.

(4) For the asymmetric hydroformylation of styrene see: (a) Sakai, N.; Nozaki, K.; Takaya, H. *J. Chem. Soc., Chem. Commun.* **1994**, 395. (b) Sakai, N.; Mano, S.; Nozaki, K.; Takaya, H. *J. Am. Chem. Soc.* **1993**, *115*, 7033. (c) Horiuchi, T.; Ohta, T.; Shirakawa, E.; Nozaki, K.; Takaya, H. *J. Org. Chem.* **1997**, *62*, 4285. (d) Buisman, G. J. H.; Kamer, P. C. J.; van Leeuwen, P. W. N. M. *Tetrahedron: Asymmetry* **1993**, *4*, 1625. (e) Buisman, G. J. H.; Martin, M. E.; Vos, E. J.; Klootwijk, A.; Kamer, P. C. J.; van Leeuwen, P. W. N. M. *Tetrahedron: Asymmetry* **1995**, *6*, 719.

(5) (a) Breit, B. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 2835. (b) Breit, B. *Chem. Commun.* **1997**, 591. (c) Breit, B. *Liebigs Ann./Recl.* **1997**, 1841.

(6) (a) Burke, S. D.; Cobb, J. E. *Tetrahedron Lett.* **1986**, *27*, 4237. (b) Jackson, W. R.; Perlmutter, P.; Tasdelen, E. E. *J. Chem. Soc., Chem. Commun.* **1990**, 763.


(7) For detailed analyses of factors affecting the branched to linear selectivities in the hydroformylation of aliphatic olefins see: (a) van der Veen, L. A.; Boele, M. D. K.; Bregman, F. R.; Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Goubitz, K.; Fraanje, J.; Schenk, H.; Bo, C. *J. Am. Chem. Soc.* **1998**, *120*, 11616. (b) Kranenburg, M.; van der Burgt, Y. E. M.; Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Goubitz, K.; Fraanje, J. *Organometallics* **1995**, *14*, 3081. (c) Casey, C. P.; Whitaker, G. T. *Isr. J. Chem.* **1990**, *30*, 299. (d) Casey, C. P.; Paulsen, E. L.; Beuttenmueller, E. W.; Proft, B. R.; Petrovich, L. M.; Matter, B. A.; Powell, D. R. *J. Am. Chem. Soc.* **1997**, *119*, 11817. (e) Brown, J. M.; Kent, A. G. *J. Chem. Soc., Perkin Trans. 2* **1987**, 1597.

\* To whom correspondence should be addressed. E-mail: [crudden@unb.ca](mailto:crudden@unb.ca)

(1) Colquhoun, H. M.; Thompson, D. J.; Twigg, M. V. *Carbonylation: Direct Synthesis of Carbonyl Compounds*; Plenum Press: New York, 1991.

(2) Pruett, R. L. *Adv. Organomet. Chem.* **1979**, *17*, 1.

for either the branched or the linear isomer (depending on the application) is critical. Optimizing the branched/linear ratio is generally accomplished by altering the nature and the number of ligands on rhodium, which are usually phosphorus-based.<sup>7</sup> We report herein the first example of carbene rhodium complexes **1** and **2** as catalysts for the hydroformylation reaction.<sup>8</sup> These complexes are highly selective for the hydroformylation of styrene derivatives, giving up to 50:1 selectivity for the branched isomer (eq 1).



Rhodium complex **1** was prepared by treatment of IMes (1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene)<sup>9</sup> with freshly recrystallized Wilkinson's complex ([RhCl(PPh<sub>3</sub>)<sub>3</sub>]). The resulting yellow solid was characterized by <sup>1</sup>H and <sup>31</sup>P NMR. In the <sup>1</sup>H NMR spectrum, the two ortho methyl residues on the IMes ligand are inequivalent. Saturation transfer experiments revealed that these two methyl groups are undergoing slow exchange on the NMR time scale.<sup>10</sup> In the <sup>31</sup>P NMR spectrum, two resonances with distinct <sup>31</sup>P–<sup>103</sup>Rh coupling constants were observed at 48.7 and 35.4 ppm, attributable to triphenylphosphine ligands that are trans and cis to IMes. It is likely that the large bulk of the IMes ligand overrides the usual preference of [LRhCl(PPh<sub>3</sub>)<sub>2</sub>]-type derivatives of Wilkinson's complex to have the two phosphine ligands in a trans arrangement.<sup>11</sup> Complex **1** proved to be relatively sensitive to oxygen in solution, decomposing to give a green solution that contained uncomplexed PPh<sub>3</sub> along with triphenylphosphine oxide.

As phosphine lability is often beneficial to catalytic activity, we decided to assess the catalytic behavior of the IMes catalyst **1** in the hydroformylation reaction.


(8) For the reaction of Rh–IMes catalysts with H<sub>2</sub> and CO, see: Huang, J.; Stevens, E. D.; Nolan, S. *Organometallics* **2000**, *19*, 1194. A related rhodium complex with Ph<sub>2</sub>CH substituents on the imidazolium nitrogens has been reported: Kocher, C.; Herrmann, W. A. *J. Organomet. Chem.* **1997**, *532*, 261. For hydroformylation and other catalytic chemistry using saturated carbene ligands see: Lappert, M. F. In *Transition Metal Chemistry*; Muller, A., Diemann, E., Eds.; Verlag Chemie: Weinheim, Germany, 1981. For an excellent review of carbene chemistry with a description of catalytic applications see: Herrmann, W. A.; Kocher, C. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2162.

(9) Arduengo, A. J., III; Harlow, R. L.; Kline, M. *J. Am. Chem. Soc.* **1991**, *113*, 361.

(10) Doyle, M. J.; Lappert, M. F. *J. Chem. Soc., Chem. Commun.* **1974**, 679.

(11) McAuliffe, C. A.; Levason, W. *Phosphine, Arsine, and Stibine Complexes of the Transition Elements*; Elsevier: Amsterdam, 1979; p 137. It should be noted that Lappert has prepared complexes of this same type employing less sterically hindered carbene ligands: Doyle, M. J.; Lappert, M. F.; McLaughlin, G. M.; McMeeking, J. *J. Chem. Soc., Dalton Trans.* **1974**, 1494. Cardin, D. J.; Cetinkaya, B.; Doyle, M. J.; Lappert, M. F. *Chem. Soc. Rev.* **1973**, *2*, 99. Complexes of the type RhL<sub>3</sub>Cl (L = saturated carbene) have also been prepared: Cetinkaya, B.; Hitchcock, P. B.; Kucukbay, H.; Lappert, M. F.; Al-Juaid, S. *J. Organomet. Chem.* **1994**, *481*, 89. Cetinkaya, B.; Hitchcock, P. B.; Lappert, M. F.; Shaw, D. B.; Spyropoulos, K.; Warhurst, N. J. W. *J. Organomet. Chem.* **1993**, *459*, 311.


**Table 1. Hydroformylation of Aromatic Olefins with Carbene Catalyst **1**<sup>a</sup>**



entry	Ar	loading (%)	yield <sup>b</sup> (%)	B:L <sup>c</sup>
1	Ph	1.0	85	96:4
2	<i>p</i> -MePh	1.0	88	94:6
3	<i>p</i> -ClPh	0.8	94	98:2
4	<i>p</i> -MeOPh	1.0	91	95:5
5	<i>p</i> - <i>i</i> -BuPh	1.1	95	97:3

<sup>a</sup> Reaction conditions: substrate (0.8 mmol), catalyst (ca. 1%), and solvent (5 mL, dry, deoxygenated benzene) were treated with 500 psi of CO and 500 psi of H<sub>2</sub> at 60 °C for 20–22 h, except for entry 5 (16 h). <sup>b</sup> Isolated yield of chromatographically and spectroscopically homogeneous material. <sup>c</sup> B:L = branched:linear ratio.

**Table 2. Hydroformylation of Styrene with Various Catalysts<sup>a</sup>**



entry	cat.	additive	convers <sup>b</sup> (%)	TOF <sup>c</sup>	B:L <sup>d</sup>
1	<b>1</b>	none	78	7	95:5
2	<b>2</b>	none	38	4	94:6
3	<b>2</b>	1 PPh <sub>3</sub>	74	8	96:4
4	<b>2</b>	2 PPh <sub>3</sub>	79	8	96:4
5	<b>2</b>	10 PPh <sub>3</sub>	80	8	96:4
6	<b>3</b> <sup>e</sup>	2 PPh <sub>3</sub>	27	3	88:12


<sup>a</sup> Reaction conditions: see Table 1 footnote, 0.5% catalyst loading employed for 19 h, except entry 1, which was run for 20 h. <sup>b</sup> convers = conversion. <sup>c</sup> TOF = turnover frequency per hour. <sup>d</sup> B:L = branched:linear ratio. <sup>e</sup> Catalyst **3** = Rh(CO)Cl(PPh<sub>3</sub>)<sub>2</sub>.

As can be seen in Table 1, aromatic olefins are hydroformylated with extremely high branched to linear selectivities. The selectivities range from 94:6 for *p*-methylstyrene to 98:2 for *p*-chlorostyrene.<sup>12</sup> Although **1** displays good selectivity, its reactivity is marginal. As shown in Table 2, under the conditions studied, approximately 7 turnovers/h are obtained.

During hydroformylation, the originally bright yellow suspension of **1** in benzene was converted into a homogeneous, pale yellow solution that underwent no appreciable color change with time, provided that an inert argon atmosphere was maintained. This prompted us to prepare and isolate the corresponding carbonyl complex. Indeed, rhodium complex **2**, a pale lime yellow powder, could be readily prepared by passing a stream of CO through a THF solution of **1**. The resulting complex (**2**) is highly stable, and no special precautions were required in its manipulation. In the <sup>1</sup>H NMR spectrum of complex **2**, the two ortho methyl substituents on IMes are equivalent. The <sup>31</sup>P NMR spectrum of **2** contained, as expected, only one phosphine doublet at 31.7 ppm, with a <sup>31</sup>P–<sup>103</sup>Rh coupling constant of 115 Hz, consistent with a phosphine ligand trans to the IMes.<sup>13</sup> X-ray-quality crystals were grown by layering

(12) Hydroformylation of aliphatic olefins was also performed with limited success. For example, with 1-octene as substrate, a linear to branched ratio of 2.7:1 was obtained. Conversion was quantitative after 16 h; the yield of the aldehyde was determined to be 97% (NMR yield, with hexamethylbenzene as internal standard) and was not accompanied by isomerization.

(13) Doyle, M. J.; Lappert, M. F.; Pye, P. L.; Terreros, P. *J. Chem. Soc., Dalton Trans.* **1984**, 2355.



**Figure 1.** Thermal ellipsoids and numbering scheme for **2**. Ellipsoids are drawn at the 30% probability level, and hydrogen atoms are omitted for clarity. Selected bond distances (Å) and angles (deg): Rh–C(2), 2.032(4); Rh–C(1), 1.704(15); Rh–P, 2.3016(12); Rh–Cl, 2.396(6); C(1)–Rh–C(2), 91.5(7); C(1)–Rh–P, 92.6(7); C(2)–Rh–P, 174.46(12); C(1A)–Rh–ClA, 176.8(7); C(2)–Rh–Cl, 89.3(2); P–Rh–Cl, 86.9(2).<sup>14</sup>

a dichloromethane solution of **2** with pentane at  $-25^{\circ}\text{C}$  (Figure 1).

The X-ray crystal structure reveals a distorted-square-planar Rh center (sum of bond angles around Rh  $360.19^{\circ}$ ) and a trans relationship between the IMes and  $\text{PPh}_3$  ligands. The C(2)–Rh–P tensor is  $174.46(12)^{\circ}$  and deviates slightly from linearity. The Cl and CO positions are disordered and were refined with site occupancies of 0.55 and 0.45, respectively. The Rh–C(2) distance of 2.032(4) Å and the Rh–P distance of 2.3016(12) Å is in accord with expected values.<sup>9,15</sup> The bulky mesityl substituents adopt a propeller-like arrangement, forming dihedral angles of  $81.3$  and  $75.5^{\circ}$  with respect to the central imidazole ring.

In the hydroformylation reaction, complex **2** is approximately half as active as **1**, but the selectivity is essentially retained (compare entries 1 and 2 in Table

**Table 3.** Hydroformylation of Vinyl Arenes with Carbene Catalyst **2**<sup>a</sup>

$$\text{Ar-CH=CH}_2 \xrightarrow[\text{H}_2/\text{CO}]{\text{2 (1\%)} + \text{PPh}_3 \text{ (2\%)}} \text{Ar-CH(CHO)-CH}_3 \quad (4)$$

entry	Ar	yield <sup>b</sup> (%)	B:L <sup>c</sup>
1 <sup>d</sup>	Ph		96:4
2	<i>p</i> -MePh	96	96:4
3	<i>p</i> -ClPh	94	97:3
4	<i>p</i> -MeOPh	98	95:5
5 <sup>e</sup>	<i>p</i> - <i>i</i> -BuPh	93	96:4

<sup>a</sup> Reaction conditions: see Table 1 footnote, reaction time 22 h. <sup>b</sup> Isolated yield of chromatographically and spectroscopically homogeneous material. <sup>c</sup> B:L = branched:linear. <sup>d</sup> 0.5% catalyst used. <sup>e</sup> 2% catalyst used.

2). Gratifyingly, activity could be recovered by the addition of triphenylphosphine. In comparison with the corresponding Rh–phosphine complex Rh(CO)Cl( $\text{PPh}_3$ )<sub>2</sub> (**3**), carbene catalyst **2** is approximately twice as active and gives higher selectivity for the branched isomer (compare entries 4 and 6 in Table 2). Under optimized conditions, a variety of vinyl arenes could be hydroformylated to give the corresponding aldehydes as shown in Table 3.

In conclusion, we have successfully prepared two new rhodium carbene catalysts capable of hydroformylating vinylarenes with high selectivity for the branched isomer. The preparation of an asymmetric version is currently underway.

**Acknowledgment.** The Natural Sciences and Engineering Research Council of Canada (NSERC) is gratefully acknowledged for support of this research in terms of equipment and operating grants to C.M.C. and a postgraduate scholarship to A.C.C. The Canada Foundation for Innovation is also thanked for funding this research. UNB and BioChem Pharma are thanked for providing scholarships to L.R. and A.C.C. We thank Professors Steven P. Nolan, Larry A. Calhoun, and David C. Forbes for useful discussions and Dr. Robert McDonald at the University of Alberta for X-ray data collection.

**Supporting Information Available:** Tables giving data and details of the crystal structure determination for **2** and text giving spectral data for **1** and **2** and experimental procedures for the synthesis of **1** and **2**, as well as for the hydroformylation reaction. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM0003739

(14) Crystal data for **2**:  $\text{C}_{40}\text{H}_{39}\text{ClN}_2\text{OPRh}$ , crystal size  $0.04 \times 0.20 \times 0.27$  mm, monoclinic,  $P2_1/c$ ,  $a = 17.7743(14)$  Å,  $b = 13.7175(11)$  Å,  $c = 15.0069(12)$  Å,  $\beta = 92.818(2)^{\circ}$ ,  $V = 3654.5(5)$  Å<sup>3</sup>,  $Z = 4$ ,  $2\theta(\text{max}) = 52.8^{\circ}$ ,  $\omega$  scans, 17 895 reflections scanned, 7485 unique reflections ( $R(\text{int}) = 0.0954$ ),  $T = 293(2)$  K, full-matrix least squares on  $F^2$ ,  $wR2 = 0.0712$ ,  $R1 = 0.0436$ . Full details for the crystallographic analysis of **2** are described in the Supporting Information.

(15) (a) Herrmann, W. A.; Gooden, L. J.; Spiegler, M. J. *Organomet. Chem.* **1997**, *547*, 357. (b) Coleman, A. W.; Hitchcock, P. B.; Lappert, M. F.; Maskell, R. K.; Muller, J. G. *J. Organomet. Chem.* **1985**, *296*, 173.