


Ni-Catalyzed Carboxylation of Unactivated Primary Alkyl Bromides and Sulfonates with CO₂Yu Liu,[†] Josep Cornella,[†] and Ruben Martin^{*,†,‡}[†]Institute of Chemical Research of Catalonia (ICIQ), Av. Països Catalans 16, 43007 Tarragona, Spain[‡]Catalan Institution for Research and Advanced Studies (ICREA), Passeig Lluís Companys 23, 08010 Barcelona, Spain

Supporting Information


ABSTRACT: A Ni-catalyzed carboxylation of *unactivated* primary alkyl bromides and sulfonates with CO₂ at atmospheric pressure is described. The method is characterized by its mild conditions and remarkably wide scope without the need for air- or moisture-sensitive reagents, which make it a user-friendly and operationally simple protocol en route to carboxylic acids.

The use of CO₂ as an alternative renewable feedstock has recently received significant attention in the scientific community.¹ Such interest is primarily associated with the fact that CO₂ is nontoxic, abundant, and nonflammable, hence constituting an opportunity for carbon sequestration and allowing the implementation of innovative, yet practical, methodologies that may be counterintuitive at first sight.¹ Beyond any doubt, the synthesis of carboxylic acids represents an ideal target in CO₂ fixation since a myriad of molecules, such as Atorvastatin, Beraprost, Artesunate, Pemetrezed, and Pregabalin, among others, display significant biological activities (Scheme 1).^{2,3}

Scheme 1. Biologically Significant Carboxylic Acids




Encouraged by the seminal work of Osakada and Yamamoto,⁴ we⁵ and others⁶ launched a program aimed at unlocking the potential of CO₂ in reductive catalytic reactions (Scheme 2, path a).⁷ Unlike carboxylation events based on stoichiometric, well-defined, and in some cases air-sensitive organometallic species (path b),^{8,9} such reductive events offer higher flexibility and ease of execution by using simpler building blocks, thus representing an added value from a simplicity, reliability, and step-economical standpoint. Unfortunately, reductive carboxylation protocols are inherently restricted to substrates that rapidly undergo oxidative addition, such as aryl^{5,6} or benzyl halides (path a).^{5a,b} Ideally, this field should include the use of *unactivated* alkyl electrophiles possessing β -hydrogens. Indeed, these substrates are the most

Scheme 2. Reductive Carboxylation Reactions with CO₂

challenging in the cross-coupling arena because of their reluctance to undergo oxidative addition and the proclivity of in situ-generated alkylmetal species for β -hydride elimination, homodimerization, or hydrogen abstraction pathways, among others.¹⁰ Therefore, at the outset of our investigations it was unclear whether a metal-catalyzed carboxylation event could ever be conducted with unactivated alkyl electrophiles.¹¹ If successful, such a process would offer an unrecognized opportunity in CO₂ fixation while opening up new possibilities via unconventional bond disconnections. Herein we report a mild Ni-catalyzed carboxylation of unactivated primary alkyl bromides and sulfonates possessing β -hydrogens with CO₂ (path c). The protocol represents a convenient method that provides rapid access to carboxylic acids from simple precursors without the need to handle air-, moisture-sensitive reagents or cyanide sources and is characterized by a wide scope and an excellent chemoselectivity profile.

We initiated our investigations with **1a** as the model substrate with CO₂ (1 atm) at room temperature (Table 1). As expected, the conditions previously employed for the carboxylation of aryl halides^{5c,6} or primary benzylic halides^{5a,b} failed to convert **1a** into **2a**. Initial screening of metal complexes identified NiCl₂·glyme as a competent catalyst with cheap Mn as a reducing agent.¹² While nitrogen donors have successfully been employed as ligands in cross-coupling reactions of unactivated alkyl halides,¹³ no conversion to **2a** was observed with commonly employed bipyridines, terpyridines, or oxazolines (**L1–L9**).¹² In these cases, dimerization, β -hydride elimination, and recovered


Received: January 26, 2014

Table 1. Ligand Influence on the Reaction Outcome^{a,b}

^aReaction conditions: **1a** (0.15 mmol), $\text{NiCl}_2 \cdot \text{glyme}$ (10 mol %), **L** (22 mol %), and Mn (0.33 mmol) in DMA (0.15 M) at rt under CO_2 (1 atm) for 12 h. ^bYields were determined by HPLC analysis using naphthalene as an internal standard. ^cIsolated yield. ^d**1a** (1.0 mmol). ^e $\text{NiCl}_2 \cdot \text{glyme}$ (5 mol %).

starting material were observed in the crude reaction mixtures. A similar reactivity pattern was found when simple phenanthroline-type ligands (**L10**–**L13**) were employed. We speculated that an increase in the steric bulk around the nitrogen-donor ligand could lead to more robust Ni complexes with enhanced stability and greater activity. In line with this notion, we found that **L14** delivered **2a** in 66% yield. Analogously, **L16**, a bench-stable ligand readily obtained in one-step and in bulk quantities,¹⁴ afforded **2a** in 76% isolated yield. Dimerization and traces of β -hydride elimination byproducts accounted for the observed mass balance.¹⁵ Importantly, the reaction could be scaled up without any erosion in yield. Intriguingly, subtle changes in the electronic or steric environment of the 1,10-phenanthroline backbone had a deleterious effect (**L15** and **L17**).¹⁶ Control experiments unambiguously revealed that all of the reaction components were necessary to promote the carboxylation of **1a**.^{12,17}


Encouraged by these findings, we set out to explore the preparative scope of our reaction. As shown in Table 2, a host of unactivated primary alkyl bromides possessing β -hydrogens could be equally accommodated in good yields.^{18,19} Particularly illustrative is the chemoselectivity profile of our protocol, as esters (**2f**, **2i**–**l**, **2r**, **2t**, **2w**, and **2y**), nitriles (**2g**), heterocycles (**2k** and **2l**), acetals (**2e**), amides (**2o**), ketones (**2j** and **2n**), and even aldehydes (**2q**) were tolerated. Notably, unprotected aliphatic alcohols (**2h**), phenols (**2s**), and carbonyl compounds containing relatively acidic α -protons (**2g**, **2i**, **2j**, **2n**, **2o**, and **2w**) did not compete with the efficacy of the carboxylation event. At the current level of development, unactivated secondary alkyl bromides cannot be employed as coupling partners.¹⁹ Surprisingly, the reaction could also be conducted in the presence of aryltin reagents (**2p**) with **L14**, thus providing ample opportunities for subsequent manipulation. Interestingly, no macrocycle resulting from an intramolecular addition of the aryltin into the $\text{C}(\text{sp}^3)\text{--Br}$ bond or a carboxylation event on the C--Sn bond were detected in the crude material.¹⁹ While conformational restrictions might account for the former, the latter is particularly interesting since organotin reagents have been reported to efficiently undergo carboxylation events.²⁰

Table 2. Ni-Catalyzed Carboxylation of Alkyl Bromides^{a,b}

^aReaction conditions: **1a–u** (0.30 mmol), $\text{NiCl}_2 \cdot \text{glyme}$ (10 mol %), **L16** (22 mol %), and Mn (0.66 mmol) in DMA (0.15 M) at rt under CO_2 (1 atm) for 12 h. ^bIsolated yields (averages of at least two independent runs) are shown. ^cUsing **L14** (22 mol %).

Similarly, **L14** provided better results for **2s**. Site selectivity could be accomplished in the presence of electrophilic sites amenable to Ni-catalyzed cross-coupling reactions such as aryl pivalates (**2t**),²¹ acetates (**2y**),²¹ carbamates (**2v**),²¹ and aryl fluorides (**2r**).²² While aryl chlorides,^{6a} tosylates,^{6a} or pivalates^{5a} have been used in reductive carboxylation reactions, we found exclusive CO_2 insertion into the $\text{C}(\text{sp}^3)\text{--Br}$ bond (**2t** and **2u–x**). The synthetic value of this transformation is illustrated by a concise synthesis of compounds that exhibit potent biological activities such as MCPB (**2x**) and α -CEHC (**2y**) from available precursors.¹²

In light of these results, we wondered whether we could extend our Ni-catalyzed reductive carboxylation event to unactivated alkyl sulfonates. While the reaction of **3b** under the optimized conditions for alkyl bromides (Table 2) resulted in lower conversions to products, the combination of $\text{NiBr}_2 \cdot \text{glyme}$, **L14**, and DMF as the solvent at 50 °C under 1 atm CO_2 was optimal, furnishing the corresponding carboxylic acid in 74% yield (Table 3).¹² Interestingly, alkyl mesylates (**3c**) and trifluoroacetates (**3d**) could also be employed, albeit in lower yields. Notably, the


Table 3. Ni-Catalyzed Carboxylation of Alkyl Sulfonates^{a,b}

^aReaction conditions: **3a–f** (0.25 mmol), NiBr₂·glyme (10 mol %), L14 (26 mol %), and Mn (2.4 equiv) in DMF (0.25 M) at 50 °C for 12 h. ^bIsolated yields (averages of at least two independent runs) are shown. ^cAt 60 °C. ^dAt 100 °C. ^eNiBr₂·glyme (7.5 mol %). ^fAt 70 °C.

presence of other C–O electrophiles such as alkyl pivalates did not interfere, resulting in the selective carboxylation of the alkyl sulfonate backbone (**3f**). Overall, we believe that the results in Tables 2 and 3 show the robustness and the prospective impact of our Ni-catalyzed carboxylative protocol when employing unactivated alkyl bromides or alkyl sulfonates.²³

Although an in-depth mechanistic study should await further investigations, we wondered whether the reaction was initiated by β -hydride elimination followed by a hydrocarboxylation event.²⁴ Thus, we subjected 5-phenylpentene (**5**) to our optimized conditions. Under the limits of detection, we did not detect any carboxylation reaction.¹² A similar result was obtained when *n*-butylmanganese bromide (**6**) was exposed to our Ni/L16 system in the presence or absence of Mn, thus leaving some doubt about the intermediacy of organomanganese species.¹² In order to shed light on the mechanism, we decided to study the carboxylation reaction of **7a** and **7b** (Scheme 3).¹²

Scheme 3. Mechanistic Experiments



Diastereomerically pure **8** was anticipated for a mechanism consisting of a “classical” oxidative addition;²⁵ on the contrary, a statistical mixture of diastereoisomers in **8** would indicate a free-radical mechanism via single-electron transfer (SET). As shown in Scheme 3, ¹H NMR spectroscopic analysis of the crude mixture revealed the loss of stereochemical integrity at C1.²⁶ Similar behavior was found for **7c** and **7d**, an observation that might indicate a scenario consisting of SET processes via Ni(I) species.^{27–31} In line with this notion, we observed that radical clocks such as (bromomethyl)cyclopropane and 1-bromo-5-hexene resulted in ring-opened products.

In summary, we have reported a new catalytic carboxylation of unactivated primary alkyl bromides and sulfonates possessing β -hydrogens with CO₂ that gives access to valuable carboxylic acids. This method is characterized by its exquisite functional group compatibility, mild conditions, ready availability of the

starting materials, and ease of execution without the need for air- or moisture-sensitive materials. Further investigations into the mechanism and the extension to more challenging substrate combinations are currently underway.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

rmartinromo@iciq.es

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This paper is dedicated to the memory of Prof. Gregory L. Hillhouse. We thank ICIQ, the European Research Council (ERC-277883), and MINECO (CTQ2012-34054 and Severo Ochoa Excellence Accreditation 2014-2018 SEV-2013-0319) for support. Johnson Matthey, Umicore, and Nippon Chemical Industrial are acknowledged for gifts of metal and ligand sources. Y.L. and J.C. thank COFUND and the European Union for a Marie Curie Fellowship (FP7-PEOPLE-2012-IEF-328381).

■ REFERENCES

- (1) For selected reviews, see: (a) Zhang, L.; Hou, Z. *Chem. Sci.* **2013**, *4*, 3395. (b) Tsuji, Y.; Fujihara, T. *Chem. Commun.* **2012**, *48*, 9956. (c) Cokoja, M.; Bruckmeier, C.; Rieger, B.; Herrmann, W. A.; Kühn, F. E. *Angew. Chem., Int. Ed.* **2011**, *50*, 8510. (d) Martin, R.; Kleij, A. W. *ChemSusChem* **2011**, *4*, 1259. (e) Huang, K.; Sun, C.-L.; Shi, Z.-J. *Chem. Soc. Rev.* **2011**, *40*, 2435. (f) Sakakura, T.; Choi, J.-C.; Yasuda, H. *Chem. Rev.* **2007**, *107*, 2365.
- (2) (a) Patai, S. *The Chemistry of Acid Derivatives*; Wiley: New York, 1992. (b) Goossen, L. J.; Rodríguez, N.; Goossen, K. *Angew. Chem., Int. Ed.* **2008**, *47*, 3100.
- (3) Maag, H. *Prodrugs of Carboxylic Acids*; Springer: New York, 2007.
- (4) Osakada, K.; Sato, R.; Yamamoto, T. *Organometallics* **1994**, *13*, 4645.
- (5) (a) Correa, A.; León, T.; Martin, R. *J. Am. Chem. Soc.* **2014**, *136*, 1062. (b) León, T.; Correa, A.; Martin, R. *J. Am. Chem. Soc.* **2013**, *135*, 1221. (c) Correa, A.; Martin, R. *J. Am. Chem. Soc.* **2009**, *131*, 15974.
- (6) (a) Fujihara, T.; Nogi, K.; Xu, T.; Terao, J.; Tsuji, Y. *J. Am. Chem. Soc.* **2012**, *134*, 9106. (b) Tran-Vu, H.; Daugulis, O. *ACS Catal.* **2013**, *3*, 2417.
- (7) For electrochemical carboxylative processes of organic halides, see, for example: (a) Ohkoshi, M.; Michinishi, J.-Y.; Hara, S.; Senboku, H. *Tetrahedron* **2010**, *66*, 7732. (b) Amatore, C.; Jutand, A.; Khalil, F.; Nielsen, M. F. *J. Am. Chem. Soc.* **1992**, *114*, 7076.
- (8) For a review, see: Correa, A.; Martin, R. *Angew. Chem., Int. Ed.* **2009**, *48*, 6201.
- (9) For selected carboxylations of alkyl organoboranes, see: (a) Ohishi, T.; Zhang, L.; Nishiura, M.; Hou, Z. *Angew. Chem., Int. Ed.* **2011**, *50*, 8114. (b) Ohmiya, H.; Tanabe, M.; Sawamura, M. *Org. Lett.* **2011**, *13*, 1086. For carboxylation of alkyl organozincs, see: (c) Ochiai, H.; Jang, M.; Hirano, K.; Yorimitsu, H.; Oshima, K. *Org. Lett.* **2008**, *10*, 2681.
- (10) For selected reviews of the use of unactivated alkyl halides, see: (a) Kambe, N.; Iwasaki, T.; Terao, J. *Chem. Soc. Rev.* **2011**, *40*, 4937. (b) Hu, X. *Chem. Sci.* **2011**, *2*, 1867. (c) Jana, R.; Pathak, T. P.; Sigman, M. S. *Chem. Rev.* **2011**, *111*, 1417.
- (11) For photoinduced events with excess Sm/SmI₂ using tungsten lamps, see: Nomoto, A.; Kojo, Y.; Shiino, G.; Tomisaka, Y.; Mitani, I.; Tatsumi, M.; Ogawa, A. *Tetrahedron Lett.* **2010**, *51*, 6580.
- (12) See the Supporting Information for details.

(13) Selected references: (a) Everson, D. A.; Jones, B. A.; Weix, D. J. *J. Am. Chem. Soc.* **2012**, *134*, 6146. (b) Choi, J.; Fu, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 9102. (c) Binder, J. T.; Cordier, C. J.; Fu, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 17003. (d) Phapale, V. B.; Guisán-Ceinos, M.; Buñuel, E.; Cárdenas, D. J. *Chem.—Eur. J.* **2009**, *15*, 12681. (e) Phapale, V. B.; Buñuel, E.; García-Iglesias, M.; Cárdenas, D. J. *Angew. Chem., Int. Ed.* **2007**, *46*, 8790. (f) Powell, D. A.; Maki, T.; Fu, G. C. *J. Am. Chem. Soc.* **2005**, *127*, 510. (g) Powell, D. A.; Fu, G. C. *J. Am. Chem. Soc.* **2004**, *126*, 7788.

(14) Pijper, P. J.; Van der Goot, H.; Timmerman, H.; Nauta, W. T. *Eur. J. Med. Chem.* **1984**, *19*, 399.

(15) Lowering the loading of the reaction components for the carboxylation of **1a** was not successful. A lower Ni:L ratio resulted in higher amounts of dimerization. At present we believe that an L:Ni ratio ≥ 2 is needed to stabilize the resting state of the active Ni(0) catalyst while avoiding undesired pathways. A similar Ni:L effect has been observed in other reductive events. See: Wu, F.; Lu, W.; Qian, Q.; Ren, Q.; Gong, H. *Org. Lett.* **2012**, *14*, 3044. Also see refs 5 and 6.

(16) At present, we believe that electron-rich ligands such as **L15** might prevent the binding of CO₂ to the Ni center.

(17) While the use of unactivated alkyl chlorides resulted in no conversion, the coupling of alkyl iodides delivered yields of 14–19% with significant amounts of dimerization events. All attempts to improve these results were unsuccessful.

(18) The coupling of phenethyl electrophiles primarily resulted in dimerization with traces of styrene derivatives.

(19) Dimerization and β -hydride elimination account for the mass balance. Although a screening was conducted for substrates with low yields, the results were not satisfactory.

(20) Selected examples: (a) Wu, J.; Hazari, N. *Chem. Commun.* **2011**, 47, 1069. (b) Shi, M.; Nicholas, K. M. *J. Am. Chem. Soc.* **1997**, *119*, 5057 and citations therein.

(21) For selected reviews dealing with C–O electrophiles, see: (a) Mesganaw, T.; Garg, N. K. *Org. Process Res. Dev.* **2013**, *17*, 29. (b) Yamaguchi, J.; Muto, K.; Itami, K. *Eur. J. Org. Chem.* **2013**, 19. (c) Rosen, B. M.; Quasdorf, K. W.; Wilson, D. A.; Zhang, N.; Resmerita, A.-M.; Garg, N. K.; Percec, V. *Chem. Rev.* **2011**, *111*, 1346.

(22) The coupling of 1-bromo-4-(6-bromohexyl)benzene resulted in recovered starting material.

(23) The carboxylative event could be conducted in the dark with no erosion in yield, thus suggesting that light is not necessary for the reaction to occur.

(24) For recent metal-catalyzed hydrocarboxylation reactions, see: (a) Greenhalgh, M. D.; Thomas, S. P. *J. Am. Chem. Soc.* **2012**, *134*, 11900. (b) Williams, C. M.; Johnson, J. B.; Rovis, T. *J. Am. Chem. Soc.* **2008**, *130*, 14936.

(25) For related Pd-catalyzed events using isotopically labeled intermediates, see: (a) Monks, B. M.; Cook, S. P. *J. Am. Chem. Soc.* **2012**, *134*, 15297. (b) Netherton, M. R.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 3910. (c) Stokes, B. J.; Opra, S. M.; Sigman, M. S. *J. Am. Chem. Soc.* **2012**, *134*, 11408.

(26) Care must be taken when selecting the appropriate model substrate for isotopic labeling. Whereas the $J_{1,2}$ values in **7a** and **7c** are significantly different (Scheme 3), other related γ,γ -unsubstituted alkyl bromides such as (5-bromopentyl)benzene had similar $J_{1,2}$ values in both the *erythro* and *threo* isomers.

(27) In line with such a hypothesis, we found that radical scavengers such as TEMPO and BHT inhibited the reaction. Furthermore, aliphatic alcohols were obtained as byproducts when alkyl tosylates were employed, an observation that is consistent with a radical pathway. For example, see: (a) Madabhushi, S.; Kumar, B. A.; Narender, R. *Tetrahedron Lett.* **1998**, *39*, 2847. (b) Closson, W. D.; Wriede, P.; Bank, S. J. *J. Am. Chem. Soc.* **1966**, *88*, 1581 and citations therein.

(28) In the absence of CO₂, **7a–d** resulted predominantly in dimerization events and traces of β -hydride elimination.

(29) We cannot rule out the possibility that **L14** and **L16** act as a “redox-noninnocent ligands”. For example, see: Jones, G. D.; Martin, J. L.; McFarland, C.; Allen, O. R.; Hall, R. E.; Haley, A. D.; Brandon, R. J.;

Konovalova, T.; Desrochers, P. J.; Pulay, P.; Vicić, D. A. *J. Am. Chem. Soc.* **2006**, *128*, 13175.

(30) For the intermediacy of Ni species in odd oxidation states, see refs 5a, 5b, 6a, and the following: (a) Laskowski, C. A.; Bungum, D. J.; Baldwin, S. M.; Del Ciello, S. A.; Iluc, V. M.; Hillhouse, G. L. *J. Am. Chem. Soc.* **2013**, *135*, 18272. (b) Breitenfeld, J.; Ruiz, J.; Wodrich, M. D.; Hu, X. *J. Am. Chem. Soc.* **2013**, *135*, 12004. (c) Biswas, S.; Weix, D. J. *J. Am. Chem. Soc.* **2013**, *135*, 16192.

(31) For a mechanistic hypothesis, see the Supporting Information.