ChemComm

COMMUNICATION

View Article Online View Journal

NHC-catalysed aerobic aldehyde-esterifications with alcohols: no additives or cocatalysts required[†]

Cite this: DOI: 10.1039/c3cc42596g

Received 9th April 2013, Accepted 3rd June 2013 Eoghan G. Delany,^a Claire-Louise Fagan,^a Sivaji Gundala,^a Alessandra Mari,^a Thomas Broja,^b Kirsten Zeitler^{*bc} and Stephen J. Connon^{*a}

DOI: 10.1039/c3cc42596g

www.rsc.org/chemcomm

A highly efficient, broad scope, additive-free mild protocol for the oxidative carbene-catalysed esterification of aldehydes (including the related aqueous oxidation to acids) has been developed.

Esters are a general class of molecules which are so commonplace that their undoubted synthetic importance defies quantification.¹ The ester functional group is most commonly accessed *via* the *stoichiometric* activation of a carboxylic acid as an acyl halide, anhydride or activated ester with subsequent acyl transfer to an alcohol nucleophile.

Recently, interest has grown rapidly in the development of an alternative oxidative esterification of aldehydes with alcohols catalysed by *N*-heterocyclic carbenes (NHCs, **1**, Fig. 1).^{2,3} These transformations are purported to involve the addition of the carbene catalyst to an aldehyde **2** to form the Breslow intermediate **3**, which is then converted by a stoichiometric oxidant *e.g.* nitrobenzene,⁴ MnO₂,^{5,6} TEMPO,⁷ diphenoquinone,^{8,9} azobenzene,^{10,11} phenazine or riboflavin;¹² (the use of anodic electrochemical oxidation to replace the stoichiometric oxidant has also been very recently demonstrated¹³) to an electrophilic acyl azolium ion **4**. This species can then transfer the acyl group to an alcohol **5** to afford ester **6** and the carbene, which re-enters the catalytic cycle. The potential of this novel strategy as a synthetic tool in natural product synthesis has recently begun to be appreciated.^{5*b*,*e*}

In the presence of molecular oxygen in place of the stoichiometric oxidant it has been proposed that a separate oxidative fate for the Breslow intermediate involving addition of O_2 to **3** affords the internal peroxy zwitterion 7^{14} which then reacts with either the substrate aldehyde **2** (*via* peroxyacid intermediates)



Fig. 1 NHC-mediated aldehyde esterification using either added oxidant or molecular oxygen: current mechanistic rationales.

or another molecule of **3** to generate a carboxylate ion **8**;¹⁵ this can be trapped by either a proton or an alkyl halide 10^{14-17} to afford either the acid **9** or the ester **11** respectively.^{18,19} von Wangelin *et al.*^{2a} (Fig. 1) have recently classified these distinct NHC-mediated esterifications as being either 'oxidative' (*i.e.* the oxidation of **3** to **4** by an added oxidant) or 'oxygenative' (*i.e.* the conversion of **3** to **7** by O₂).²⁰ While significant progress has been made – a mild, additive-free NHC-mediated protocol of broad scope for the aerobic oxidation of aldehydes has yet to be developed.

During an investigation into the use of magnetite nanoparticles as a co-catalyst for the NHC-mediated oxidative esterification of *aldehydes*, we observed partial esterification of benzaldehyde (12) in methanol under aerobic conditions *in the absence of the nanoparticle*. After reproducing the result several times, we embarked on a study to determine the factors which influence the efficiency of this catalytic process (Table 1).

Preliminary experiments revealed that 12 could be esterified to a conveniently detectable extent in methanolic THF (1:1 v/v) in the presence of triazolium precatalysts (15 mol%) and a

^a Centre for Synthesis and Chemical Biology, Trinity Biomedical Sciences Institute, School of Chemistry, The University of Dublin, Trinity College, Dublin 2, Ireland. E-mail: connons@tcd.ie; Fax: +353 16712826

^b Institut für Organische Chemie, Universität Leipzig, D-04103 Leipzig, Germany. E-mail: kzeitler@uni-leipzig.de; Fax: +49 341-97-36599

^c Institut für Organische Chemie, Universität Regensburg, D-93053 Regensburg, Germany

[†] Electronic supplementary information (ESI) available: Experimental procedures and spectroscopic data for all new compounds. See DOI: 10.1039/c3cc42596g.

O precatalyst (x mol%) O									
		12	base (y THF/M rt, 24 h	eOH n, air	13				
	precatalysts								
	CI ⁻ 14				BF4 16	5			
Ent	try Precat.	x (mol%)	Base	y (mol%)	Solv. ^a (v/v)	Yield ^b (%)			
1	_	0	DBU	110	1:1	0			
2	14	15	DBU	110	1:1	48			
3	15	15	DBU	110	1:1	56			
4	16	15	DBU	110	1:1	61			
5	17	15	DBU	110	1:1	0			
6	18	15	DBU	110	1:1	96			
7 ^c	18	15	DBU	110	1:1	73			
8	18	5	DBU	110	1:1	74			
9	18	15	DBU	15	1:1	17			
10	18	15	DBU	50	1:1	54			
11	18	15	TEA	110	1:1	9			
12	18	15	DMAP	110	1:1	12			
13	18	15	DABCO	110	1:1	0			
14	18	15	K_2CO_3	110	1:1	60			
15	18	15	DBU	110	12:1	44			
16	18	15	DBU	110	6:1	61			
^{<i>a</i>} THF held constant (2.5 mL). ^{<i>b</i>} Determined by ¹ H NMR spectroscopy									

"THF held constant (2.5 mL). " Determined by "H NMR spectroscopy using styrene (114 μ L, 1 mmol, 1 equiv.) as an internal standard. " THF replaced with CH₂Cl₂ solvent.

small excess of base (DBU, 110 mol%) under an air atmosphere. We therefore began by examining the influence of precatalyst structure on the efficiency of the process. It is important to note that in the absence of triazolium ions, no conversion to methyl benzoate (13) is observed (entry 1). However, in the presence of the *N*-phenyl bicyclic triazolium ion 14 a moderate yield of 13 is formed after 24 h at ambient temperature (entry 2). The use of the more acidic carbene precursor 15^{21} led to a marginal improvement in product yield (entry 3), which increased further when the reaction was mediated by the hindered carbene derived from 16 (entry 4). An open-chain analogue of precatalyst 14 (*i.e.* 17) proved completely inactive under these conditions (entry 5). Modification of this structure *via* the exchange of the *N*-phenyl substituent for a methyl group (*i.e.* precatalyst 18) brought about a dramatic increase in activity (entry 6).

Given the mechanistic discussion outlined above (Fig. 1), this efficient, room temperature aerobic oxidative esterification in the absence of either an added stoichiometric oxidant, an alkyl halide or a dedicated catalyst for the activation of triplet oxygen^{22–24} is remarkable. We next proceeded to determine the influence of the other reaction components on the process. THF could be exchanged for CH_2Cl_2 , with an attendant loss in product yield (entry 7). Reduction of the loading of the triazolium ion to 5 mol% led to a similar reduction in efficacy (entry 8). We were also surprised to find that the oxidation process also strongly depends on both the loading (entries 9 and 10) and the identity (entries 11–14) of the base – with DBU Table 2 Reaction scope: aldehyde component

	R 18 (15 BBU (1 THF/Me rt,	5 mol%) 10 mol%) eOH (1:1) air	$\stackrel{O}{R} \stackrel{O}{\longleftarrow} OMe \underbrace{ \underbrace{ \begin{array}{c} 18 \\ 18 \\ 1 \\ 1 \\ 1 \\ 1 \end{array} }^{+} N \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\$	
Entry	Substrate	Time	Product	Yield ^a (%)
1		24 h	O OMe 13	94
2	0	12 h	O OMe 31	90
3	20	30 h	OMe 32	92
4	MeO 21	92 h	MeO 33	90
5	CI22	24 h	CI OMe	92
6		12 h	CI OMe	72
7 ^{<i>b</i>}	MeO 24	68 h	MeO OMe	77
8	25	24 h	O 37 OMe	28 ^c
9		24 h	CI O OMe 38	53 (53) ^d
10	CF3 0 27	24 h	CF ₃ O OMe	0
11	S 28	18 h	O S 40	$65(69)^d$
12	0 29	20 min	OMe 41	93
13	30	13	OMe 42	15 ^c

^{*a*} Yield of isolated product. ^{*b*} Reaction at 45 °C. ^{*c*} Determined by ¹H NMR spectroscopy using styrene (114 μ L, 1 mmol, 1 equiv.) as an internal standard. ^{*d*} The value in parenthesis is the yield of isolated product obtained when this reaction was repeated in the collaborating research group using independently synthesised/purified materials and precatalyst.

emerging comfortably superior to the others evaluated in this study. The esterification can also be carried out using considerably less MeOH (entries 15 and 16), at the expense of rate.

With a useful protocol in hand, we next investigated the compatibility of the new process with different aldehydes (Table 2). Electron-neutral (*i.e.* **12** and **19**, entries 1 and 2)

and deactivated (*i.e.* **20** and **21**, entries 3 and 4) aromatic aldehydes could be converted to the corresponding methyl esters in excellent isolated yield in the presence of **18**, DBU, methanol and air. Activated benzaldehydes provided interesting results: *m*-chlorobenzaldehyde (**22**) proved an excellent substrate, while the esterification of its *para*-substituted isomer **23** and a *m*-methoxy-substituted variant (*i.e.* **24**) proceeded in 77–92% yields (entries 5–7). *Ortho*-substitution is not well tolerated: *o*-tolualdehyde (**25**) and *o*-chlorobenzaldehyde (**26**) underwent conversion to **37** and **38** in low to moderate yields respectively (entries 8 and 9), while *o*-trifluoromethyl-benzaldehyde was inert (entry 10).

Substrates incorporating oxidisable heterocyclic functionality (*i.e.* **28** and **29**) could be converted to their methyl ester analogues in good-excellent yields (entries 11 and 12).²⁵ In contrast, the aliphatic aldehyde **30** resulted in a poor yield of the ester **42** and a complex product mixture (entry 13).

In summary, we have developed an efficient NHC-catalysed esterification of aldehydes involving alcohols and air (*i.e.* oxygen) as the oxidant. No other added stoichiometric oxidants or catalysts to activate molecular oxygen are required. Unhindered aromatic aldehydes (including heterocyclic analogues) can be converted to the corresponding methyl esters in good to excellent yields at ambient temperature.

Financial support from IRCSET, Science Foundation Ireland and the DFG is gratefully acknowledged.

Notes and references

- 1 (a) J. Otera, Esterification: Methods, Reactions and Applications, Wiley, New York, 2003; (b) R. C. Larock, Comprehensive Organic Transformations, VCH, New York, 1989.
- 2 For reviews on NHC-catalysed oxidative transformations see: (*a*) C. E. I. Knappke, A. Imami and A. Jacobi von Wangelin, *ChemCatChem*, 2012, 4, 937; (*b*) S. DeSarkar, A. Biswas, R. C. Samanta and A. Studer, *Chem.-Eur. J.*, 2013, **19**, 4664.
- 3 Recent reviews concerning various aspects of NHC organocatalysis:
 (a) H. U. Vora and T. Rovis, Aldrichimica Acta, 2011, 44, 3; (b) A. T. Biju, N. Kuhl and F. Glorius, Acc. Chem. Res., 2011, 44, 1182;
 (c) V. Nair, R. S. Menon, A. T. Biju, C. R. Sinu, R. R. Paula, A. Josea and V. Sreekumar, Chem. Soc. Rev., 2011, 40, 5336;
 (d) P.-C. Chiang and J. W. Bode, RSC Catal. Ser., 2010, 6, 399;
 (e) C. D. Campbell, K. B. Ling and A. D. Smith, in Catalysis by Metal Complexes, 2010, vol. 32, p. 263; (f) J. L. Moore and T. Rovis, Top. Curr. Chem., 2009, 291, 77; (g) E. M. Phillips, A. Chan and K. A. Scheidt, Aldrichimica Acta, 2009, 42, 55; (h) T. Rovis, Chem. Lett., 2008, 1; (i) K. Zeitler, Ernst Schering Found. Symp. Proc., 2007, 2, 183;
 (j) D. Enders, O. Niemeier and A. Henseler, Chem. Rev., 2007, 107, 5506;
 (k) N. Marion, S. Diez-Gonzalez and S. P. Nolan, Angew. Chem., Int. Ed., 2007, 46, 2988; (l) K. Zeitler, Angew. Chem., Int. Ed., 2005, 44, 7506.
- 4 (a) A. Miyashita, Y. Suzuki, M. Kobayshi, N. Kuriyama and T. Higashino, *Heterocycles*, 1996, 43, 509; (b) A. Miyashita, Y. Suzuki, I. Nagsaki, C. Ishiguro, K.-I. Iwamoto and T. Higashino, *Chem. Pharm. Bull.*, 1997, 45, 1254; (c) J. Castells, F. Pujol, H. Llitjós and M. Moreno-Mańas, *Tetrahedron*, 1982, 38, 337.
- 5 (a) B. E. Maki and K. A. Scheidt, Org. Lett., 2008, 10, 4331; (b) B. Maki,
 A. Chan and K. A. Scheidt, Synthesis, 2008, 1306; (c) H. Kim, Y. Park
 and J. Hong, Angew. Chem., Int. Ed., 2009, 48, 7577; (d) B. Maki,
 A. Chan, E. M. Philips and K. A. Scheidt, Tetrahedron, 2009, 65, 3102;
 (e) K. Lee, H. Kim and J. Hong, Angew. Chem., Int. Ed., 2012, 51, 5735.
- 6 Related reactions mediated by MnO₂: (a) B. E. Maki, A. Chan, E. M. Phillips and K. A. Scheidt, Org. Lett., 2007, 9, 371; (b) M. Rueping, H. Sundén, L. Hubener and E. Sugiono, Chem. Commun., 2012, 48, 2201.
- 7 J. Guin, S. De Sarkar, S. Grimme and A. Studer, *Angew. Chem., Int. Ed.*, 2008, **47**, 8727.

- 8 J. Guin, S. De Sarkar, S. Grimme and A. Studer, J. Am. Chem. Soc., 2010, 132, 1190.
- 9 For examples of related reactions not involving ester formation see: (*a*) S. De Sarkar and A. Studer, *Org. Lett.*, 2010, **12**, 1992; (*b*) S. De Sarkar and A. Studer, *Angew. Chem., Int. Ed.*, 2010, **49**, 9266; (*c*) A. Biswas, S. De Sarkar, R. Fröhlich and A. Studer, *Org. Lett.*, 2011, **13**, 4966.
- 10 (a) H. Inoue and K. Higashiura, J. Chem. Soc., Chem. Commun., 1980, 549; (b) C. Noonan, L. Baragwanath and S. J. Connon, *Tetrahedron Lett.*, 2008, **49**, 4003; (c) C. A. Rose and K. Zeitler, Org. Lett., 2010, **12**, 4552.
- 11 For related NHC-mediated oxidations see: (a) J. Castells, H. Llitjós and M. Moreno-Mańas, *Tetrahedron Lett.*, 1977, **18**, 205; (b) H. Inoue and S. J. Tamura, J. Chem. Soc., Chem. Commun., 1985, 141; (c) S. Shinkai, T. Yamashita, Y. Kusano and O. Manabe, J. Org. Chem., 1980, **45**, 4947; (d) S. Shinkai, T. Yamashita, Y. Kusano and O. Manabe, *Tetrahedron Lett.*, 1980, **21**, 2543; (e) S. Shinkai, Y. Hara and O. Manabe, *Bull. Chem. Soc. Jpn.*, 1983, **56**, 770; (f) Y. Yano, Y. Hoshino and W. Takagi, Chem. *Lett.*, 1980, 749; (g) Y. Yano and Y. Tsukagoshi, J. Chem. Res., Synop., 1984, 406; (h) W. H. Rastetter, J. Adams, J. W. Frost, L. J. Nummy, J. E. Frommer and K. B. Roberts, J. Am. Chem. Soc., 1979, **101**, 2752; (i) W. H. Rastetter and J. Adams, J. Org. Chem., 1981, **46**, 1882; (j) D. Hilvert and R. Breslow, *Bioorg. Chem.*, 1984, **12**, 206; (k) L. Jimenez and F. Diederich, *Tetrahedron Lett.*, 1989, **30**, 2759.
- 12 (a) S.-W. Tam, L. Jimenez and F. Diederich, J. Am. Chem. Soc., 1992, 114, 1503; (b) T. Uno, T. Inokuma and Y. Takemoto, Chem. Commun., 2012, 48, 1901; (c) Rovis et al. have noted the application of catalytic oxidant where imine substrates are supposed to re-oxidize riboflavin tetraacetate: X. Zhao, K. E. Ruhl and T. Rovis, Angew. Chem., Int. Ed., 2012, 51, 12330.
- 13 E. E. Finney, K. A. Ogawa and A. J. Boydston, *J. Am. Chem. Soc.*, 2012, 134, 12374.
- 14 L. Lin, Y. Li, W. Du and W.-P. Deng, *Tetrahedron Lett.*, 2010, **51**, 3571. 15 B. Maji, S. Vedachalan, X. Ge, S. Cai and X.-W. Liu, *J. Org. Chem.*,
- 2011, 76, 3016.
- 16 Y.-C. Xin, S.-H. Shi, D.-D. Xie, X.-P. Hui and P.-F. Xu, Eur. J. Org. Chem., 2011, 6527.
- 17 For a related process involving intramolecular cyclisation of the carboxylate ion onto an alkyne see: J. H. Park, S. V. Bhilare and S. W. Youn, *Org. Lett.*, 2011, **13**, 2228.
- 18 Recent claims that CO₂ can be involved in NHC-mediated oxidations of aldehydes as an oxidant (ref. 18*a* and *b*) were subsequently investigated by Bode *et al.* (ref. 18*c*), who ascertained that O₂ is the actual oxidant in these processes, with CO₂ serving an ancillary role as a suppressant of side-reactions. (*a*) L. Gu and Y. Zhang, *J. Am. Chem. Soc.*, 2010, 132, 914; (*b*) V. Nair, V. Varghese, R. R. Paul, A. Jose, C. R. Sinu and R. S. Menon, *Org. Lett.*, 2010, 12, 2653; (*c*) P.-C. Chiang and J. W. Bode, *Org. Lett.*, 2011, 13, 2422; for a NHC-catalysed aerobic oxidation of ArCHO to their acids using "Bode-type" conditions:; (*d*) W. Yang, G.-Z. Gou, Y. Wang and W.-F. Fu, *RSC Adv.*, 2013, 3, 6334.
- 19 Du and Deng (ref. 14) have proposed an alternative mechanism for oxygenative esterification involving *O*-alkylation of the Breslow intermediate by the alkyl halide electrophile.
- 20 Intermediate 7 has also been proposed to for a recent related NHC-catalysed esterification under O₂-atmosphere: I. N. C. Kiran, K. Lalwani and A. Sudalai, *RSC Adv.*, 2013, **3**, 1695.
- 21 (a) M. S. Kerr, J. Read de Alaniz and T. Rovis, J. Org. Chem., 2005, 70, 5725; (b) H. U. Vora, S. P. Lathrop, N. T. Reynolds, M. S. Kerr, J. Read de Alaniz and T. Rovis, Org. Synth., 2010, 87, 350; (c) The pK_a of catalyst 14 (H₂O) has been determined to be 17.7: C. D. Campbell, N. Duguet, K. A. Gallagher, J. E. Thomson, A. G. Lindsay, A. C. O'Donoghue and A. D. Smith, Chem. Commun., 2008, 3528; (d) For a recent compilation of more pK_a values of triazolium salts: R. S. Massey, C. J. Collett, A. G. Lindsay, A. D. Smith and A. C. O'Donoghue, J. Am. Chem. Soc., 2012, 134, 20421.
- 22 S. Iwahana, H. Iida and E. Yashima, Chem.-Eur. J., 2011, 17, 8009.
- 23 For a recent flavin-mediated aerobic oxidation of 12 in a Dakin-type process see: S. Chen and F. W. Foss Jr., *Org. Lett.*, 2012, 14, 5150.
 24 For a recent report concerning the use of Fe²⁺ ions as a co-catalyst
- 24 For a recent report concerning the use of Fe²⁺ ions as a co-catalyst and coupling with phenols see: R. S. Reddy, J. N. Rosa, L. F. Veiros, S. Caddick and P. M. P. Gois, *Org. Biomol. Chem.*, 2011, 9, 3126.
- 25 In this context it is interesting to note that Goswami and Hazra had previously reported the aerobic esterification of heterocyclic aromatic aldehydes (only) involving thiamine hydrochloride and NEt₃ (2.0 equiv.) in MeOH (heating under reflux was required): S. Goswami and A. Hazra, *Chem. Lett.*, 2009, 484.