# K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O supported on silica gel: An efficient and selective reagent for the cleavage of oximes to their corresponding carbonyl compounds in aqueous medium

ABBAS AMINI MANESH\* and BEHZAD SHIRMARDI SHAGHASEMI

Department of Chemistry, Payame Noor University, 19395-4697 Tehran, I. R. of IRAN e-mail: a\_aminima@yahoo.com

MS received 5 June 2014; revised 27 August 2014; accepted 29 August 2014

**Abstract.** K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O supported on silica gel, a new oxidant, for efficient, simple and selective cleavage of ketoximes and aldoximes to their corresponding carbonyl compounds in aqueous medium is described. Further oxidation of aldehydes to carboxylic acid and formation of by-products were not observed.  $\alpha - \beta$  unsaturated oxime was deoximated smoothly without oxidation of the double bond.

Keywords. K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O, Silica gel; oxime; aldehyde; ketone.

# 1. Introduction

Oximes can be prepared from carbonyl and noncarbonyl compounds, and compounds having active methylene groups.<sup>1,2</sup> These solid compounds are used for isolation, purification and characterization of the parent carbonyl compounds.<sup>3</sup> Carbonyl groups can be protected by oximation, and in turn, these oximes can serve as an intermediate for the preparation of nitriles<sup>4</sup> of amides via the Beckmann rearrangement or to activate the carbonyl group.<sup>5</sup> Thus, there has been an increasing interest in the development of methods for the conversion of oximes into their corresponding carbonyl compounds and a number of methods and compounds such as 1,3-Dibromo-5,5dimethylhydantoin,<sup>6</sup> microwave,<sup>7</sup> CeO<sub>2</sub>-ZrO<sub>2</sub>,<sup>8</sup> Potassium bromide and Ammonium heptamolybdate,<sup>9</sup> Silica Sulfuric acid,<sup>10</sup> Bi(OTf)<sub>3</sub>·4H<sub>2</sub>O,<sup>11</sup> Ionic liquid,<sup>12</sup> poly[4-vinyl-N, N-dichlorobenzenesulfonamide],<sup>13</sup> metalloporphyrins,<sup>14</sup> 1,3-Dichloro-5,5-dimethylhydantoin,<sup>15</sup> 2-Nitro-4,5-dichloropyridazin-3(2H)-one,<sup>16</sup> N-bromo-*N*-benzoyl-4-toluenesulfonamide,<sup>17</sup> Tetra-n-alkylammonium bromates<sup>18</sup> have been explored.

In a nutshell, the regeneration of carbonyl compounds from oximes can be achieved using a variety of methods, including oxidative, reductive, and photochemical techniques.<sup>19</sup> In continuation of our systematic study and research on oxidation methods,<sup>20–24</sup> we report a new, simple and selective method for the conversion of aliphatic, aromatic and cyclic oximes to their corresponding aldehydes and ketones by  $K_3[Fe(CN)_6].3H_2O$  supported on SiO<sub>2</sub> as a new oxidant (scheme 1).

### 2. Experimental

### 2.1 General methods and materials

In this study, all of the oximes were prepared as reported elsewhere,<sup>25</sup> except benzaldehyde oxime, which purchases from Merck (Darmstadt, Germany). IR and <sup>1</sup>H NMR spectra were recorded using a Shimadzu Infrared Spectrophotometer FT-IR Model IR Prestige 21 (KBr pellets) and a 90 MHz Jeol FT-NMR spectrometer, respectively. <sup>1</sup>H NMR chemical shifts were measured relative to TMS. Silica-supported K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O (oxidant B) was prepared by impregnating 2.0 g of silica with aqueous solutions of K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O (1 g in 30 mL H<sub>2</sub>O). The mixture was stirred overnight at reflux temperature, followed by drying using a rotary evaporator at 110°C which gives yellow powders.

### 2.2 General Procedure for Deoximation

To a solution of oxime (1 mmol) in ethanol (10 mL), 0.1 g oxidant B (0.393 mmol) was added. Then H<sub>2</sub>O (1 mL) was added to the solution. The resulting mixture was stirred at reflux conditions and the progress of the reaction monitored by thin layer chromatography (hexane-acetone, 8:2 v/v). After completion of the reaction, the oxidant was removed by filtration and washed with  $2 \times 5$  mL ethanol. The solvent was removed from

<sup>\*</sup>For correspondence



R1, R<sub>2</sub> = Aromatic , Aliphatic, Cyclic, H

Scheme 1. Cleavage of oximes by  $K_3[Fe(CN)_6].3H_2O$  supported on SiO2.

the filtrate and the product was purified by column chromatography.

### 3. Results and Discussion

In order to optimize the reaction conditions, first of all, 0.2 g oxidant A (0.787 mmol  $K_3$ [Fe(CN)<sub>6</sub>].3H<sub>2</sub>O) was added to a solution of acetophenone oxime (1 mmol, 0.135 g) in acetonitrile (10 mL). According to the proposed mechanism, H<sub>2</sub>O (1 mL) was added to the solution (scheme 2). The resulting mixture was stirred at reflux conditions (the reaction did not proceed after prolonged time at room temperature) and the progress of the reaction was monitored by thin layer chromatography (hexane-acetone, 8:2 v/v). Several trials of oxidant A were tried to find the suitable amount of oxidant for the reaction (table 1).

During the optimization of the reaction conditions, we compared the effect of different solvents on the rate and yields of the reactions. Our findings showed that EtOH was generally the best of the solvents tested in terms of yield and time (table 2).

After using oxidant A in EtOH as the suitable solvent, we conducted the reaction for different amounts of oxidant B and oxidant C (table 3). Having analyzed these results, we conclude that the best reaction conditions were achieved when we used 0.1 g of oxidant B in EtOH media in reflux condition. It is noteworthy that, addition of more  $H_2O$  to the reaction mixture, does not improve the reaction.

Different conditions for the preparation of oxidant A, B and C are shown in table 3.

### 3.1 Characterization of the oxidant

The XRD profiles of the samples at 298 K are shown in figure 1. Powder X-ray diffraction (XRD) patterns were



Scheme 2. Optimization of the reaction in acetonitrile.

Table 1.Optimization of Oxidant A.

Entry	Oxidant A (g)	Conditions	Time (h)	Yield (%)
1	0.2	r.t.	5	0
2	0.2	reflux	2.5	73
3	0.1	reflux	3.5	65
4	0.3	reflux	3.45	57
5	0.25	reflux	2.5	85

Table 2.Optimization of solvent.

Entry	Solvent	Time (h)	Yield (%)
1	CH <sub>3</sub> CN	2:30	85
2	EtOH	2:15	91
3	CH <sub>2</sub> Cl <sub>2</sub>	5	0
4	Hexan	5	0
5	MeOH	2:30	86

**Table 3.**Preparation of oxidant.

Oxidant	$SiO_2(g)$	$K_3[Fe(CN)_6].3H_2O(g)^8$
A	2	2
B C	2 2	1 0.5

<sup>a</sup>K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O was dissolved in 30 mL of H<sub>2</sub>O.



**Figure 1.** XRD pattern of A:  $SiO_2$ , B:  $K_3[Fe(CN)_6]$ .3H<sub>2</sub>O supported on silica gel.



**Figure 2.** IR (KBr) Spectrum of  $K_3[Fe(CN)_6]$ .3H<sub>2</sub>O supported on silica gel.

Entry	Oxime	Product	Time (min)	Yield (%) <sup>a,b</sup>
	Г <sup>N</sup> _ОН	<b>1</b> 0		
1	$\bigcirc$		50	95
	Мон			
2	q	CI	50	90
	Лон	C C C C C C C C C C C C C C C C C C C		
3	$\square$	$\square$	45	95
	С ОН	, o	UT UT	75
4	$\square$	$\square$		
	о Он	0. 	45	95
5	$\bigcirc$	$\bigcirc$		
	NO₂   <sup>N</sup> _OH	No <sub>2</sub>	65	87
6	$\bigcirc$	$\bigcirc$		
	I Br	l Br O	50	90
7			55	90
	N OH	L'a		
8	N <sup>_OH</sup>	Ŷ	60	86
9			75	93
10			105	85
10	HON	~ ~	105	05
11		СТ Сн I	_	_
12	Ň	× Co	_	_
	OH NOH	OH O		
13			_	_

 Table 4.
 The results of the conversion of oximes to the carbonyl compounds by oxidant.

<sup>a</sup> Products were characterized by their physical constants, comparison with authentic samples and melting points of 2,4-dinitro phenyl hydrazone derivatives and IR and NMR spectra; <sup>b</sup>Isolated yields

recorded on a Philips PW-1840 X-ray diffractometer (XRD) with Cu K $\alpha$  radiation. The intensity data were collected over a  $2\theta$  range of 10–90°. The XRD results revealed the presence of K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O over amorphous SiO<sub>2</sub> matrix in the case of K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O-SiO<sub>2</sub> (see  $2\theta = 20 - 40^{\circ}$ ), and some overlapped regions.

The particles size of the  $K_3[Fe(CN)_6].3H_2O$  on  $SiO_2$  is 82 nm (figure 1). The main features of

 $K_3$ [Fe(CN)<sub>6</sub>].3H<sub>2</sub>O supported on silica gel are connected to the similarity of bands associated with the inorganic backbone including (i) a broad band between 3400 and 3200 cm<sup>-1</sup>, which is assigned to the O-H stretching mode of silanol groups and also to the remaining adsorbed water, (ii) the intense band related to siloxane stretching of these groups at 1100 cm<sup>-1</sup>, (iii) a band assigned to Si-O stretching mode of the silanol group at 800 cm<sup>-1</sup> and (iv) the band around 1650 cm<sup>-1</sup>,



Scheme 3. Proposed mechanism of deoximation by the oxidant.

assigned to the water bending mode, and (v) an intense band assigned to CN band at  $2117 \text{ cm}^{-1}$  (figure 2).

### 3.2 Proposed mechanism

Various methods were adopted for the deoximation processes to effectively and suitably obtain the parent or new aldehydes and ketones. Deoximation can be achieved using various processes like oxidation, hydrolysis, hydrogenation, etc., using organic and inorganic reagents. Besides, some surface templated processes and photochemical processes were also used to give products in a remarkably effective way.<sup>26</sup>

Mechanism of deoximation by the  $K_3[Fe(CN)_6]$ .  $3H_2O$  supported on silica gel is similar to other electron transfer type mechanisms that are described elsewhere.<sup>27</sup> Proposed mechanism for deoximation of variety of oximes to their corresponding aldehydes and ketones (table 4) is shown in scheme 3. The first step is simply the formation of the radical cation (I) of oxime by electron transfer from oxime to the potassium ferricyanide. Alternatively, a sequence can be written involving removal of a proton from (I) followed oxidation of iminoxyl radical (II) or nitroso radical (III) by potassium ferricyanide to give the nitroso carbocation (IV).

Hydration of the carbocation (**IV**) produces the intermediate (**V**), and finally decomposition of the (**VI**) produces carbonyl compounds.

# 4. Conclusion

It is noteworthy that in this study, over oxidation of aldehydes to carboxylic acid and formation of byproducts were not observed.  $\alpha$ -  $\beta$  unsaturated oxime (Entry 7) was deoximated smoothly without oxidation of the double bond. Furthermore, acid sensitive groups (Entry 4 and 8) remained unaffected during the oxidation reaction. The water insolubility of the reagent and by-products made product recovery easy. Due to sterical hindrance, benzoin oxime (Entry 11), Camphor oxime (Entry 12), and 2-hydroxy acetophenone oxime (Entry 13) could not be converted to their corresponding carbonyl compounds. Also, the reactions did not proceed using K<sub>3</sub>[Fe(CN)<sub>6</sub>].3H<sub>2</sub>O or SiO<sub>2</sub> alone, even after a prolonged period of time. The striking features of our method are short reaction times, heterogeneous reaction conditions, absence of formation of over oxidation products due to high selectivity and mild nature of oxidant, easy workup procedure, high yields and preservation of carboncarbon double bond functional group in the oxime structure.

### **Supplementary Information**

The electronic supporting information can be seen at www.ias.ac.in/chemsci.

## Acknowledgement

The authors acknowledge the support of the Hamedan Payame Noor University to this work.

### References

- Barhate N B, Gajare A S, Wakharkar R D and Sudalai A 1997 Tetrahedron Lett. 38 653
- 2. Hajipour A R, Mohammadpoor-Baltork I, Nikbaghat K and Imanzadeh G 1999 *Synth. Commun.* **29** 1697
- 3. Bosch A I, De La Crus P, Diez-Barra E, Loupy A and Langa F 1995 *Synlett* 1259
- Varma R S, Dahiya R and Saini R K 1997 Tetrahedron Lett. 38 8819
- 5. Kabalka G W, Pace R D and Wadgaonkar P P 1990 Synth. Commun. 20 2453
- Khazaei A and Rostami A 2006 Org. Prep. Proced. Int. 38 484
- 7. Khazaei A and Amin Manesh A 2005 J. Braz. Chem. Soc. 16 874
- 8. Deshpande S S, Sonavane S U and Jayaram R V 2008 *Catal. Commun.* **9** 639
- 9. Ganguly N C and Barik S K 2008 Synthesis 425
- Li J T, Sun M X and Yin Y 2010 Ultrason. Sonochem. 17 359
- 11. Arnold J N, Hayes P D, Kohaus R L and Mohan R S 2003 *Tetrahedron Lett.* **44** 9173
- Li D, Shi F, Guo S and Deng Y 2004 Tetrahedron Lett. 45 265

- 13. Khazaei A and Vaghei R G 2002 Tetrahedron Lett. 43 3073
- Zhou X-T, Yuan Q-L and Ji H-B 2010 Tetrahedron Lett. 51 613
- Khazaei A and Amini Manesh A 2005 J. Chin. Chem. Soc. 52 1017
- 16. Kim B R, Lee H G, Kim E J, Lee S G and Yoon Y J 2010 *J. Org. Chem.* **75** 484
- Khazaei A and Amini Manesh A 2006 Mendeleev Commun. 16 109
- 18. Nath U, Sandhy Das S, Deb D and Jyoti Das P 2004 *New J. Chem.* **28** 1423
- 19. Lijser H J P, Fardoun F H, Sawyer J R and Quant M 2002 Org. Lett. 4 2325
- 20. Khazaei A, Amini Manesh A and Rostami A 2005 J. Chem. Res. 6 391
- 21. Khazaei A and Amini Manesh A 2005 J. Braz. Chem. Soc. 16 874
- 22. Khazaei A, Amini Manesh A and Rostami A 2004 J. Chem. Res. 10 695
- 23. Khazaei A, Amini Manesh A and Rostami A 2004 Phosphorus, Sulfur Silicon Relat. Elem. **179** 2483
- 24. Khazaei A and Amini Manesh A 2005 Synthesis 12 1929
- 25. Khoramabadi-Zad A, Azadmanesh M and Rezaee A 2010 S. Afr. J. Chem. 63 192
- 26. Sahu S, Sahu S, Patel S, Dash S and Mishra B K 2008 Indian J. Chem. **47B** 259
- 27. Shevlin P B and Neufeld H A 1970 *J. Org. Chem.* **35** 2178