## Palladium-Catalyzed Aerobic Oxidative Carbonylation of **Arylboronate Esters under Mild Conditions\*\***

Qiang Liu, Gang Li, Jun He, Jing Liu, Peng Li, and Aiwen Lei\*

The transition-metal-catalyzed carbonylation involving CO gas is a fundamental chemical transformation, which not only extends the carbon chain length, but also introduces a synthetically versatile carbonyl group. Theoretically, as shown in Scheme 1, the Ar group could originate from an



Scheme 1. Two theoretical aryl carbonylation pathways (ArM; M = H, In).

electrophile (ArX; Path I) or a nucleophile (ArM; Path II). The carbonylation of organic halides (ArX) with CO, pioneered by Heck and co-workers,<sup>[1a]</sup> has attracted broad interest in the past 40 years, and is widely applied to the synthesis of fine chemicals in industry.<sup>[1-3]</sup> In the reaction systems shown in Scheme 1, the  $\pi$ -acceptor property of CO renders low-valent metal catalysts such as Pd<sup>0</sup> species to be relatively electron deficient, and makes the oxidative addition of organic halides towards Pd<sup>0</sup> species difficult.<sup>[4]</sup> Consequently, the reaction in Path I usually requires relatively harsh conditions; for example, high temperatures or high CO pressures.<sup>[5,6]</sup>

Recently, oxidative carbonylation of ArM (M = H, In)with CO has emerged as a conceptually new alternative to "classic carbonylation processes".<sup>[7-11]</sup> The use of CO avoids the reluctant oxidative addition step, which is mentioned above in traditional carbonylations, and has the potential to accommodate milder reaction conditions. The oxidative carbonylation of ArH shows promise when catalyzed by

| [*]       | Q. Liu, G. Li, J. He, Dr. J. Liu, P. Li, Prof. A. Lei<br>The College of Chemistry and Molecular Sciences<br>Wuhan University, Wuhan, Hubei (China)<br>Fax: (+86) 27-6875-4067<br>E-mail: aiwenlei@whu.edu.cn<br>Homepage: http://www.chem.whu.edu.cn/GCI_WebSite/<br>main.htm |
|-----------|---|
|           | Prof. A. Lei<br>State Key Laboratory for Oxo Synthesis and Selective Oxidation,<br>Lanzhou Institute of Chemical Physics<br>Chinese Academy of Sciences, 730000 Lanzhou (China)   |
| [**]      | This work was supported by the National Natural Science<br>Foundation of China (20772093, 20972118, and 20832003), and the<br>Doctoral Fund of the Ministry of Education of China (20060486005).  |
| ( 100 k - |   |

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201000460.

transition metals, whereas the current transformation usually requires the use of directing groups.<sup>[7-10]</sup> Furthermore, the oxidative carbonylation of arylindium derivatives requires the use of stoichiometric amounts of the oxidant desyl chloride.[11] Herein, we report the first example of the oxidative carbonylation of arylboronic acid derivatives under balloon pressure of CO with air as the oxidant at  $40 \rightarrow 50$  °C.

Arylboronic acid derivatives are air and moisture stable, and are compatible with a broad range of common functional groups.<sup>[12–14]</sup> For these reasons, they have been widely applied in chemical syntheses.<sup>[15-17]</sup> Recently, the research groups of Hou<sup>[18]</sup> and Iwasawa<sup>[19]</sup> have reported a nucleophilic addition of arylboronate esters toward CO<sub>2</sub> in the presence of copper or rhodium catalysts, respectively. In the presence of CO, the Suzuki carbonylation reaction was also realized by Beller and co-workers, and resulted in the formation of biaryl ketones.<sup>[20]</sup> However, there is no literature precedent of their oxidative carbonylation to form carboxylate esters. Furthermore, with respect to cost and simplicity, air is the ideal oxidant for oxidative carbonylation.<sup>[21-26]</sup> The oxidative carbonylation of amines or alcohols was investigated by employing air or oxygen as the oxidant.<sup>[22,27-32]</sup> However, to the best of our knowledge, no example has been reported regarding the oxidative carbonylation of arylmetal reagents by employing oxygen or air as the sole oxidant.

To probe the feasibility of our proposed study, we chose the model substrates phenylBneop (1a;  $neop = OCH_2C$ - $(CH_3)_2CH_2O$  and *n*-butanol to afford the carbonylation product 2a, oxidative protodeboronation product 3a and homocoupled product 4a. Upon variation of the reaction conditions, different product distributions were observed (Table 1). Among the palladium catalysts used as precursors (Table 1, entries 1–11), [PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>] showed the best result (56% yield for 2a and 44% for 3a; Table 1, entry 7), whereas the bidentate ligands gave poor results (Table 1, entries 1, 2, and 6). Interestingly,  $Pd(OAc)_2/PPh_3$  (1:2) and  $[Pd(dba)_2]/$  $PPh_3$  (1:2) were not effective catalysts for this transformation (Table 1, entries 10 and 11). Using [PdCl<sub>2</sub>(MeCN)<sub>2</sub>]/PPh<sub>3</sub> (1:2) was equivalent to  $[PdCl_2(PPh_3)_2]$ , whereas  $[PdCl_2 (MeCN)_2$ /PPh<sub>3</sub> (1:4) was less effective at producing the carbonylation product 2a, however, it was an efficient catalytic system for the oxidative homocoupling of 1a (Table 1, entries 8 and 9).

Next, different bases were then studied (Table 1, entries 12–19). The results of triethylamine  $(\mathbf{B}^1)$  were similar to those of 2,2,6,6-tetramethylpiperidine  $(\mathbf{B}^2)$  and 2,6-lutidine  $(\mathbf{B}^3)$ , however, other bases tested were less effective in this reaction. Furthermore, the ratio of air to CO was also screened. Interestingly, the yield of *n*-butyl benzoate (2a) improved with an increase in the ratio of air to CO (Table 1,

Angew. Chem. Int. Ed. 2010, 49, 3371-3374

© 2010 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

the

## Communications

Table 1: Oxidative carbonylation of phenylboronate ester 1 a.<sup>[a]</sup>

|       |   | Pd](5 mol %)                   | PhCO <sub>2</sub> <i>n</i> Bu | + Pł | י<br>ח H + | PhPh              |
|-------|---|--------------------------------|-------------------------------|------|------------|-------------------|
|       |   | , 40 °C, 24 h                  | 111002/100                    |      | - non - n  |                   |
| 1a    |   | ase (2 equiv)                  | 2a                            | 3    | Ba         | 4a                |
| Entry | [Pd] Catalyst   | Base                           | Air/CO                        | Yi   | eld [%     | 5] <sup>[b]</sup> |
|       |   |                                |                               | 2a   | 3 a        | 4 a               |
| 1     | $Pd(OAc)_2/L^1$   | Et₃N                           | 1:3                           | 21   | 46         | 0                 |
| 2     | [PdCl <sub>2</sub> (dppf)]                                    | Et <sub>3</sub> N              | 1:3                           | 0    | 0          | 0                 |
| 3     | [PdCl <sub>2</sub> (MeCN) <sub>2</sub> ]/L <sup>2</sup> (1:2) | $Et_3N$                        | 1:3                           | 0    | 0          | 0                 |
| 4     | [PdCl <sub>2</sub> (MeCN) <sub>2</sub> ]/L <sup>3</sup> (1:2) | $Et_3N$                        | 1:3                           | 0    | 0          | 0                 |
| 5     | Pd(OAc) <sub>2</sub> /py (1:4)                                | Et <sub>3</sub> N              | 1:3                           | 0    | 0          | 0                 |
| 6     | Pd(OAc) <sub>2</sub> /bpy                                     | $Et_3N$                        | 1:3                           | 0    | 0          | 0                 |
| 7     | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | Et₃N                           | 1:3                           | 56   | 44         | 0                 |
| 8     | [PdCl <sub>2</sub> (MeCN) <sub>2</sub> ]/PPh <sub>3</sub> (1  | :4) Et₃N                       | 1:3                           | 0    | 12         | 80                |
| 9     | [PdCl <sub>2</sub> (MeCN) <sub>2</sub> ]/PPh <sub>3</sub> (1) | :2) Et₃N                       | 1:3                           | 56   | 44         | 0                 |
| 10    | Pd(OAc) <sub>2</sub> /PPh <sub>3</sub> (1:2)                  | Et₃N                           | 1:3                           | 26   | 34         | 0                 |
| 11    | [Pd(dba) <sub>2</sub> ]/PPh <sub>3</sub> (1:2)                | Et <sub>3</sub> N              | 1:3                           | 0    | 0          | 0                 |
| 12    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | <i>n</i> BuONa                 | 1:3                           | 32   | 68         | 0                 |
| 13    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | CsF                            | 1:3                           | 32   | 67         | 0                 |
| 14    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | K <sub>3</sub> PO <sub>4</sub> | 1:3                           | 45   | 53         | 0                 |
| 15    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | dbu                            | 1:3                           | 47   | 50         | 0                 |
| 16    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | dabco                          | 1:3                           | 0    | 0          | 0                 |
| 17    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | B <sup>1</sup>                 | 1:3                           | 51   | 47         | 0                 |
| 18    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | B <sup>2</sup>                 | 1:3                           | 54   | 42         | 0                 |
| 19    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | B <sup>3</sup>                 | 1:3                           | 54   | 38         | 0                 |
| 20    | $[PdCl_2(PPh_3)_2]$   | $Et_3N$                        | 1:10                          | 43   | 30         | 0                 |
| 21    | [PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]         | Et₃N                           | 1:5                           | 47   | 36         | 0                 |
| 22    | $[PdCl_2(PPh_3)_2]$   | Et₃N                           | 1:1                           | 58   | 40         | 0                 |
| 23    | $[PdCl_2(PPh_3)_2]$   | Et₃N                           | 3:1                           | 75   | 24         | 0                 |
| 24    | $[PdCl_2(PPh_3)_2]$   | Et₃N                           | 5:1                           | 74   | 24         | 0                 |

[a] Reaction conditions: Using **1a** (0.25 mmol), base (0.5 mmol), Pd catalyst (0.0125 mmol), and air/CO in 1 atm pressure. [b] Yield determined by GC analysis. bpy=2,2'-bipyridine , dabco=1,4-diazabicyclo-[2.2.2]octane, dbu=1,8-diazabicyclo[5.4.0]undec-7-ene, dppf=diphenylphosphinoferrocene, py=pyridine.



entries 20–24). Therefore, the optimized conditions for the oxidative carbonylation were 5 mol% of  $[PdCl_2(PPh_3)_2]$  as the catalyst, triethylamine as the base, and a ratio of air/CO 3:1 at 40 °C.

In Table 2, different arylboronic acid derivatives were compared under the optimized reaction conditions. The phenylboronic anhydride (63% yield of **2a**; Table 2, entry 1) performed slightly worse than **1a** (75% yield of **2a**; Table 1, entry 23). The oxidative carbonylation of phenylboronic acid produced **2a** in 55% yield (Table 2, entry 2). When potassium phenyltrifluoroborate was employed, the yields of both **2a** and **3a** were poor.

A further investigation of the substrate scope was carried out under the optimized reaction conditions. In Table 3, various ArBneop substrates were tested. Substrates containing electron-donating groups provided higher yields (Table 3, Table 2: Oxidative carbonylation of different arylboronic acid derivatives.  $^{[a]}$ 

| Entry | y 1 Yiel             |     | [%] <sup>[b]</sup> |
|-------|----------------------|-----|--------------------|
|       |                      | 2 a | 3a                 |
| 1     | (PhBO)₃              | 63  | 36                 |
| 2     | PhB(OH) <sub>2</sub> | 55  | 45                 |
| 3     | PhBF₃K               | 15  | 15                 |

[a] Reaction conditions: Using 1 (0.25 mmol),  $Et_3N$  (0.5 mmol),  $[PdCl_2-(PPh_3)_2]$  (0.0125 mmol), CO/air (1:3) at 1 atm pressure. [b] Yield determined by GC analysis.

**Table 3:** Palladium-catalyzed aerobic oxidative carbonylation of ArBneopin nBuOH.<sup>[a]</sup>

| $\Lambda_{r} = P^{O} / (PdCl_2(PPh_3)_2) (5 \text{ mol } \%), Et_3N$ |                                    |  |  |  |
|--|------------------------------------|--|--|--|
|  | CO/air (1:3), <i>n</i> BuOH (2 ml) |  |  |  |
| 1  | 40 °C, 24 h                        |  |  |  |

| Entry | 1   | Ar  | Yield [%] <sup>[b]</sup> |
|-------|-----|-----|--------------------------|
| 1     | la  |     | 75                       |
| 2     | 1 b | Me  | 81                       |
| 3     | lc  | Me  | 85                       |
| 4     | ٦d  | Me  | 80                       |
| 5     | le  | MeO | 79                       |
| 6     | 1 f | MeO | 95                       |
| 7     | lg  | ci– | 66                       |
| 8     | 1 h | F   | 61                       |
| 9     | 11  |     | 74                       |
| 10    | 1j  | Me  | 69                       |

[a] Reaction Conditions: Using 1 (0.25 mmol), Et<sub>3</sub>N (0.5 mmol), [PdCl<sub>2</sub>- (PPh<sub>3</sub>)<sub>2</sub>] (0.0125 mmol), CO/air (1:3) at 1 atm pressure. [b] Yield of isolated product.

entries 1–6). The yields for *p*-MeO phenylBneop **1f** and *p*-fluoro phenylBneop **1h** are 95% and 61%, respectively (Table 3, entries 6 and 8). Ortho-substituted substrates, such as *o*-Me phenylBneop **1b** and naphthylBneop **1i** and **1j** gave satisfactory results (Table 3, entries 2, 9, and 10).

Furthermore, a variety of alcohols were tested and the results are listed in Table 4. In the presence of isopropanol, the reactions gave good results at  $50 \,^{\circ}$ C for 24 hours in the presence of  $10 \,\text{mol}\,\%$  of palladium catalyst (Table 4,

**Table 4:** Palladium-catalyzed aerobic oxidative carbonylation of ArBneop in different alcohols.<sup>[a]</sup>

|  | 0-<br>Ar-B   |                  | 13)2] (5 mol %), Et <sub>3</sub> N  |  |  |
|--|--|------------------|---|--|--|
|  | Ar -B, CO/air (1:3), ROH (2 ml)<br>40 °C, 24 h     |                  |   |  |  |
| Entry  | 1  | Ar               | Alcohol (ROH)   | Yield [%] <sup>[b]</sup>                           |  |
| 1 <sup>[c]</sup>   | 1 d  |                  | iPrOH   | 78   |  |
| 2 <sup>[c]</sup>   | 1 f  |                  | <i>i</i> PrOH   | 75   |  |
| 3 <sup>[c]</sup>   | 1 g  |                  | iPrOH   | 70   |  |
| 4 <sup>[c]</sup>   | 1 k  |                  | iPrOH   | 85   |  |
| 5 <sup>[c]</sup>   | 11   | F <sub>3</sub> C | iPrOH   | 71   |  |
| 6  | le   |                  | EtOH  | 77   |  |
| 7  | 1i   |                  | EtOH  | 70   |  |
| 8  | la   |                  | EtOH  | 73   |  |
| 9  | 1 f  |                  | MeOH  | 90   |  |
| 1 <sup>[c]</sup><br>2 <sup>[c]</sup><br>3 <sup>[c]</sup><br>4 <sup>[c]</sup><br>5 <sup>[c]</sup><br>6<br>7<br>8<br>9 | ld<br>lf<br>lg<br>lk<br>ll<br>le<br>li<br>la<br>lf |                  | iPrOH<br>iPrOH<br>iPrOH<br>iPrOH<br>iPrOH<br>EtOH<br>EtOH<br>EtOH<br>EtOH<br>MeOH | 78<br>75<br>70<br>85<br>71<br>77<br>70<br>73<br>90 |  |

[a] Reaction conditions: Using 1 (0.25 mmol), Et<sub>3</sub>N (0.5 mmol), [PdCl<sub>2</sub>- (PPh<sub>3</sub>)<sub>2</sub>] (0.0125 mmol), and CO/air (1:3) at 1 atm pressure. [b] Yield of isolated product. [c] Used [PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>] (0.025 mmol) at 50 °C.

entries 1–5).<sup>[33]</sup> Ethanol and methanol worked well under the standard reaction conditions (Table 4, entries 6–9).

It is known that the reaction of  $[Pd(PPh_3)_4]$  with oxygen affords the  $[(\eta-O_2)Pd(PPh_3)_2]$  complex.<sup>[34]</sup> Jutand and coworkers investigated the transmetalation between the  $[(\eta-O_2)Pd(PPh_3)_2]$  complex and PhB(OH)<sub>2</sub>, and a fast reaction generated the *trans*-[PhPd(OH)(PPh\_3)<sub>2</sub>] complex.<sup>[35,36]</sup> A proposed catalytic cycle of the aerobic oxidative carbonylation was shown in Scheme 2. Carbonylation of *trans*-



Scheme 2. Proposed mechanism.

 $[ArPd(OH)(PPh_3)_2]$  in the presence of CO, ROH, and base would result in the formation of product ArCOOR, and a Pd<sup>0</sup>L<sub>n</sub> species. The  $[(\eta$ -O<sub>2</sub>)Pd(PPh<sub>3</sub>)<sub>2</sub>] complex would be regenerated from the reaction of  $[Pd(PPh_3)_n]$  with air.

In Scheme 3, a simplified model was used to compare the energy difference between *cis*- and *trans*-[ArPd(OH)(L)<sub>2</sub>]. The *trans* complex was 4.8 kcalmol<sup>-1</sup> more stable than the *cis* complex. Thus, we speculated that in this oxidative carbonylation, *cis*-[ArPd(OH)(PPh<sub>3</sub>)<sub>2</sub>], initially produced from the transmetalation of  $[(\eta$ -O<sub>2</sub>)Pd(PPh<sub>3</sub>)<sub>2</sub>] with PhB(OH)<sub>2</sub>, would quickly transform into *trans*-[ArPd(OH)-(PPh<sub>3</sub>)<sub>2</sub>].

In the presence of CO gas, two experiments revealed the different transmetalation reactivities of  $[PdCl_2(PPh_3)_2]$  and  $[(\eta-O_2)Pd(PPh_3)_2]$  complex with **1a**. No biphenyl or any

**Scheme 3.** The Gibbs free energy difference between *cis*- and *trans*- $[ArPd(OH)(L)_2]$ .

carbonylation products were identified in the reaction of 1a with a stoichiometric amount of  $[PdCl_2(PPh_3)_2]$  when  $NEt_3$  was employed as the base [Eq. (1)]. In fact, starting material

$$Ph-B'_{O} \xrightarrow{V} 2Et_{3}N + [PdCl_{2}(PPh_{3})_{2}] \xrightarrow{CO, nBuOH,} no reaction$$
(1)

**1a** was quantitatively recovered. However, when  $[PdCl_2-(PPh_3)_2]$  was replaced by a stoichiometric amount of  $[(\eta-O_2)Pd(PPh_3)_2]$ , *n*-butyl benzoate (**2a**) and phenol (**3a**) were obtained in 50% yield, respectively [Eq. (2)]. Similarly, in the

$$\begin{array}{c} \mathsf{Ph}-\mathsf{B}_{O}^{\prime} \xrightarrow{O} + 2\mathsf{Et}_{3}\mathsf{N} + \bigcup_{O}^{\mathsf{O}}\mathsf{Pd}(\mathsf{PPh}_{3})_{2} \xrightarrow{\mathsf{CO}, n\mathsf{BuOH},}{40 \ ^{\circ}\mathsf{C}, 18\mathsf{h}} \xrightarrow{\mathsf{PhCO}_{2}n\mathsf{Bu}^{+} \mathsf{PhOH}} \mathsf{PhOH} (2) \\ \mathbf{1a} \end{array}$$

absence of CO and air at 40 °C using NEt<sub>3</sub> as the base and *n*BuOH as the solvent, **1a** did not react with a stoichiometric amount of  $[PdCl_2(PPh_3)_2]$  either [Eq. (3)]. However, in air,

$$Ph-B \stackrel{O}{\underset{O}{\longrightarrow}} + 2Et_{3}N + [PdCl_{2}(PPh_{3})_{2}] \xrightarrow{nBuOH,} \text{ no reaction}$$
(3)  
1a

the homocoupled product biphenyl (4a) and oxidative protodeboronation product phenol (3a) were generated in 56% and 44% yield, respectively [Eq. (4)]. These comparative

$$Ph-B' \xrightarrow{O} + 2Et_{3}N \xrightarrow{[PdCl_{2}(PPh_{3})_{2}](5 \text{ mol }\%)}{air, nBuOH, 55 °C, 5h} \xrightarrow{PhOH + PhPh} (4)$$
1a

results indicated that the transmetalation reaction could only occur between phenylBneop and  $[(\eta-O_2)Pd(PPh_3)_2]$  under these reaction conditions;  $[PdCl_2(PPh_3)_2]$  as the catalyst precursor needs to convert into  $[(\eta-O_2)Pd(PPh_3)_2]$  before initiating the catalytic cycle.

To study whether peroxide species were formed in this reaction, aliquots were tested using potassium iodide starch. Interesting, the test paper did not turn blue in the early stage of the reaction (the solution of the reaction was yellow). When the yellow reaction solution turned red, the potassium iodide starch test paper turned blue, thus proving that the peroxides were formed during the reaction.

## Communications

Another interesting finding is that there was almost no biaryl product formed in this oxidative carbonylation. However, arylboronic acid derivatives could be homocoupled in the presence of oxygen.<sup>[37–39]</sup> The difference between the oxidative homocoupling and this oxidative carbonylation was mainly attributed to the presence or absence of CO. Thus, we presumed that it was CO which led the high selectivities between the oxidative carbonylation and oxidative homocoupling, but we do not have enough data to confirm this at present.

In conclusion, we have developed a novel oxidative carbonylation reaction. Under a balloon pressure of CO/air and at temperatures of  $40 \rightarrow 50$  °C, readily available arylboronic acid derivatives could be converted into their corresponding carbonylative products using [PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>] as the catalyst precursor. It is the first example whereby arylboronic acid derivatives can be transformed into carbonylative products using air as the sole oxidant under mild conditions.<sup>[34]</sup>

## **Experimental Section**

General procedure:  $[PdCl_2(PPh_3)_2]$  (0.0125 mmol) and arylboronate ester (0.25 mmol) was added to a Schlenk tube equipped with a stirrer bar. A balloon filled with air and CO (the ratio is 3:1) was connected to the Schlenk tube through the side arm. The mixture was purged with the mixed gas for 1 min before triethylamine (0.5 mmol) and 1butanol (2 mL) were added. The Schlenk tube was placed in an oil bath and heated to 40 °C for 24 h, and then cooled to room temperature. After the crude mixture was concentrated under vacuum, the reaction mixture was purified by flash chromatography on silica gel (eluent: petroleum ether/ethyl acetate 60:1) to afford the product.

Received: January 26, 2010 Published online: April 1, 2010

**Keywords:** air oxidation · arylboronate esters · homogeneous catalysis · oxidative carbonylation · palladium

- a) A. Schoenberg, R. F. Heck, J. Org. Chem. 1974, 39, 3327– 3331; b) A. Brennführer, H. Neumann, M. Beller, Angew. Chem. 2009, 121, 4176–4196; Angew. Chem. Int. Ed. 2009, 48, 4114– 4133.
- [2] R. Skoda-Földes, L. Kollar, Curr. Org. Chem. 2002, 6, 1097– 1119.
- [3] M. Beller, B. Cornils, C. D. Frohning, C. W. Kohlpaintner, J. Mol. Catal. A 1995, 104, 17–85.
- [4] G. Zanti, D. Peeters, Eur. J. Inorg. Chem. 2009, 3904-3911.
- [5] C. F. J. Barnard, Org. Process Res. Dev. 2008, 12, 566-574.
- [6] C. F. J. Barnard, Organometallics 2008, 27, 5402-5422.
- [7] S. Inoue, H. Shiota, Y. Fukumoto, N. Chatani, J. Am. Chem. Soc. 2009, 131, 6898–6899.
- [8] Z. H. Guan, Z. H. Ren, S. M. Spinella, S. C. Yu, Y. M. Liang, X. M. Zhang, J. Am. Chem. Soc. 2009, 131, 729–733.
- [9] R. Giri, J. Q. Yu, J. Am. Chem. Soc. 2008, 130, 14082-14083.

- [10] K. Orito, A. Horibata, T. Nakamura, H. Ushito, H. Nagasaki, M. Yuguchi, S. Yamashita, M. Tokuda, J. Am. Chem. Soc. 2004, 126, 14342–14343.
- [11] Y. Zhao, L. Jin, P. Li, A. Lei, J. Am. Chem. Soc. 2008, 130, 9429– 9433.
- [12] A. Suzuki, J. Synth. Org. Chem. Jpn. 2005, 63, 312-324.
- [13] A. Suzuki in Organoboranes for Syntheses, Vol. 783, American Chemical Society, New York, 2001, pp. 80–93.
- [14] Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine (Ed.: D. G. Hall), Wiley-VCH, Weinheim, 2005.
- [15] R. Martin, S. L. Buchwald, Acc. Chem. Res. 2008, 41, 1461-1473.
- [16] A. Suzuki, J. Synth. Org. Chem. Jpn. 2004, 80, 359-371.
- [17] F. Bellina, A. Carpita, R. Rossi, Synthesis 2004, 2419-2440.
- [18] T. Ohishi, M. Nishiura, Z. Hou, Angew. Chem. 2008, 120, 5876– 5879; Angew. Chem. Int. Ed. 2008, 47, 5792–5795.
- [19] K. Ukai, M. Aoki, J. Takaya, N. Iwasawa, J. Am. Chem. Soc. 2006, 128, 8706–8707.
- [20] H. Neumann, A. Brennfuehrer, M. Beller, *Chem. Eur. J.* 2008, 14, 3645–3652, and references therein.
- [21] S. S. Stahl, Angew. Chem. 2004, 116, 3480-3501; Angew. Chem. Int. Ed. 2004, 43, 3400-3420.
- [22] J. L. Brice, J. E. Harang, V. I. Timokhin, N. R. Anastasi, S. S. Stahl, J. Am. Chem. Soc. 2005, 127, 2868–2869.
- [23] M. M. Rogers, J. E. Wendlandt, I. A. Guzei, S. S. Stahl, Org. Lett. 2006, 8, 2257–2260.
- [24] C. C. Scarborough, S. S. Stahl, Org. Lett. 2006, 8, 3251-3254.
- [25] S. S. Stahl, Science 2005, 309, 1824-1826.
- [26] V. Kotov, C. C. Scarborough, S. S. Stahl, *Inorg. Chem.* 2007, 46, 1910–1923.
- [27] B. Gabriele, R. Mancuso, G. Salerno, E. Lupinacci, G. Ruffolo, M. Costa, J. Org. Chem. 2008, 73, 4971–4977.
- [28] B. Gabriele, P. Plastina, G. Salerno, R. Mancuso, M. Costa, Org. Lett. 2007, 9, 3319–3322.
- [29] B. Gabriele, G. Salerno, L. Veltri, R. Mancuso, Z. Y. Li, A. Crispini, A. Bellusci, J. Org. Chem. 2006, 71, 7895-7898.
- [30] B. Gabriele, G. Salerno, M. Costa, *Top. Organomet. Chem.* 2006, 18, 239–272.
- [31] K. Orito, M. Miyazawa, T. Nakamura, A. Horibata, H. Ushito, H. Nagasaki, M. Yuguchi, S. Yamashita, T. Yamazaki, M. Tokuda, J. Org. Chem. 2006, 71, 5951–5958.
- [32] J. Dupont, C. S. Consorti, J. Spencer, Chem. Rev. 2005, 105, 2527–2571.
- [33] At 40°C with 5 mol% of palladium catalyst, the reactions need 48 hours to reach completion. When the reaction temperature was increased to 60°C, low conversion and palladium-black formation were observed.
- [34] Y. Izawa, I. Shimizu, A. Yamamoto, Bull. Chem. Soc. Jpn. 2004, 77, 2033–2045.
- [35] C. Adamo, C. Amatore, I. Ciofini, A. Jutand, H. Lakmini, J. Am. Chem. Soc. 2006, 128, 6829–6836.
- [36] This transmetalation step did not involve the ligand dissociation, see Ref. [35].
- [37] J. S. Chen, K. Krogh-Jespersen, J. G. Khinast, J. Mol. Catal. A 2008, 285, 14–19.
- [38] H. Yoshida, Y. Yamaryo, J. Ohshita, A. Kunai, *Tetrahedron Lett.* 2003, 44, 1541–1544.
- [39] K. A. Smith, E. M. Campi, W. R. Jackson, S. Marcuccio, C. G. M. Naeslund, G. B. Deacon, *Synlett* **1997**, 131–132.