

## ORIGINAL ARTICLE

# Absolute stereochemistry of pentaceciliides, new inhibitors of lipid droplet formation in mouse macrophages, produced by *Penicillium cecidicola* FKI-3765-1

Hiroyuki Yamazaki, Narihiro Ugaki, Daisuke Matsuda and Hiroshi Tomoda

The structure of a new pentacecilide congener, pentacecilide D, produced by *Penicillium cecidicola* FKI-3765-1 was elucidated by various NMR experiments. The absolute stereochemistry of pentaceciliides was elucidated by using the modified Mosher method for pentacecilide C. The inhibitory activity of all pentaceciliides against lipid droplet formation and acyl-CoA:cholesterol acyltransferase isozymes was compared.

*The Journal of Antibiotics* (2010) 63, 315–318; doi:10.1038/ja.2010.39; published online 23 April 2010

**Keywords:** absolute stereochemistry; acyl-CoA:cholesterol acyltransferase; fungal metabolite; lipid droplet formation; *Penicillium cecidicola*; pentaceciliides D

## INTRODUCTION

As we reported previously, pentaceciliides A to C (Figure 1) were isolated from the whole culture of *Penicillium cecidicola* FKI-3765-1 as inhibitors of lipid droplet formation in mouse macrophages.<sup>1,2</sup> From the biochemical study,<sup>1</sup> pentaceciliides inhibited lipid droplet formation in mouse macrophages, probably due to the blockade of acyl-CoA:cholesterol acyltransferase (ACAT) activity. Furthermore, pentaceciliides appeared as dual inhibitors of ACAT1 and ACAT2 isozymes.<sup>1,3–7</sup> The structures including the relative stereochemistry were also elucidated;<sup>2</sup> however, the indication of the relative stereochemistry at C-2 was revised<sup>8,9</sup> because erroneous structures of pentaceciliides had been shown in our previous reports.<sup>1,2</sup> Further precise analysis of metabolites in the whole culture led to the discovery of a new congener named pentacecilide D (Figure 1). In this study, the structure of the new pentacecilide congener was elucidated. Furthermore, the absolute stereochemistry of pentaceciliides was determined by using the modified Mosher method<sup>10</sup> for pentacecilide C.

## RESULTS

### Isolation of pentacecilide D

The 13-day-old whole culture (1000 g) was extracted with 2.0 l of acetone. After the acetone extracts were filtered and concentrated to remove acetone, the aqueous solution was extracted with ethyl acetate. The extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo* to dryness to yield a red-brown material (1.7 g). The material was dissolved in 30% CH<sub>3</sub>CN, applied to an octadecyl silyl (ODS) column chromatography (100 g), and eluted stepwise with 30, 50, 70, 100%

CH<sub>3</sub>CN containing 0.050 % trifluoroacetic acid (250 ml×2 tubes for each solvent). Pentaceciliides A and B were present in the first tube of 100% CH<sub>3</sub>CN. The first tube of 70% CH<sub>3</sub>CN containing pentaceciliides C and D was concentrated *in vacuo* to dryness to give a red-brown material. The material (172 mg) was finally purified by preparative HPLC (column, PEGASIL ODS, 20×250 mm; Senshu Scientific, Tokyo, Japan; solvent, 55% CH<sub>3</sub>CN containing 0.050% trifluoroacetic acid; detection, UV at 210 nm; flow rate, 8.0 ml min<sup>-1</sup>). Under these conditions, pentaceciliides C and D were eluted as peaks with retention times of 15.8 and 25.8 min, respectively. The respective fractions were concentrated *in vacuo* to dryness to give pure pentaceciliides C (56.6 mg) and D (5.1 mg) as white crystals.

### Structural elucidation of pentacecilide D

The physicochemical properties of pentacecilide D are summarized in Table 1. Pentacecilide D showed UV absorption maxima at 219, 276 and 310 nm, similar to pentaceciliides A to C.<sup>2</sup> IR absorption at 3426 and 1712–1617 cm<sup>-1</sup> suggested the presence of hydroxy and carbonyl groups in the structure.

The molecular formula was determined to be C<sub>25</sub>H<sub>32</sub>O<sub>7</sub> on the basis of HRESI-TOF-MS measurement, indicating that pentacecilide D was smaller than pentacecilide C.<sup>2</sup> The <sup>13</sup>C NMR spectrum (in CDCl<sub>3</sub>) showed 25 resolved signals, which were classified into five methylene carbons, four methylene carbons, two *sp*<sup>3</sup> methine carbons, one *sp*<sup>2</sup> methine carbon, three oxygenated *sp*<sup>3</sup> methine carbons, two *sp*<sup>3</sup> quaternary carbons, one oxygenated *sp*<sup>3</sup> quaternary carbon, three *sp*<sup>2</sup> quaternary carbons, two oxygenated *sp*<sup>2</sup> quaternary carbons and two




Figure 1 Structures of pentacecillides A–D.

Table 1 Physicochemical properties of pentacecillide D

Pentacecillide D	
Appearance	White crystalline solid
Molecular weight	444
Molecular formula	C <sub>25</sub> H <sub>32</sub> O <sub>7</sub>
<i>HRESI-TOF-MS (m/z)</i>	
Calcd:	445.2226 (M+H) <sup>+</sup>
Found:	445.2216 (M+H) <sup>+</sup>
UV (MeOH) λ <sub>max</sub> , nm (ε)	219 (34800), 276 (24700), 310 (11800)
[α] <sub>D</sub> <sup>26</sup>	−51.4° (c 0.10, CHCl <sub>3</sub> )
IR (KBr) ν <sub>max</sub> (cm <sup>−1</sup> )	3426, 1712, 1666, 1617, 1473

carbonyl carbons by analysis of the DEPT and HSQC spectra. The <sup>1</sup>H NMR spectrum (in CDCl<sub>3</sub>) displayed 30 proton signals, one of which was suggested to be a hydroxy proton (δ 11.09). Taking the molecular formula into consideration, the presence of two hydroxy protons was suggested. The connectivity of proton and carbon atoms was established by the <sup>13</sup>C-<sup>1</sup>H HSQC spectrum (Table 2). These spectral data from NMR and physicochemical properties showed that pentacecillide D has the same skeleton as pentacecillide C.<sup>2</sup> Comparison of the <sup>1</sup>H NMR spectra between pentacecillides C and D indicated that the methyl proton signals (C-17, δ 2.17) derived from the acetyl group in pentacecillide C disappeared from pentacecillide D. In addition, the chemical shift of C-2 (δ 69.9) and the molecular formula showed the presence of a hydroxy group at C-2. These data led to the conclusion that the structure of pentacecillide D was 2-deacetyl pentacecillide C (Figure 1). The structure satisfied the degree of unsaturation and the molecular formula.

Regarding the relative stereochemistry of pentacecillide D, NOESY experiments were carried out showing coupling constants in <sup>1</sup>H NMR similar to pentacecillide C.<sup>2,8,9</sup> Accordingly, the relative stereochemistry of pentacecillide D was elucidated as shown in Figure 1.

#### Absolute stereochemistry of pentacecillides

To elucidate the absolute stereochemistry of pentacecillides, the (*S*)- (1) and (*R*)- α-methoxy-α-(trifluoromethyl)phenylacetyl (MTPA) esters (2) were prepared by treatment of pentacecillide C with (*R*)- (−)- and (*S*)-(+)- MTPA chloride in the presence of 4-(dimethylamino) pyridine and triethylamine (in CH<sub>2</sub>Cl<sub>2</sub>, at room temperature, 80.6 and 61.5% yield, respectively) (Scheme 1). The <sup>1</sup>H NMR spectra of 1 and 2 were completely assigned. As shown in Figure 2, the

Table 2 <sup>1</sup>H and <sup>13</sup>C NMR chemical shifts of pentacecillide D


No.	Pentacecillide D	
	δ <sub>C</sub>	δ <sub>H</sub> (J in Hz)
1	44.9	1.60 m
		2.38 m
2	69.9	4.64 m
3	214.8	—
4	47.6	—
5	46.4	1.80 m
6	26.6	1.86 m
		2.18 m
7	71.8	4.13 dd (10.0, 3.0)
8	79.2	—
9	43.4	2.21 m
10	36.0	—
11	21.5	2.52 m
12	20.9	1.23 s
13	20.9	1.20 s
14	25.4	1.19 s
15	23.9	1.54 s
1'	110.9	—
2'	139.4	—
3'	103.6	—
4'	162.4	—
5'	103.6	6.34 s
6'	159.1	—
7'	31.9	2.71 dd (17.0, 11.0)
		2.84 dd (17.0, 3.5)
8'	74.7	4.64 m
9'	20.9	1.56 d (7.0)
10'	170.0	—
4'-OH		11.09 s

calculated Δδ values (Δδ = δ<sub>S</sub> − δ<sub>R</sub>) of H<sub>2</sub>-11, H<sub>3</sub>-12, H-5' and H-8' were positive, whereas those (Δδ = δ<sub>S</sub> − δ<sub>R</sub>) of H<sub>2</sub>-1, H-2, H-5, H<sub>2</sub>-6, H-9, H<sub>3</sub>-13, H<sub>3</sub>-14, H<sub>3</sub>-15 and H<sub>3</sub>-17 were negative, indicating that the absolute stereochemistry of C-7 was elucidated to be 7*S*. Accordingly, the absolute stereochemistry of the seven chiral centers in pentacecillide C was concluded to be 2*R*5*R*7*S*8*R*9*R*10*S*8'*S*, as shown in Figure 1. It is plausible that all pentacecillides are biosynthesized by the same pathway; therefore, the absolute stereochemistry of the other pentacecillides was deduced to be the same as that of pentacecillide C (Figure 1).


#### Biological properties

*Inhibition of CE synthesis in macrophages and ACAT1- and ACAT2-CHO cells.* In the previous study, pentacecillides inhibited cholesteryl ester (CE) synthesis in mouse macrophages and ACAT1- and ACAT2-CHO cells, strongly suggesting that they were dual inhibitors of ACAT1 and ACAT2 isozymes.<sup>1</sup> Pentacecillide D and semisynthetic 3*S*'-hydroxy pentacecillide A (3) were evaluated in these three cell-based assays, and the IC<sub>50</sub> values of CE synthesis were compared. Pentacecillide A is the most potent inhibitor of CE synthesis in the three assays, followed by pentacecillide B.<sup>1</sup> Unfortunately, pentacecillides D and 3 showed almost no inhibitory activity at 22.5–24.2 μM.

No pentacecillides showed any cytotoxic effects on macrophages and ACAT1- and ACAT2-CHO cells, even at 20.6–24.3 μM.



**Scheme 1** Semisynthetic preparation of 1–3.



**Figure 2** Absolute stereochemistry determination.  $\Delta\delta$  values ( $\Delta\delta$  (in p.p.m.) =  $\delta_S - \delta_R$ ) obtained for di-Mosher esters 1 and 2.

## DISCUSSION

As described above, from various spectral analyses, the structure of a new congener, pentacecillide D, was elucidated to be 2-deacetyl pentacecillide C with the same relative stereochemistry as shown in Figure 1.

To determine the absolute stereochemistry of pentacecillides, the modified Mosher method<sup>10</sup> was adapted to pentacecillide C. Calculation of the  $\Delta\delta$  values led to the clear conclusion that pentacecillide C has the absolute stereochemistry of 2*R*5*R*7*S*8*R*9*R*10*S*8'*S* (Figure 1). Furthermore, assuming that all pentacecillides are generated through the same biosynthetic pathway, pentacecillides A, B and D have the same absolute stereochemistry as pentacecillide C. This

**Table 3** Effects of pentacecillides on cholesteryl ester synthesis in macrophages, and ACAT1- and ACAT2-CHO cells

Pentacecillide	<i>IC</i> <sub>50</sub> ( $\mu$ M)		
	Macrophages	ACAT1-CHO	ACAT2-CHO
A	3.65	1.09	0.69
B	4.76	10.8	3.97
C	> 20.6	> 20.6	> 20.6
D	> 22.5	> 22.5	> 22.5
3	> 24.2	> 24.2	19.0

conclusion was supported by evidence that all pentacecillides have negative optical rotations.<sup>2</sup>

In a cell-based assay using mouse macrophages, ACAT1- and ACAT2-CHO cells,<sup>11–13</sup> pentacecillides A and B showed more potent inhibition of CE synthesis than pentacecillides C and D<sup>1</sup> (Table 3). Furthermore, 3 showed no inhibition of CE synthesis in mouse macrophages, and ACAT1- and ACAT2-CHO cells, even at 24.2  $\mu$ M (Table 3). Taking the structural differences into consideration (Figure 1), the presence of a hydroxy group at C-7 is unfavorable for inhibiting lipid droplet formation and ACAT activity. Regarding the C-3 position, 3 with a hydroxy group at C-3 showed weaker inhibitory activity than pentacecillide A, indicating that the presence of a carbonyl group at C-3 is important for activity. Regarding the C-2 position, introduction of acethoxy and hydroxy groups resulted in decreased activity. Namely, no modification at this position is preferable for inhibitory activity. Among these pentacecillides, it was concluded that pentacecillide A has the most favorable structure for showing potent inhibitory activity against lipid droplet formation and ACAT isozymes. Regarding the structure–activity relationships of

the hydroxy group at C-4', synthesis of acyl derivatives at this position is now in progress.

## METHODS

### Materials

(S)-(+)- and (R)-(-)-MTPA chlorides, and sodium borohydride (NaBH<sub>4</sub>) were purchased from Wako Pure Chemical Industries (Osaka, Japan). Acetic anhydride (Ac<sub>2</sub>O) and 4-(dimethylamino) pyridine were purchased from Kanto Chemical (Tokyo, Japan). Triethylamine was purchased from Nacalai Tesque (Kyoto, Japan).

### General experimental procedures

Fungal strain *P. cecidicola* FKI-3765-1 was used for the production of pentacecillides.<sup>1,2</sup> Pentacecillide D was from 13-day-old culture broth fermented as reported previously.<sup>1</sup> For determination of the amounts of pentacecillide D in culture broths, samples (ethyl acetate extracts) dissolved in methanol were analyzed by the HP1100 system (Hewlett-Packard, Palo Alto, CA, USA) under the following conditions: column, Symmetry (2.1×150 mm; Waters, Milford, MA, USA); flow rate, 0.2 ml min<sup>-1</sup>; mobile phase, a 20-min linear gradient from 60% CH<sub>3</sub>CN to 100% CH<sub>3</sub>CN containing 0.050% H<sub>3</sub>PO<sub>4</sub>; detection, UV at 210 nm. Under these conditions, pentacecillide D was eluted with a retention time of 3.59 min, respectively.

SSC-ODS-7515-12 (Senshu Scientific) was used for ODS column chromatography. HPLC was carried out using the L-6200 system (Hitachi, Tokyo, Japan).

UV spectra were recorded on a spectrophotometer (8453 UV-Visible spectrophotometer; Agilent Technologies, Santa Clara, CA, USA). IR spectra were recorded on a Fourier transform infrared spectrometer (FT-710; Horiba, Kyoto, Japan). Optical rotations were measured with a digital polarimeter (DIP-1000; JASCO, Tokyo, Japan). ESI-TOF-MS and HRESI-TOF-MS spectra were recorded on a mass spectrometer (JMS-T100LP; JEOL, Tokyo, Japan). Various NMR spectra were measured with a spectrometer (XL-400; Varian, Palo Alto, CA, USA).

### Preparation of the (S)-(+)- and (R)-(-)-MTPA ester derivatives of pentacecillide C

To a solution of pentacecillide C (5.0 mg, 0.010 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (500 μl), (R)-(-)-MTPA chloride (16.5 mg, 0.065 mmol), 4-(dimethylamino) pyridine (5.3 mg, 0.043 mmol) and triethylamine (10 μl, 0.073 mmol) were added. The reaction mixture was stirred at room temperature. After 10 h, the reaction mixture was diluted with 1.0 N HCl and the aqueous phase was extracted with EtOAc. The organic layer was recovered, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo* to dryness. The product was purified by preparative HPLC (column; PEGASIL ODS, 20×250 mm; Senshu Scientific; solvent, 80% CH<sub>3</sub>CN; detection, UV at 210 nm; flow rate, 8.0 ml min<sup>-1</sup>) to give **1** (7.40 mg, 0.0081 mmol, 80.6%) as a white powder. Similarly, **2** (4.63 mg, 0.0050 mmol, 61.5%) was obtained using (S)-(+)-MTPA chloride.

**7**, 4'-(S)-(+)-α-methoxy-α-(trifluoromethyl)phenylacetoxypentacecillide C (**1**): <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.05 (s, 3H), 1.06 (s, 3H), 1.10 (s, 3H), 1.36 (s, 3H), 1.52 (d, 3H, *J*=7.0 Hz), 1.68, 2.28 (m, 2H), 1.85, 2.14 (m, 2H), 2.17 (s, 3H), 2.57 (m, 2H), 2.74, 2.82 (dd, 2H, *J*=17.0, 11.0 Hz, *J*=17.0, 3.5 Hz), 3.52 (s, 3H), 3.73 (s, 3H), 4.58 (m, 1H), 5.54 (m, 1H), 5.56 (m, 1H), 6.35 (s, 1H), 7.26 (m, 3H), 7.44 (m, 3H), 7.52 (m, 2H), 7.75 (m, 2H); HRESI-TOF-MS (*m/z*) found: 941.2935, calcd: 941.2948 [M+Na]<sup>+</sup> for C<sub>47</sub>H<sub>48</sub>F<sub>8</sub>NaO<sub>12</sub>.

**7**, 4'-(R)-(-)-α-methoxy-α-(trifluoromethyl)phenylacetoxypentacecillide C (**2**): <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.14 (s, 3H), 1.17 (s, 3H), 1.30 (s, 3H), 1.33 (s, 3H), 1.53 (d, 3H, *J*=7.0 Hz), 1.87, 2.16 (m, 2H), 2.31 (m, 2H), 2.17 (s, 3H), 2.50 (m, 2H), 2.74, 2.82 (dd, 2H, *J*=17.0, 11.0 Hz, *J*=17.0, 3.5 Hz), 3.53 (s, 3H), 3.76 (s, 3H), 4.57 (m, 1H), 5.60 (m, 1H), 5.61 (m, 1H), 6.31 (s, 1H), 7.32 (m, 3H), 7.44 (m, 3H), 7.52 (m, 2H), 7.74 (m, 2H); HRESI-TOF-MS (*m/z*) found: 941.2980, calcd: 941.2948 [M+Na]<sup>+</sup> for C<sub>47</sub>H<sub>48</sub>F<sub>8</sub>NaO<sub>12</sub>.

### Preparation of 3S<sup>+</sup>-Hydroxy-pentacecillide A

To a solution of pentacecillide A (2.9 mg, 0.0070 mmol) in CH<sub>3</sub>OH (500 μl), NaBH<sub>4</sub> (3.3 mg, 0.087 mmol) was added, and the resulting solution was stirred at room temperature. After 10 h, the reaction mixture was diluted with H<sub>2</sub>O and the aqueous phase was extracted with EtOAc. The organic layer was recovered, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo* to dryness. The product was purified by preparative HPLC (column; PEGASIL ODS, 20×250 mm; Senshu Scientific; solvent, 80% CH<sub>3</sub>CN; detection, UV at 210 nm; flow rate, 8.0 ml min<sup>-1</sup>) to give **3** (2.5 mg, 0.0060 mmol, 85.7%) as a white powder.

3S<sup>+</sup>-Hydroxy-pentacecillide A (**3**): <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.86 (s, 3H), 1.02 (s, 3H), 1.09 (s, 3H), 1.26 (m, 1H), 1.29 (s, 3H), 1.46, 1.56 (m, 2H), 1.55 (d, 3H, *J*=6.0 Hz), 1.66, 1.80 (m, 2H), 1.70 (m, 2H), 1.76 (m, 1H), 2.00, 2.08 (m, 2H), 2.40 (m, 2H), 2.68, 2.85 (dd, 2H, *J*=17.0, 11.0 Hz, *J*=17.0, 3.5 Hz), 3.26 (dd, *J*=10.0, 4.0 Hz, 1H), 4.63 (m, 1H), 6.26 (s, 1H), 11.0 (s, 1H); HRESI-TOF-MS (*m/z*) found: 437.2290, calcd: 437.2304 [M+Na]<sup>+</sup> for C<sub>25</sub>H<sub>34</sub>NaO<sub>5</sub>.

### Biological assays

An assay for the synthesis of CE, triacylglycerol and phospholipids by mouse macrophages was carried out according to the method described previously.<sup>11</sup> An assay for ACAT1 and ACAT2 activities in ACAT1- and ACAT2-CHO cells was carried out by our established method.<sup>12,13</sup>

### ACKNOWLEDGEMENTS

This study was supported in part by a Sasakawa Scientific Research Grant (to HY) from The Japan Science Society. We express our thanks to Ms N Sato for NMR experiments, and Dr K Nagai and Ms A Nakagawa for measuring mass spectra.

- 1 Yamazaki, H. *et al.* Pentacecillides, new inhibitors for lipid droplet formation in mouse macrophages produced by *Penicillium cecidicola* FKI-3765-1:1. Taxonomy, fermentation, isolation and biological properties. *J. Antibiot.* **62**, 195–200 (2009).
- 2 Yamazaki, H., Omura, S. & Tomoda, H. Pentacecillides, new inhibitors for lipid droplet formation in mouse macrophages produced by *Penicillium cecidicola* FKI-3765-1:1. Structure elucidation. *J. Antibiot.* **62**, 207–211 (2009).
- 3 Chang, C. C., Huh, H. Y., Cadigan, K. M. & Chang, T. Y. Molecular cloning and functional expression of human acyl-coenzyme A:cholesterol acyltransferase cDNA in mutant Chinese hamster ovary cells. *J. Biol. Chem.* **268**, 20747–20755 (1993).
- 4 Anderson, R. A. *et al.* Identification of a form of acyl-CoA:cholesterol acyltransferase specific to liver and intestine in nonhuman primates. *J. Biol. Chem.* **273**, 26747–26754 (1998).
- 5 Cases, S. *et al.* ACAT-2, a second mammalian acyl-CoA:cholesterol acyltransferase. Its cloning, expression, and characterization. *J. Biol. Chem.* **273**, 26755–26764 (1998).
- 6 Oelkers, P., Behari, A., Cromley, D., Billheimer, J. T. & Sturley, S. L. Characterization of two human genes encoding acyl coenzyme A:cholesterol acyltransferase-related enzymes. *J. Biol. Chem.* **273**, 26765–26771 (1998).
- 7 Parini, P. *et al.* ACAT2 is localized to hepatocytes and is the major cholesterol-esterifying enzyme in human liver. *Circulation* **110**, 2017–2023 (2004).
- 8 Corrigenda. *J. Antibiot.* **62**, 651 (2009).
- 9 Corrigenda. *J. Antibiot.* **62**, 653–654 (2009).
- 10 Ohtani, I., Kusumi, T., Kashman, Y. & Kakisawa, H. High-field FT NMR application of Mosher's method. The absolute configuration of marine terpenoids. *J. Am. Chem. Soc.* **113**, 4092–4096 (1991).
- 11 Namatame, I., Tomoda, H., Arai, H., Inoue, K. & Omura, S. Complete inhibition of mouse macrophage-derived foam cell formation by triacsin C. *J. Biochem.* **125**, 319–327 (1999).
- 12 Lada, A. T. *et al.* Identification of ACAT1- and ACAT2-specific inhibitors using a novel, cell-based fluorescence assay: individual ACAT uniqueness. *J. Lipid. Res.* **45**, 378–386 (2004).
- 13 Ohshiro, T., Rudel, L. L., Omura, S. & Tomoda, H. Selectivity of microbial acyl-CoA:cholesterol acyltransferase inhibitors toward isozymes. *J. Antibiot.* **60**, 43–51 (2007).