

A Practical Ruthenium-Catalyzed Cleavage of the Allyl Protecting Group in Amides, Lactams, Imides, and Congeners

Benito Alcaide,*^[a] Pedro Almendros,*^[b] and Jose M. Alonso^[a]

Abstract: A convenient methodology for the deprotection of *N*-allylic amide-like moieties was developed. The first examples accounting for the ruthenium-catalyzed deallylation of amides, lactams, imides, pyrazolidones, hydantoins, and oxazolidinones have been achieved by the sequential use of Grubbs carbene (isomerization step) and RuCl₃ (oxidation step). A variety of substrates, including enantio-pure multifunctional β - and γ -lactams, can be employed.

Keywords: amides • cleavage reactions • lactams • protecting groups • ruthenium

Introduction

NH-Amides, imides, lactams, and related nitrogen-containing compounds have long attracted synthetic interest due to their relatively simple structural features and wide range of pharmacologic activity; these compounds also serve as crucial intermediates for numerous natural products. In particular, *N*-unsubstituted β -lactams are essential building blocks for the preparation of several antibiotics^[1] and anticancer agents, such as paclitaxel (Taxol) and docetaxel.^[2] Very often, the required free amide, imide, or lactam cannot be prepared by the established methods. However, there are relatively few methods available for the deprotection of an amide or amide-like moiety.^[3] Oxidative cleavage by ceric

ammonium nitrate of an activated aromatic moiety attached to the nitrogen offers the most direct synthesis of *NH*-amides, imides, or lactams.^[4] However, in many cases the yields are poor as this method requires quite harsh conditions that may damage sensitive functionalities. Therefore, due to the increasing complexity of the molecules synthesized, new protecting groups with modulated reactivity and new cleavage techniques for existing protecting groups are still needed.

The allyl protecting group, which is stable under both acidic and basic conditions, permits orthogonal protection strategies with a wide range of protecting groups, allowing its use in multistep synthetic schemes. Despite the fact that allyl protecting group cleavage in ethers and amines is a well-documented methodology, especially by using palladium chemistry,^[5] its extension to the deprotection of allylic imides and lactams has been scarcely documented. To the best of our knowledge, the *N*-allyl cleavage of a 4-unsubstituted- β -lactam^[6] and a 2-hydroxy- γ -lactam,^[7] both induced by rhodium(III) followed by KMnO₄ or acidic treatments, respectively, are the only available examples. In our ongoing project directed toward the asymmetric synthesis of nitrogen-containing products of biological interest,^[8] we faced serious difficulties with the elimination of nitrogen protecting groups on *N*-protected lactams. For this reason, and taking into account the highly selective properties of various transition metal derived reagents that would seem to recommend their application to the removal of the allyl protecting group in amides, imides, and lactams, we were tempted to study alternative strategies for the cleavage of the N–C bond in amides and congeners. We report herein full details of a novel, mild, and efficient ruthenium-catalyzed methodology for the catalytic deprotection of *N*-allyl lactams,^[9] as well as

[a] Prof. Dr. B. Alcaide, Dr. J. M. Alonso
Departamento de Química Orgánica I
Facultad de Química
Universidad Complutense de Madrid
28040 Madrid (Spain)
Fax: (+34) 91-394-4103
E-mail: alcaideb@quim.ucm.es


[b] Dr. P. Almendros
Instituto de Química Orgánica General
Consejo Superior de Investigaciones Científicas
CSIC, Juan de la Cierva 3
28006 Madrid (Spain)
Fax: (+34) 91-564-4853
E-mail: iqoa392@iqog.csic.es

Supporting information for this article is available on the WWW under <http://www.chemurj.org/> or from the author. It contains compound characterization data and experimental procedures for compounds **1a–d**, **2a–c**, **3a–d**, **4a–e**, **5a–c**, (+)-**6b**, (±)-**6d**, (+)-**6e**, (+)-**7**, (+)-**8**, (+)-**9**, (–)-**10**, (+)-**11**, **13a**, **13b**, **14b**, **15a**, **15b**, **16a–e**, **17a**, and **17c–e**.

its extension to the allyl cleavage of amides, imides, pyrazolidones, hydantoin, and oxazolidinones.

Results and Discussion

In recent years there have been several reports of nonmetallic transformations promoted by Grubbs carbene.^[10] Our exploration of this type of ruthenium-promoted reaction began after noting for the first time that allylic amines can be catalytically cleaved by using reagents other than palladium catalysts.^[11] The higher stability of enamide-like moieties compared with enamines favors the double bond isomerization,^[12] which prevents (CO)N-allyl cleavage. On the basis of these principles, it was to be expected that successful catalytic C–N deprotection in *N*-allyl amides, imides, lactams, and congeners would require an additional step (Scheme 1). We chose ruthenium(III) chloride, which has been probed as an excellent catalyst for the oxidative cleavage of olefins to aldehydes,^[13] because it has been reported that *N*-formyl lactams smoothly lose CO under slightly basic conditions to give the corresponding NH-amides.^[14]



Scheme 1. Retrosynthetic analysis for the cleavage of the allyl protecting group in lactams, imides, and analogous compounds.

First, as a model reaction we tested the deprotection of *N*-allyl γ -butyrolactam **1a**. Treatment of allylic lactam **1a** with first generation Grubbs carbene [(PCy₃)₂Cl₂Ru=CHPh] under optimized reaction conditions (5 mol% catalyst, 0.03 M, toluene, 110 °C) resulted in the clean formation of the corresponding enamide **2a** as an isomeric mixture (*E/Z* 1.2:1) in good isolated yield (90%) after chromatographic purification. Attempts to effect the reaction at lower temperatures (50 °C) slowed the reaction considerably. Incom-

plete conversion was observed on decreasing the amount of catalyst. The catalytic scission of the internal C=C was attained by using the system RuCl₃–NaIO₄ in 1,2-dichloroethane–H₂O (1:1), followed by an aqueous workup under slightly basic conditions (sat. aq NaHCO₃ containing a catalytic amount of Na₂CO₃). In this way, the *N*-unsubstituted 2-pyrrolidone **3a** was obtained in an 87% yield. Exposure of *N*-allyl L-pyrroglutamic acid ethyl ester (+)-**1b**, *N*-allyl δ -valerolactam **1b**, and *N*-allyl ϵ -caprolactam **1c** to the above sequential catalytic conditions smoothly afforded the *NH*-lactams (+)-**3b**, **3c**, and **3d** (Table 1). Under the reaction con-

Table 1. Ruthenium-catalyzed deprotection of allylic γ -, δ -, and ϵ -lactams.^[a]

Substrate	<i>n</i>	R	Product	<i>E/Z</i> ratio ^[b]	Yield [%] ^[c]	Product	Yield [%] ^[c]
1a	1	H	2a	55:45	90	3a	87
(+)- 1b	1	COOEt	(+)- 2b	20:80	75	(+)- 3b	73
1c	2	H	2c	53:47	71	3c	70
1d	3	H	2d	0:100	72	3d	65

[a] Reagents and conditions: a) 5 mol% [(PCy₃)₂Cl₂Ru=CHPh], toluene, 110 °C; b) RuCl₃–NaIO₄, 1,2-dichloroethane–H₂O (1:1), aqueous workup (sat. aq NaHCO₃ + cat. Na₂CO₃). [b] The ratio was determined by the integration of well-resolved signals in the ¹H NMR spectra (300 MHz) of the crude reaction mixtures before purification. [c] Yield of pure, isolated product with correct analytical and spectral data.

ditions, the intermediate *N*-formyl lactams accumulate in the reaction mixture immediately after the RuCl₃–NaIO₄ system is added over the enamide, and can be isolated by using a neutral workup.

Next, we decided to test the *N*-allyl cleavage protocol in the strained four-membered lactam series. Racemic and enantiopure allylic β -lactams **4a–e** were conveniently deprotected to give the corresponding *NH*- β -lactams by using both ruthenium catalysts (Table 2). This transformation tolerates different substituents on the lactam ring, such as aryl, heteroaryl, alkoxy, silyloxy, dioxolanyl, and carboxyalkyl moieties. Of special interest are the furan and electron-rich arene moieties, both of which are sensitive to oxidative deprotection conditions, as well as the acid labile silyl ether and acetonide groups. Importantly, the stereochemical integrity of the stereogenic centres on the lactam rings, when applicable, remained unaltered during the transformation of *N*-allyl compounds **4** into *NH*-products **6**.


To demonstrate that the above ruthenium-catalyzed *N*-deallylation protocol can be considered in complex synthetic planning, it was successfully used for the cleavage of highly functionalized *N*-allyl bis- β - and bis- γ -lactams.^[15] The precursors were obtained in an optically pure form from 4-oxoazetidine-2-carbaldehyde (+)-**7**.^[16] Thus, C₄,C_{4'}-bis- β -lactam (+)-**8** was prepared by the treatment of aldehyde (+)-**7** with

Table 2. Ruthenium-catalyzed deprotection of allylic β -lactams^[a]

Substrate	R ¹	R ²	Product	<i>E/Z</i> ratio ^[b]	Yield [%] ^[c]	Product	Yield [%] ^[c]
($\pm$)- 4a	Me		($\pm$)- 5a	76:24	62	($\pm$)- 6a	64
(+)- 4b	Me		(+)- 5b	80:20	77	(+)- 6b	85
(+)- 4c	TBS		(+)- 5c	88:12	79	(+)- 6c	95
($\pm$)- 4d	Ph		($\pm$)- 5d	95:5	72	($\pm$)- 6d	57
(+)- 4e	Ph		(+)- 5e	50:50	79	(+)- 6e	78

[a] Reagents and conditions: a) 5 mol % [(PCy₃)₂Cl₂Ru=CHPh], toluene, 110 °C. b) RuCl₃-NaIO₄, 1,2-dichloroethane-H₂O (1:1), aqueous workup (sat. aq. NaHCO₃ + cat. Na₂CO₃). [b] The ratio was determined by integration of well-resolved signals in the ¹H NMR spectra (300 MHz) of the crude reaction mixtures before purification. [c] Yield of pure, isolated product with correct analytical and spectral data.

allylamine, followed by ketene-imine cyclization. Bis- β -lactam (+)-**8** was conveniently deallylated by ruthenium catalysis to afford the *NH*-bis- β -lactam (+)-**9** (Scheme 2).



Scheme 2. Ruthenium-catalyzed deprotection of *N*-allyl bis- β -lactams. Reagents and conditions: a) Allylamine, MgSO₄, CH₂Cl₂, RT; b) MeOCH₂COCl, Et₃N, CH₂Cl₂, RT; c) 5 mol % [(PCy₃)₂Cl₂Ru=CHPh], toluene, 110 °C; d) RuCl₃-NaIO₄, 1,2-dichloroethane-H₂O (1:1), aqueous workup (sat. aq. NaHCO₃ + cat. Na₂CO₃). PMP=4-MeOC₆H₄.


N-Allyl bis- γ -lactam (+)-**11** was obtained from compound (+)-**8** by a two-step route through sequential sodium methoxide rearrangement to give the homopyroglutamic derivative (–)-**10** followed by acid-promoted cyclization. The *NH*-bis- γ -lactam (+)-**12** was obtained in a 64% yield by using ruthenium-catalyzed allyl breakage (Scheme 3). Worthy of note are the examples in Schemes 2 and 3, which show the selective N–C bond breakage of the *N*-allyl moiety in the presence of the *N*-*p*-methoxyphenyl functionality, allowing differentiation between two chemically equivalent functional groups within the same molecule.

Next we moved to the alicyclic series, testing the *N*-allyl cleavage protocol in both aromatic and aliphatic amides **13**.

As expected, the corresponding free amides **15** could be smoothly obtained via the corresponding enamides **14** through ruthenium catalysis (Scheme 4).


The extension of the above methodology to related systems bearing extra heteroatoms was studied next. Fortunately the ruthenium-catalyzed allyl cleavage worked nicely for imides, pyrazolidones, hydantoin, and oxazolidinones **16a–d** (Scheme 5). Although the 3-allyl-2-thioxo-4-thiazolidinone **16e** tolerates the isomerization reaction conditions, it was necessary to increase the catalyst loading from 5 to 15 mol %.^[17] However, the 2-thioxo-4-thiazolidinone moiety was not compatible with the RuO₄ step.^[18]

It has been reported that well-characterized ruthenium hydride species are formed by thermal decomposition of *N*-




Scheme 3. Ruthenium-catalyzed deprotection of *N*-allyl bis- γ -lactams. Reagents and conditions: a) MeONa, MeOH, RT; b) PTSA cat., Dean–Stark apparatus, toluene, reflux; c) 5 mol % [(PCy₃)₂Cl₂Ru=CHPh], toluene, 110 °C. d) RuCl₃-NaIO₄, 1,2-dichloroethane-H₂O (1:1), aqueous workup (sat. aq. NaHCO₃ + cat. Na₂CO₃). PMP=4-MeOC₆H₄; PTSA = *p*-toluenesulfonic acid.

heterocyclic-based Grubbs carbene,^[19] and it has been noted that ruthenium hydride complexes are formed as byproducts during the preparation of second-generation Grubbs catalyst.^[20] In addition, a substrate-induced decomposition route involving β -hydride transfer from a ruthenacyclobutane intermediate has been investigated.^[21] We believe that the isomerization process from *N*-allyl lactams, imides, and analo-




Scheme 4. Ruthenium-catalyzed deprotection of amides. Reagents and conditions: a) 5 mol % $[(\text{PCy}_3)_2\text{Cl}_2\text{Ru}=\text{CHPh}]$, toluene, 110 °C; b) RuCl_3 – NaIO_4 , 1,2-dichloroethane– H_2O (1:1), aqueous workup (sat. aq. NaHCO_3 + cat. Na_2CO_3).



Scheme 5. Ruthenium-catalyzed deprotection of imides, pyrazolidones, hydantoin, and oxazolidinones. Reagents and conditions: a) 5 mol % $[(\text{PCy}_3)_2\text{Cl}_2\text{Ru}=\text{CHPh}]$, toluene, 110 °C; b) RuCl_3 – NaIO_4 , 1,2-dichloroethane– H_2O (1:1), aqueous workup (sat. aq. NaHCO_3 + cat. Na_2CO_3).

gous enamide-like moieties is catalyzed by either a hydride decomposition compound or alternatively by impurities remaining from catalyst synthesis. The isomerization may occur according to the hydride mechanism, by hydrometallation followed by β -elimination, analogous to the double bond migration of allyl ethers promoted by related ruthenium complexes.^[22] This metal hydride mechanism, which requires the presence of a coordinatively unsaturated ruthenium hydride species is shown in Scheme 6. Coordination of the olefin to the metal centre gives a π -allyl complex **19**, initiating the metal hydride addition to the olefin in a Markov-



Scheme 6. Mechanistic explanation for the ruthenium Grubbs' carbene-catalyzed amide-enamide isomerization.

nikov fashion to afford a σ -alkyl complex **20**. Subsequent β -hydride elimination gives the enamide π complex **21**, which decomplexes to the free enamide regenerating the catalytically active species.

Conclusion

In conclusion, we have presented a convenient methodology for the catalytic deprotection of *N*-allylic amide-like moieties. In addition to the novelty of the method, it is general, selective, and operationally simple, offering the first ruthenium-catalyzed deallylation of amides, lactams, imides, pyrazolidones, hydantoin, and oxazolidinones. This protocol is tolerant towards different functionalities as well as the stereocentres present in the molecule. Taking together these observations, this C–N bond cleavage has the potential to significantly extend the scope of the allyl protecting group in synthesis.

Experimental Section

General methods: Melting points were taken by using a Gallenkamp apparatus and are uncorrected. IR spectra were recorded on a Perkin-Elmer 781 spectrophotometer. ^1H and ^{13}C NMR spectra were recorded on a Bruker Avance-300, Varian VRX-300S, or Bruker AC-200. NMR spectra were recorded in CDCl_3 , except when otherwise stated. Chemical shifts are given in ppm relative to TMS (^1H , 0.0 ppm) or CDCl_3 (^{13}C , 76.9 ppm). LRMS and HRMS were taken on a HP5989 A spectrometer by using the electronic impact (EI) or electrospray modes (ES) unless otherwise stated. Optical rotations were measured by using a Perkin-Elmer 241 polarimeter. Specific rotation $[\alpha]_D$ is given in $10^{-1} \text{ deg cm}^2 \text{ g}^{-1}$ at 25 °C, and the concentration (c) is expressed in g per 100 mL. All commercially available compounds were used without further purification. THF was distilled from Na-benzophenone. Benzene, dichloromethane, and triethylamine were distilled from CaH_2 . Flame-dried glassware and standard Schlenk techniques were used for moisture sensitive reactions. Flash chromatography was performed by using Merck silica gel 60 (230–400 mesh).

General procedure for the isomerization reaction of *N*-allyl amides, lactams, imides, and congeners: $[\text{Cl}_2(\text{Cy}_3\text{P})_2\text{Ru}=\text{CHPh}]$ (0.01 mmol) was added in portions under argon to a solution, protected from the sunlight, of the corresponding allylic compound (0.20 mmol) in anhydrous toluene (6 mL), and the mixture was heated at reflux. The reaction was monitored by TLC. After completion, the mixture was concentrated under reduced pressure, and was purified by column chromatography, eluting with EtOAc/hexanes mixtures to give analytically pure enamide-like compounds. Spectroscopic and analytical data for some representative pure forms of the above enamides follow.^[23]

(Z)-2d: From 110 mg (0.72 mmol) of 1-allyl-azepan-2-one **1d** and after column chromatography (hexanes/EtOAc 3:1) 79 mg (72%) of compound (Z)-**2d** was obtained as a yellow oil. ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 6.19 (dd, *J* = 8.4, 1.7 Hz, 1H), 5.14 (m, 1H), 3.40 (m, 2H), 2.51 (m, 2H), 1.63 (m, 6H), 1.51 ppm (dd, *J* = 7.1, 1.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 175.6, 133.7, 118.1, 51.0, 37.3, 29.9, 28.7, 23.3, 14.1 ppm; IR (CHCl₃): $\tilde{\nu}$ = 1652 cm⁻¹; MS (ES): *m/z* (%): 154 (100) [*M*+H]⁺, 153 (8) [*M*]⁺; elemental analysis calcd (%) for C₉H₁₅NO (153.2): C 70.55, H 9.87, N 9.14; found C 70.42, H 9.83, N 9.18.

(E)-(±)-5d: From 150 mg (0.56 mmol) of 1-allyl-β-lactam (±)-**4d** and after column chromatography (hexanes/EtOAc 5:1), 108 mg (72%) of compound (E)-(±)-**5d** was obtained as a colorless oil. ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 7.28 (m, 1H), 7.14 (m, 2H), 6.59 (m, 3H), 6.36 (m, 1H), 6.25 (m, 1H), 6.07 (dd, *J* = 9.2, 1.7 Hz, 1H), 5.42 (d, *J* = 4.6 Hz, 1H), 5.30 (d, *J* = 4.6 Hz, 1H), 4.93 (dd, *J* = 9.2, 7.3 Hz, 1H), 1.56 ppm (dd, *J* = 7.3, 1.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 163.3, 157.2, 147.6, 143.3, 129.4 (2C), 122.4, 120.8, 115.7 (2C), 114.8, 110.9, 110.5, 81.6, 58.2, 15.1 ppm; IR (CHCl₃): $\tilde{\nu}$ = 1751 cm⁻¹; MS (ES): *m/z* (%): 208 (100) [*M*+H]⁺, 207 (9) [*M*]⁺; elemental analysis calcd (%) for C₁₁H₁₃NO₃ (207.2): C 63.76, H 6.32, N 6.76; found C 63.64, H 6.29, N 6.73.

Preparation of (+)-5e: From 200 mg (0.66 mmol) of 1-allyl-β-lactam (+)-**4e** and after column chromatography (hexanes/EtOAc 2:1), 81 mg (40%) of the less polar compound (E)-(+)-**5e** and 79 mg (39%) of the more polar compound, its *Z* isomer were obtained.

(E)-(+)-5e: Colorless oil; [α]_D = +184.3 (*c* = 0.6 in CHCl₃); ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 7.23 (m, 2H), 6.99 (m, 3H), 6.36 (dd, *J* = 14.3, 1.6 Hz, 1H), 5.79 (dd, *J* = 14.3, 6.9 Hz, 1H), 5.16 (d, *J* = 5.4 Hz, 1H), 4.42 (m, 1H), 4.22 (m, 1H), 3.97 (m, 1H), 3.68 (m, 1H), 1.67 (dd, *J* = 6.8, 0.6 Hz, 3H), 1.39 (s, 3H; Me), 1.27 ppm (s, 3H; Me); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 163.0, 157.4, 129.7 (2C), 122.8, 121.9, 115.9 (2C), 112.9, 109.9, 79.7, 76.9, 67.1, 61.9, 26.9, 26.8, 25.1, 15.4 ppm; IR (CHCl₃): $\tilde{\nu}$ = 1753 cm⁻¹; MS (ES): *m/z* (%): 304 (100) [*M*+H]⁺, 303 (7) [*M*]⁺; elemental analysis calcd (%) for C₁₇H₂₁NO₄ (303.3): C 67.31, H 6.98, N 4.62; found C 67.43, H 6.94, N 4.59.

(Z)-(+)-5e: Colorless oil; [α]_D = +233.0 (*c* = 0.7 in CHCl₃); ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 7.24 (m, 2H), 6.97 (m, 3H), 5.96 (m, 1H), 5.18 (m, 2H), 4.42 (m, 1H), 4.10 (m, 2H), 3.68 (m, 1H), 1.73 (dd, *J* = 7.1, 0.6 Hz, 3H), 1.39 (s, 3H), 1.14 ppm (s, 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 163.2, 157.4, 129.7 (2C), 122.7, 121.9, 118.7, 115.9 (2C), 115.7, 109.9, 79.4, 66.9, 61.8, 26.8, 25.1, 14.3 ppm; IR (CHCl₃): $\tilde{\nu}$ = 1752 cm⁻¹; MS (ES): *m/z* (%): 304 (100) [*M*+H]⁺, 303 (10) [*M*]⁺; elemental analysis calcd (%) for C₁₇H₂₁NO₄ (303.3): C 67.31, H 6.98, N 4.62; found C 67.44, H 6.90, N 4.65.

Compound (E)-14a: From 120 mg (0.59 mmol) of 1-allyl-acetamide **13a** and after column chromatography (hexanes/EtOAc 3:1), 85 mg (71%) of compound (E)-**14a** was obtained as a colorless oil; ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 7.35 (dd, *J* = 14.3, 1.5 Hz, 1H), 6.99 (m, 2H), 6.91 (m, 2H), 4.33 (m, 1H), 3.78 (s, 3H), 1.77 (s, 3H), 1.56 ppm (dd, *J* = 6.7, 1.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 167.5, 159.1, 133.2, 129.9, 129.2, 115.1, 109.1, 55.5, 23.2, 15.1 ppm; IR (CHCl₃): $\tilde{\nu}$ = 1654 cm⁻¹; MS (EI): *m/z* (%): 205 (100) [*M*]⁺; elemental analysis calcd (%) for C₁₂H₁₅NO₂ (205.2): C 70.22, H 7.37, N 6.82; found C 70.10, H 7.34, N 6.78.

Compound (E)-17b. From 200 mg (0.926 mmol) of 2-allyl-4-methyl-1-phenyl-pyrazolidin-3-one **16b** and after column chromatography (hexanes/EtOAc 3:1), 174 mg (87%) of compound (E)-**17b** was obtained as a orange oil; ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 7.20 (m, 5H), 6.35

(dd, *J* = 9.6, 1.6 Hz, 1H), 4.83 (m, 1H), 3.89 (dd, *J* = 11.3, 7.6 Hz, 1H), 3.50 (t, *J* = 11.3 Hz, 1H), 2.70 (m, 1H), 1.53 (dd, *J* = 4.4, 1.6 Hz, 1H), 1.05 ppm (d, *J* = 7.0 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 173.0, 150.1, 129.7, 129.3, 123.6, 120.6, 118.5, 111.5, 62.9, 34.0, 13.1, 12.7 ppm; IR (CHCl₃): $\tilde{\nu}$ = 1722 cm⁻¹; MS (ES): *m/z* (%): 217 (100) [*M*+H]⁺, 216 (10) [*M*]⁺; elemental analysis calcd (%) for C₁₃H₁₆N₂O (216.3): C 72.19, H 7.46, N 12.95; found C 72.30, H 7.50, N 13.00.

General procedure for the cleavage of enamide derivatives. Preparation of free *NH*-amides, lactams, imides, and congeners: Aqueous RuCl₃ (0.2 mL, 3.5 mol %) and solid NaIO₄ (0.4 mmol) were sequentially added to a solution of the corresponding enamide derivative (0.20 mmol) in a 1,2-dichloroethane-water mixture (2 mL, 1:1). The reaction mixture was stirred at room temperature until complete disappearance of the starting material was observed (TLC), before being quenched with aqueous Na₂S₂O₃ and extracted with EtOAc. The organic phase was concentrated and the resulting residue was dissolved in acetone and stirred with saturated aq NaHCO₃ (0.33 mL) and Na₂CO₃ (0.02 mmol). The mixture was extracted with EtOAc, washed with water, dried over MgSO₄, filtered, and concentrated to give pure *NH*-compounds **3**, **6**, **9**, **12**, **15**, and **18**. Spectroscopic and analytical data for some representative pure forms of the above free *NH*-compounds follow.

***NH*-β-Lactam (±)-6a:** From 87 mg (0.385 mmol) of enamide derivative (±)-**5a**, compound (±)-**6a** (58 mg, 64%) was obtained as a colorless oil; ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 6.90 (s, 1H), 6.70 (m, 3H), 5.11 (d, *J* = 4.7 Hz, 1H), 4.65 (d, *J* = 5.7 Hz, 1H), 3.73 and 3.69 (s, each 3H), 3.16 ppm (s, 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 168.2, 153.6, 151.2, 125.4, 113.7, 113.4, 111.1, 86.9, 58.6, 55.8, 55.7, 53.4 ppm; IR (CHCl₃): $\tilde{\nu}$ = 3408, 1766 cm⁻¹; MS (ES): *m/z* (%): 238 (100) [*M*+H]⁺, 237 (7) [*M*]⁺; elemental analysis calcd (%) for C₁₂H₁₅NO₄ (237.3): C 60.75, H 6.37, N 5.90; found C 60.88, H 6.40, N 5.87.

***NH*-β-Lactam (+)-6c:** From 51 mg (0.15 mmol) of enamide derivative (+)-**5c**, compound (+)-**6c** (42 mg, 95%) was obtained as a colorless oil; [α]_D = +12.0 (*c* = 1.0 in CHCl₃); ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 6.20 (s, 1H), 4.70 (d, *J* = 4.9 Hz, 1H), 4.01 (m, 2H), 3.51 (m, 2H), 1.32 and 1.20 (s, each 3H), 0.77 (s, 9H), 0.09 and 0.01 ppm (s, each 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 166.2, 107.1, 76.9, 66.5, 57.7, 26.8, 25.5, 24.9, 17.8 ppm; IR (CHCl₃): $\tilde{\nu}$ = 3392, 1764 cm⁻¹; MS (ES): *m/z* (%): 302 (100) [*M*+H]⁺, 301 (14) [*M*]⁺; elemental analysis calcd (%) for C₁₄H₂₇NO₄Si (301.5): C 55.78, H 9.03, N 4.65; found C 55.80, H 8.99, N 4.62.

***NH*-Bis-γ-lactam (+)-12:** From 20 mg (0.049 mmol) of the corresponding enamide derivative, compound (+)-**12** (14 mg, 79%) was obtained as a orange oil; [α]_D = +67.1 (*c* = 0.8 in CHCl₃); ¹H NMR (300 MHz, CDCl₃, 25°C): δ = 7.21 (m, 5H), 6.70 (m, 4H), 4.78 (d, *J* = 6.7 Hz, 1H), 4.56 (d, *J* = 1.7 Hz, 1H), 4.30 (d, *J* = 5.1 Hz, 1H), 3.64 (s, 3H), 3.62 (m, 1H), 3.49 ppm (s, 3H); ¹³C NMR (75 MHz, CDCl₃, 25°C): δ = 172.2, 171.6, 158.7, 157.5, 129.9, 129.2, 125.4, 123.1, 116.8, 115.1, 82.3, 77.6, 68.4, 60.8, 59.2, 55.9 ppm; IR (CHCl₃): $\tilde{\nu}$ = 3390, 1722, 1719 cm⁻¹; MS (ES): *m/z* (%): 409 (100) [*M*+H]⁺, 408 (11) [*M*]⁺; elemental analysis calcd (%) for C₂₀H₂₀N₂O₅ (368.4): C 65.21, H 5.47, N 7.60; found C 65.33, H 5.44, N 7.64.

Acknowledgements

Support for this work by the DGI-MCYT (Project BQU2003-07793-CO2-01) is gratefully acknowledged. J. M. Alonso thanks the Universidad Complutense de Madrid for a fellowship.

- [1] For a pertinent review see: J. Kant, D. G. Walker in *The Organic Chemistry of β-Lactams* (Ed.: G. I. Georg), Wiley-VCH, New York, 1993.
- [2] M. Suffness, *Taxol Science and Applications*, CRC Press, Boca Raton, FL, USA, 1995.
- [3] a) S. Escoubet, S. Gastaldi, M. Bertrand, *Eur. J. Org. Chem.* **2005**, 3855; b) P. J. Kocienski, *Protecting Groups*, 3rd ed., Thieme, Stuttgart, 1994.

- gart, **2003**; c) T. W. Greene, G. M. Wuts in *Protective Groups in Organic Synthesis*, 3rd ed., Wiley, New York, **1999**; d) M. Schelhaas, H. Waldmann, *Angew. Chem.* **1996**, *108*, 2192; *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 2056.
- [4] D. K. Kronenthal, C. Y. Han, M. K. Taylor, *J. Org. Chem.* **1982**, *47*, 2765.
- [5] This methodology requires the presence of both the palladium catalyst and a nucleophilic compound as an allyl group scavenger. For a review see: F. Guibé, *Tetrahedron*, **1998**, *54*, 2967.
- [6] G. Cainelli, M. DaCol, P. Galletti, D. Giacomini, *Synlett* **1997**, 923.
- [7] O. Kanno, M. Miyauchi, I. Kawamoto, *Heterocycles* **2000**, *53*, 173.
- [8] a) B. Alcaide, P. Almendros, R. Rodríguez-Acebes, *Chem. Eur. J.* **2005**, *11*, 5708; b) B. Alcaide, P. Almendros, G. Cabrero, M. P. Ruiz, *Org. Lett.* **2005**, *7*, 3981; c) B. Alcaide, P. Almendros, R. Rodríguez-Acebes, *J. Org. Chem.* **2005**, *70*, 3198; d) B. Alcaide, P. Almendros, M. C. Redondo, *Org. Lett.* **2004**, *6*, 1765; e) B. Alcaide, P. Almendros, J. M. Alonso, M. F. Aly, *Chem. Eur. J.* **2003**, *9*, 3415; f) B. Alcaide, P. Almendros, C. Aragoncillo, *Chem. Eur. J.* **2002**, *8*, 3646; g) B. Alcaide, P. Almendros, C. Aragoncillo, *Chem. Eur. J.* **2002**, *8*, 1719.
- [9] For a preliminary communication of a part of this work see: B. Alcaide, P. Almendros, J. M. Alonso, *Tetrahedron Lett.* **2003**, *44*, 8693.
- [10] For reviews on this subject see: a) B. Alcaide, P. Almendros, *Chem. Eur. J.* **2003**, *9*, 1258; b) B. Schmidt, *Eur. J. Org. Chem.* **2004**, 1865.
- [11] a) B. Alcaide, P. Almendros, J. M. Alonso, A. Luna, *Synthesis* **2005**, 668; b) B. Alcaide, P. Almendros, J. M. Alonso, *Chem. Eur. J.* **2003**, *9*, 5793; c) B. Alcaide, P. Almendros, J. M. Alonso, M. F. Aly, *Org. Lett.* **2001**, *3*, 3781.
- [12] For metal-catalyzed isomerizations of *N*-allylamides to enamides see: a) B. Neugnot, J.-C. Cintrat, B. Rousseau, *Tetrahedron* **2004**, *60*, 3575; b) S. Sergeyev, M. Hesse, *Synlett* **2002**, 1313; c) S. Krompiec, M. Pigulla, W. Szczepankiewicz, T. Bieg, N. Kuznik, K. Leszczynska-Sejda, M. Kubicki, T. Borowiak, *Tetrahedron Lett.* **2001**, *42*, 7095; d) J. K. Stille, Y. Becker, *J. Org. Chem.* **1980**, *45*, 2139.
- [13] a) D. Yang, C. Zhang, *J. Org. Chem.* **2001**, *66*, 4814; b) P. K. Sharma, P. Nielsen, *J. Org. Chem.* **2004**, *69*, 5742; c) for a review on Ru-based oxidation catalysis see: M. Pagliaro, S. Campestrini, R. Ciri-minna, *Chem. Soc. Rev.* **2005**, *34*, 837.
- [14] a) J. M. Aizpurua, F. P. Cossío, C. Palomo, *Tetrahedron Lett.* **1986**, *27*, 4359; b) C. Palomo, J. M. Aizpurua, M. Legido, A. Mielgo, R. Galarza, *Chem. Eur. J.* **1997**, *3*, 1432; c) T. G. Back, J. E. Wulff, *Angew. Chem.* **2004**, *116*, 6655; *Angew. Chem. Int. Ed.* **2004**, *43*, 6496.
- [15] For the synthesis of related bis- β - and bis- γ -lactams see: a) B. Alcaide, Y. Martín-Cantalejo, J. Pérez-Castells, M. A. Sierra, A. Monge, *J. Org. Chem.* **1996**, *61*, 9156; b) B. Alcaide, Y. Martín-Cantalejo, J. Pérez-Castells, J. Rodríguez-López, M. A. Sierra, A. Monge, V. Pérez-García, *J. Org. Chem.* **1992**, *57*, 5921.
- [16] Enantiopure 2-azetidinone (+)-**7** was obtained as a single *cis*-enantiomer by using our own methodology from the imine of (*R*)-2,3-*O*-isopropylidene-glyceraldehyde and *p*-anisidine, through a Staudinger reaction with the phenoxyacetyl chloride in the presence of Et₃N, followed by sequential acidic acetone hydrolysis and oxidative cleavage. See reference [8].
- [17] The higher load of Grubbs carbene required may be due to the poisoning of the catalyst by the sulfur functionalities.
- [18] It is presumed that overoxidation of the sulfur moieties takes place.
- [19] For the characterization of a decomposition product arising from a methylidene Ru NHC complex see: a) S. H. Hong, M. W. Day, R. H. Grubbs, *J. Am. Chem. Soc.* **2004**, *126*, 7414; b) For decomposition studies of the first generation Grubbs carbene on the basis of variable temperature ¹H NMR experiments see: J. Faulkner, C. D. Edlin, D. Fengas, I. Preece, P. Quayle, S. N. Richards, *Tetrahedron Lett.* **2005**, *46*, 2381.
- [20] A. Fürstner, L. Ackermann, B. Gabor, R. Goddard, C. W. Lehmann, R. Mynott, F. Stelzer, O. R. Thiel, *Chem. Eur. J.* **2001**, *7*, 3236.
- [21] W. J. van Rensburg, P. J. Steynberg, W. H. Meyer, M. M. Kirk, G. S. Forman, *J. Am. Chem. Soc.* **2004**, *126*, 14332.
- [22] D. V. McGrath, R. H. Grubbs, *Organometallics* **1994**, *13*, 224.
- [23] Full spectroscopic and analytical data for compounds not included in this Experimental Section are described in the Supporting Information.

Received: October 6, 2005
Published online: January 18, 2006