

A Convenient Method for the Preparation of Acid Anhydrides from Metallic Carboxylates

Tsujiaki HATA, Kiyohiko TAJIMA and Teruaki MUKAIYAMA


Laboratory of Organic Chemistry, Tokyo Institute of Technology, Ookayama, Meguro-ku, Tokyo

(Received May 2, 1968)

A convenient method for the preparation of carboxylic anhydrides by use of one mole of *N,N'*-dicyclohexylthiourea and two moles of silver or mercurous carboxylate has been established. According to this method, several carboxylic anhydrides were obtained in high yields under mild conditions even by microscale experiments.

In previous papers,^{1,2)} it has been reported that carboxylic anhydrides were prepared from the corresponding mercuric carboxylates by treatment with the addition compound of dibenzoyl ethylene and triphenylphosphine, or by treatments with *S*-ethylthioamidium iodide.

In the present study, a general preparative method of acid anhydrides was investigated with the expectation that acid anhydride would be formed by the reaction of *N,N'*-dicyclohexylthiourea with silver carboxylates through an intermediate, imidoacylate. Indeed, when one mole of *N,N'*-dicyclohexylthiourea was added to a suspension of two moles of finely pulverized silver benzoate in anhydrous acetone at room temperature, benzoic anhydride was obtained in 93% yield along with *N,N'*-dicyclohexylurea and silver sulfide.




In a similar manner, acetic, propionic, phthalic and succinic anhydrides were obtained from the corresponding silver carboxylates. These data are recorded in Table 1.

TABLE 1. ACID ANHYDRIDES FROM SILVER CARBOXYLATES


Silver salt (0.04 mol)	Yield of acid anhydride, %	Solvent	ml
Acetate	68.0	Acetone	13
Propionate	88.4	Acetone	20
Benzoate	93.0	Acetone	10
Phthalate	86.9	Acetonitrile	30
Succinate	90.0	Chloroform	30

The reaction is considered to proceed through an intermediate (IV), produced from a silver carbox-

ylate and *N,N'*-dicyclohexylthiourea. The IV in turn reacts with another silver carboxylate to form silver sulfide and the imidoacylate (V), which further reacts with carboxylic acid to give an acid anhydride and *N,N'*-dicyclohexylurea.



In addition, it was found that mercuric acetate also reacts with *N,N'*-dicyclohexylthiourea to give acetic anhydride, *N,N'*-dicyclohexylurea and mercuric sulfide along with a considerable amount of *N*-acetyl-*N,N'*-dicyclohexylurea (VI), a rearrangement product of imidoacylate (Va).



Some mercuric carboxylate, such as mercuric benzoate, contains one mole of water of crystallization, which cannot be removed by the ordinary procedure. Since water reacts with imidoacylate faster than with carboxylate ion, mercuric carboxylates containing water of crystallization cannot be used in this reaction. Next, since anhydrous mercurous carboxylates are easily prepared from sodium carboxylates and mercurous nitrate,

1) I. Kuwajima and T. Mukaiyama, *J. Org. Chem.*, **29**, 1385 (1964).

2) T. Yamaguchi, K. Inomata and T. Mukaiyama, *This Bulletin*, **41**, 673 (1968).

