

- (4) Rupp, H.; Weser, U. *Biochim. Biophys. Acta* **1976**, *446*, 151.
 (5) Brisk, M. A.; Baker, A. D. *J. Electron Spectrosc. Relat. Phenom.* **1975**, *7*, 197.
 (6) Peeling, J.; Haslett, B. G.; Evans, I. M.; Clark, D. T.; Boulter, D. *J. Am. Chem. Soc.* **1977**, *99*, 1025.
 (7) Best, S. A.; Brant, P.; Feltham, R. D.; Rauchfuss, T. B.; Roundhill, D. M.; Walton, R. A. *Inorg. Chem.* **1977**, *16*, 1976.
 (8) Amundsen, A. R.; Whelan, J.; Bosnich, B. *J. Am. Chem. Soc.* **1977**, *99*, 6730.
 (9) Metal-catalyzed oxidation of protein sulfur ligands has also been suggested as the origin of the 168-eV S_{2p} peaks by Walton and co-workers (Srinivasan, V.; Steifel, E. I.; Elsberry, A.; Walton, R. A. *J. Am. Chem. Soc.* submitted for publication); it should be noted that metals other than copper may also be effective, as a 168-eV S_{2p} peak is observed¹ in the XPE spectrum of cobalt(II) plastocyanin.
 (10) Larsson, S. *J. Am. Chem. Soc.* **1977**, *99*, 7708.
 (11) Grunthaner, F. J.; Wurzbach, J. A., personal communication.
 (12) Ghirelli-Magaldi, A.; Nuzzolo, C.; Ghirelli, F. *Biochemistry* **1966**, *5*, 1943.

Michael Thompson,* John Whelan
 Deborah J. Zemon, B. Bosnich*

Department of Chemistry, University of Toronto
 Toronto, Ontario, Canada M5S 1A1

E. I. Solomon*

Department of Chemistry
 Massachusetts Institute of Technology
 Cambridge, Massachusetts 02139

Harry B. Gray*


Contribution No. 5849
 Arthur Amos Noyes Laboratory of Chemical Physics
 California Institute of Technology
 Pasadena, California 91125

Received August 18, 1978

Anthracylines and Related Substances. 2. An Efficient and Regiospecific Synthesis of dl-7,9-Dideoxydaunomycinone¹


Sir:

Of the group of compounds that comprise the anthracylines,² two substances, namely daunomycin³ (**1**) and adriamycin⁴ (**2**), have achieved preeminence as antitumor agents, despite the fact that they frequently induce an irreversible cardiomyopathy. This has initiated a search for partially or totally synthetic derivatives that might lack this side effect.⁵




However efforts aimed at the total syntheses of **1** and **2** frequently have been plagued⁶ by the problem of regioisomerism posed by the relationship of the C-4 methoxyl and the substituents on ring A. In cases where this problem has been solved, the methods⁷ appear to lack efficiency or alternatively, seem inapplicable to large-scale work. We now describe an efficient synthesis (20% overall yield) of dl-7,9-dideoxydaunomycinone (**3**), a compound whose conversion into the aglycones of **1** and **2** is known.^{6b,8} The procedure is regiospecific and is adaptable to bulk preparative work.

The overall method of construction follows a convergent B + D → BD → ABD → ABCD pattern and called initially for the synthesis of an intermediate belonging to the class represented by **4** (X and Y being suitably functionalized carbon



atoms). The crucial step in the synthesis of molecules of this type is a new Friedel-Crafts *alkylation* which directly introduces a phthalido residue (an inductively deactivating group⁹) into the aromatic nucleus destined to be ring B. In the case under consideration the reaction of 3-bromo-4-methoxyphthalide **5** (obtained almost quantitatively by the action of *N*-bromosuccinimide on 4-methoxyphthalide¹⁰) with the triester **6a** (SnCl_4 , CH_2Cl_2 , 25 °C, 6 h) led to **7a**¹¹ in 94% yield: mp 135–137 °C. Selective hydrolysis of the terminal methyl ester of **7a** (Scheme I) (1.3 equiv¹² of KOH, THF- CH_3OH -

Scheme I



(62% from 11a) as yellow needles: mp 124–125 °C; ¹H NMR (CDCl₃) δ 7.2–7.8 (m, 3 H), 4.25 (q, 4 H, *J* = 7 Hz), 4.05 (s, 3 H), 3.9 (s, 3 H), 3.3 (s, 2 H), 2.9 (t, 2 H, *J* = 6 Hz), 2.3 (t, 2 H, *J* = 6 Hz), 1.3 (t, 6 H, *J* = 7 Hz). The conversion of 14a into 7,9-dideoxydaunomycinone dimethyl ether (17a) was then effected by the following four-part procedure.

Saponification (KOH, aqueous ethanol (1:2), 90 °C, 3 h, 98%) of 14a led to the diacid 15a as yellow needles, mp 222–224 °C from CH₂Cl₂/Et₂O. This was the decarboxylated (CH₃CO₂H, piperidine, 120 °C, 1 h) to give monocarboxylic acid 16a (85% from 14a): mp 133.5–135 °C; ¹H NMR (acetone-*d*₆) δ 7.6–7.9 (m, 3 H), 4.1 (s, 3 H), 4.0 (s, 6 H), 2.7–3.1 (m, 7 H), ~9 (very br, 1 H). The crude acid chloride derived (SOCl₂, C₆H₆, 25 °C, 15 h) from 16a was then treated with lithium dimethylcuprate¹⁵ (THF/Et₂O, –78 to 0 °C, 3 h) and afforded 7,9-dideoxydaunomycinone dimethyl ether (17a, 80% based on 16a) as yellow needles: mp 185–186 °C; ¹H NMR (acetone-*d*₆) δ 7.6–7.9 (m, 3 H), 4.1 (s, 3 H), 3.9 (s, 6 H), 2.8–3.1 (m, 7 H), 2.3 (s, 3 H).

Selective demethylation of 17a to give *dl*-7,9-dideoxydaunomycinone (3) was possible only by a two-part sequence, namely oxidation^{6b,16} (AgO/HNO₃, aqueous acetone, 70 °C, 1 h) to the 4,12:6,11-bisquinone, followed by reduction (Et₂NOH, EtOAc, 25 °C, 30 min) of the crude product. This afforded 3 in 83% yield after recrystallization from CH₂Cl₂/Et₂O: mp 243–245 °C, no depression in melting point when admixed with an authentic sample (mp 243–245 °C); ¹H NMR (CDCl₃) δ 13.78 (s, 1 H), 13.43 (s, 1 H), 8.1–7.2 (m, 3 H), 2.27 (s, 3 H), 2.15 (m, 1 H), 1.55 (m, 2 H). The NMR, IR (Nujol), visible (CH₂Cl₂), and mass spectra were identical with those recorded in the literature^{8a} for 3.

Although the demethylation of 17a is an efficient process, the initial oxidation is rather vigorous and one could envisage that more delicate molecules may not survive. To avoid this difficulty we have developed an alternative procedure based on the fact that aryl ethyl ethers are more readily cleaved¹⁷ by Lewis acids than the corresponding methyl ethers. Repetition then, of the complete synthetic sequence¹⁸ starting with 6b produced in comparable yields the corresponding diethoxy homologues 7b through 17b. Selective deethylation of 17b to give 3 was then easily accomplished in one step under mild conditions (AlCl₃/PhNO₂, 45 °C, 40 min, 80%).

We believe that the methods presented above, together, constitute a very versatile approach to the anthracyclines in general. Variations in the substitution patterns of rings A, B, and D and in the nature of the C-9 side chain now seem possible, not only because of the convergent nature of the synthesis and its regiospecificity but also because of the relatively simple nature of the reactions involved. Investigations into the use of these procedures for the synthesis of other classes of anthracyclines are underway.

Acknowledgment. The authors are grateful to Dr. F. Arcamone (Farmitalia) for a generous financial gift in support of the research. The project was also partially supported by grants from the National Cancer Institute (Grant CA20197) and from the State University of New York. The technical assistance of Mr. John Winter, Mr. Michael W. Spatz, and Miss Nancy Stambler is acknowledged.

Reference and Notes

- (1) K. S. Kim, M. W. Spatz, and Francis Johnson, *Tetrahedron Lett.*, 331 (1979).
- (2) H. Brockmann, *Prog. Chem. Org. Nat. Prod.*, **21**, 121 (1963).
- (3) (a) F. Arcamone, G. Cassinelli, G. Fantini, A. Grein, P. Orezzi, C. Pol, and C. Spalla, *Biotechnol. Bioeng.*, **11**, 1101 (1969); (b) F. Arcamone, G. Franceschi, P. Orezzi, G. Cassinelli, W. Barbieri, and R. Mondelli, *J. Am. Chem. Soc.*, **86**, 5334 (1964).
- (4) A. DiMarco, M. Gaetani, and B. Scarpinato, *Cancer Chemother. Rep.*, **53**, 33 (1969).
- (5) D. W. Henry in "Cancer Chemotherapy", A. C. Sartorelli, Ed., American Chemical Society, Washington, D.C., 1976, Chapter 2.
- (6) (a) C. M. Wong, R. Schwenk, D. Dopien, and T. L. Ho, *Can. J. Chem.*, **51**, 466 (1973); (b) A. S. Kende, Y. G. Tsay, and J. E. Mills, *J. Am. Chem. Soc.*, **98**, 1967 (1976).
- (7) (a) P. W. Reynolds, M. J. Manning, and J. S. Swenton, *Tetrahedron Lett.*, 2383 (1977); J. S. Swenton and P. W. Reynolds, *J. Am. Chem. Soc.*, **100**, 6188 (1978); (b) F. Suzuki, S. Trenbeath, R. D. Gleim, and C. J. Sih, *ibid.*, **100**, 2272 (1978).
- (8) (a) R. D. Gleim, S. Trenbeath, R. S. D. Mittal, and C. J. Sih, *Tetrahedron Lett.*, 3385 (1976); (b) F. Arcamone, G. Franceschi, and S. Penco, U.S. Patent 3 803 124 (1974).
- (9) Details of other research work which shows that the 3-phthalido group does indeed have an electron-withdrawing effect may be found in part 1 of this series.¹ However, the effect, in part at least, can be equated with the substitution of chlorine or acetoxy in the α position of an aralkane which leads to diminished reactivity of the aromatic ring toward electrophilic substitution. This is well illustrated for example, by the fact that chloromethylmesitylene is 46 times less reactive than mesitylene itself toward chloromethylation: G. Vavon, J. Bolle, and J. Calin, *Bull. Soc. Chim. Fr.*, **6**, 1025 (1930). The coordination of the Friedel-Crafts catalyst with the lactone of the product probably also makes a contribution to the diminished reactivity of the aromatic ring.
- (10) C. A. Buehler, T. A. Powers, and J. G. Michels, *J. Am. Chem. Soc.*, **66**, 417 (1944).
- (11) Suitable analytical data were obtained for all new compounds discussed in this paper.
- (12) The base-induced opening of the lactone is competitive with the hydrolysis of the methyl ester. However, the former reaction appears to be a reversible reaction so that, with considerably less than 2 equiv of base, the desired ester hydrolysis goes to completion.
- (13) It should be noted that the presence of the two ethyl esters on the side chain is essential to good yields in this reaction. In the analogous compound having only one ester group at this position, cyclization with this reagent cannot be induced. This appears to be an outstanding example of the effects of B strain: H. C. Brown, H. Bartholamy, and M. D. Taylor, *J. Am. Chem. Soc.*, **94**, 5106 (1972). For another example, see E. Testa and L. Fontanella, *Justus Liebigs Ann. Chem.*, **625**, 94 (1959).
- (14) K. Bowden, I. M. Heilbron, E. R. H. Jones, and B. C. L. Weedon, *J. Chem. Soc.*, 39 (1946).
- (15) G. H. Posner, C. E. Whitten, and P. E. McFarland, *J. Am. Chem. Soc.*, **94**, 5106 (1972).
- (16) C. D. Snyder and H. Rapoport, *J. Am. Chem. Soc.*, **94**, 277 (1972).
- (17) C. Szantay, *Acta Chim. Acad. Sci. Hung.*, **12**, 83 (1957).
- (18) The 1,4-diethoxybenzene needed for this synthesis is easily obtained (88%) by the ethylation of hydroquinone (Et₂SO₄/KOH, H₂O–CH₂Cl₂, Pr₄, NOH, 25 °C, 36 h). Formylation (Cl₂CHOCH₃, SnCl₄, CH₂Cl₂, 0 °C, 20 min) then yields 2,5-diethoxybenzaldehyde (85%). Conversion of the latter into 6b follows exactly the three-step procedure (93% overall) used for 6a, namely condensation with diethyl malonate, catalytic reduction (H₂, Pd/C, EtOAc), and alkylation with methyl bromoacetate (NaH, C₆H₆, 25 °C). The melting points (°C) of the b series are as follows: 2,5-diethoxybenzaldehyde, 60–61; diethyl 2,5-diethoxybenzylidenemalonate 52–53; 7b, 122–126; 8b, 150–151; 9b, 111–113; 11b, 110–111; 14b, 111–113; 15b, 198–200 dec; 16b, 213–214; 17b, 147.

K. S. Kim, Ermes Vanotti

Antonino Suarato, Francis Johnson*

Departments of Pharmacological Sciences and Chemistry
State University of New York at Stony Brook
Stony Brook, New York 11794

Received September 20, 1978

2-Hydroperoxyhexafluoro-2-propanol.

A Low-Cost, Catalytic Oxidant for Synthesis and a Structural Analogue of Naturally Occurring Flavin Hydroperoxides

Sir:

Organic chemists have long been interested in utilizing hydrogen peroxide directly for the epoxidation of simple, unactivated alkenes. Efforts to devise a workable process using H₂O₂ to drive the carboxylic acid-peracid exchange have been unsuccessful to date since a strong acid catalyst is required.¹ Transition metal oxides and peroxides achieve a ready equilibrium but are poor oxidants for isolated double bonds.² Only recently have the corresponding seleninic-peroxyseleninic acid systems been described as satisfactory alternatives, although they offer little, if any, regio- or stereoselectivity.^{3,4}

Since our discovery⁵ that peroxytrifluoroacetic acid esterifies alcohols by a Fischer-type mechanism (eq 1), we have been exploring the chemistry of electron-deficient hydroperoxides related to 1. We now report that 2-hydroperoxyhexafluoro-