

Cleavage of unactivated amide bonds by ammonium salt-accelerated hydrazinolysis†

Cite this: DOI: 10.1039/c4cc02014f

Yuhei Shimizu, Megumi Noshita, Yuri Mukai, Hiroyuki Morimoto* and Takashi Ohshima*

 Received 18th March 2014,
Accepted 23rd May 2014

DOI: 10.1039/c4cc02014f

www.rsc.org/chemcomm

Hydrazinolysis of unactivated amide bonds is significantly accelerated by the addition of ammonium salts. The reactions proceed at 50–70 °C to give amines with broad substrate scope that outperforms existing amide bond cleavage reactions. Application to peptide and amino sugar derivatives is also demonstrated.

An amide is a ubiquitous structural motif found in various organic molecules such as peptides, natural products, and pharmaceuticals, because of the high stability of amide bonds.¹ The recent development of catalytic methods, such as C–H activation,² hydrogenation,³ and kinetic resolution,⁴ expands the utility of amides as useful directing/protecting groups in organic reactions. In contrast to the utility of amides in organic synthesis, cleavage of amide bonds is less explored due to the high stability of amide bonds, and the most general method to cleave amide bonds is conventional hydrolysis under harsh acidic/basic conditions with limited functional group tolerance.⁵ Recent efforts to circumvent the problem have provided several elegant solutions,⁶ but these reactions still require anhydrous conditions and/or high reaction temperatures and have limited functional group tolerance. To expand the utility of simple, unactivated amides for the synthesis of organic molecules, the development of general and practical methods to cleave amide bonds under mild conditions is necessary.⁷

To overcome these limitations, we were interested in the hydrazinolysis of amide bonds. Hydrazinolysis of activated amide bonds is widely utilized, especially for the removal of phthaloyl groups under mild conditions in Gabriel amine synthesis.⁸ Hydrazinolysis of simple unactivated amides is generally very difficult, however, because of the lack of intramolecular assistance during cleavage of the amide bonds, and a high temperature and long reaction time are necessary to provide amine **2** in a reasonable yield in most cases.⁹ Hydrazinolysis of simple amide bonds under mild conditions was limited to only amino acid derivatives

and readily cleavable formamides,¹⁰ and has not been recognized as a general method to cleave simple, unactivated amide bonds because of the limited reactivity of hydrazinolysis under the reported reaction conditions (*vide infra*). To improve the applicability of hydrazinolysis, we hypothesized that the addition of ammonium salts would activate amide bonds and promote hydrazinolysis at a lower reaction temperature similar to the transamidation reaction using ethylenediamine.^{11,12} Herein, we report ammonium salt-accelerated hydrazinolysis of unactivated amides using readily available hydrazine monohydrate as the cleaving agent. The reactions proceeded at 50–70 °C to provide amines in good yields with broad substrate scope that outperforms that of existing amide bond cleavage reactions.

To test our hypothesis, we first explored the reaction conditions using hydrazine hydrate at 60 °C for 6 h under conventional heating conditions (Table 1). The reactions proceeded only sluggishly under the conditions reported in the previous literature (entries 1^{10a} and 2^{10b}).¹³ By contrast, the addition of ammonium salts significantly accelerated the reaction to provide amine **2a**, and the addition of ammonium iodide showed the highest reactivity

Table 1 Optimization of reaction conditions

Entry	Ammonium salt	Temp. (°C)	<i>t</i> (h)	Yield ^a (%)
1	None	60	6	n.d. ^b
2	H ₂ NNH ₂ ·AcOH ^c	60	6	7
3	NH ₄ Cl	60	6	35
4	NH ₄ Br	60	6	45
5	NH ₄ I	60	6	52
6	<i>n</i> Bu ₄ NI	60	6	n.d. ^b
7	NH ₄ I	60	24	90 ^d
8	NH ₄ I	50	48	89 ^d
9 ^e	NH ₄ I	50	5	90 ^d

^a Yield was determined by ¹H NMR analysis of the crude mixture. ^b Not detected. ^c 3.0 equiv. of H₂NNH₂·H₂O and AcOH were used in EtOH (1.0 M). ^d Isolated yield. ^e Under microwave irradiation conditions.

Graduate School of Pharmaceutical Sciences, Kyushu University, Fukuoka, 812-8582, Japan. E-mail: ohshima@phar.kyushu-u.ac.jp, hmorimoto@phar.kyushu-u.ac.jp

† Electronic supplementary information (ESI) available: Experimental details and characterization data. See DOI: 10.1039/c4cc02014f

(entries 3–5). No reaction proceeded with tetrabutylammonium iodide, suggesting that ammonium salt functions as a proton source (entry 6). Prolonging the reaction time to 24 h gave satisfactory results (entry 7), and the reaction proceeded even at 50 °C in 89% yield (entry 8). Although the conventional heating conditions were sufficient to promote amide bond cleavage as described above, a significant acceleration of the amide bond cleavage was observed under microwave irradiation conditions to give **2a** in 90% yield for 5 h at 50 °C (entry 9).¹⁴

With the optimized conditions in hand, we performed hydrazinolysis of various amides **1** (Table 2). The reactions proceeded at 50–70 °C even under conventional heating conditions to give amine **2** in 78–99% yield. Some of the reactions were performed under microwave irradiation conditions to provide amines in a shorter reaction time (entries 3, 5 and 8). Various secondary and tertiary amides were also reacted, and broad functional groups were tolerant under the current hydrazinolysis conditions. Amides **1b**, **1c**, and **1d** having acetamide, *tert*-butoxycarbonyl, and benzyl-oxycarbonyl functionalities, respectively, also gave products in good yields, indicating the mildness of the current reaction conditions.¹³ It is noted that the cleavage of sterically congested formamide **1i** was slow without addition of ammonium iodide (entry 8 in the parenthesis), indicating the importance of ammonium iodide as an additive to accelerate reactions.

The broad functional group tolerance under the current reaction conditions was further confirmed for amide **1p**, which has an acid- and base-sensitive *tert*-butyldimethylsilyloxy group (Scheme 1). As expected, the desired product **2p** was obtained in 89% yield with retention of the TBS group under our reaction conditions. On the other hand, undesired desilylated product **1l** and **2l** were obtained as major products under conventional acidic and basic hydrolysis conditions.


Finally, we applied our reaction conditions to cleave peptide and amino sugar amide bonds (Scheme 2). Selective deacetylation of alanine derivative **1q** was realized under the current conditions. Furthermore, selective cleavage of the Gly–Phe bond was realized when peptide **1r** and **1s** were exposed to the current reaction conditions to give amine **2r** in 84% and 74% yields (92% based on recovered starting material), respectively, without racemization at the α -position of the phenylalanine residue.¹³ The observed site-selectivity was presumably affected by both steric and electronic nature of amide bonds.¹³ In addition, this reaction condition was also applied to cleavage of an *N*-acetyl group from the amino sugar derivative **1t**. Without the addition of ammonium salt, a high temperature and long reaction time were required to obtain **2t**,^{9a} but with ammonium iodide the reaction proceeded at 60 °C in a shorter reaction time under microwave irradiation conditions to give the desired product **2t** in 81% yield.

In summary, we developed ammonium salt-accelerated hydrazinolysis of unactivated amides. The reactions proceeded by using readily available ammonium iodide and hydrazine monohydrate under mild conditions (50–70 °C without strong acids and bases) with broad substrate generality, including protic functional groups, which is superior to the conventional acidic/basic hydrolysis conditions and recently reported amide bond cleavage reactions. Application to peptide and amino sugar derivatives further confirmed the


Table 2 Scope and limitations of hydrazinolysis of amides

Entry	Amide 1	Temp. (°C)	<i>t</i> (h)	Yield ^a (%)
1		60	36	78
2 ^b		70	45	85
3 ^c		60	4	79
4		60	48	92
5 ^c		70	10	84
6		50	36	83
7 ^b		50	18	85
8 ^{b,c}		70	5	92 (26) ^d
9 ^b		60	42	97
10		70	48	84
11		50	18	94
12 ^b		60	45	94
13		50	12	97
14		50	12	99

^a Isolated yield. ^b Reaction was performed in EtOH. ^c Under microwave irradiation conditions. ^d Without addition of NH₄I.



Scheme 1 Comparison with conventional acidic/basic hydrolysis conditions.



Scheme 2 Application to peptide and amino sugar derivatives.

applicability of our method. Further experiments are underway in our laboratory to gain a detailed understanding of the reaction mechanism and to determine further applications.¹⁵

This work was supported by the Grant-in-Aid for Scientific Research (B) (#24390004), the Scientific Research on Innovative Areas (#24106733) and Platform for Drug Discovery, Informatics, and Structural Life Science from MEXT, CREST from JST, Uehara Memorial Foundation, and the Naito Foundation. H.M. thanks

Takasago Award in Synthetic Organic Chemistry, Japan and Y.S. thanks JSPS Research Fellowships for Young Scientists. We thank the research group of Prof. Hiroshi Suemune at Kyushu University for the use of a polarimeter and Dr Yoshimasa Matsushima of Takasago Int. Co. for DSC analysis.

Notes and references

- W. Mabej and T. Mill, *J. Phys. Chem. Ref. Data*, 1978, 7, 383.
- Review of C–H activation: (a) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, 110, 1147; (b) L. Ackermann, R. Vicente and A. R. Kapdi, *Angew. Chem., Int. Ed.*, 2009, 48, 9792; (c) X. Chen, K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2009, 48, 5094; (d) O. Daugulis, H.-Q. Do and D. Shabashov, *Acc. Chem. Res.*, 2009, 42, 1074.
- Review of hydrogenation: (a) K. Gopalaiah and H. B. Kagan, *Chem. Rev.*, 2011, 111, 4599; (b) J.-H. Xie, S.-F. Zhu and Q.-L. Zhou, *Chem. Rev.*, 2011, 111, 1713; (c) T. C. Nugent and M. El-Shazly, *Adv. Synth. Catal.*, 2010, 352, 753.
- Review of kinetic resolution: (a) E. Busto, V. Gotor-Fernández and V. Gotor, *Chem. Rev.*, 2011, 111, 3998; (b) C. E. Müller and P. R. Schreiner, *Angew. Chem., Int. Ed.*, 2011, 50, 6012; (c) H. Pellissier, *Adv. Synth. Catal.*, 2011, 353, 1613.
- P. G. M. Wuts and T. W. Greene, *Protective Groups in Organic Synthesis*, Wiley-Interscience, Hoboken, 4th edn, 2006, pp. 773–789.
- For selected recent reactions involving unactivated amide bond cleavage step, see: (a) P. R. Sultane, T. B. Mete and R. G. Bhat, *Org. Biomol. Chem.*, 2014, 12, 261; (b) Y. Kita, Y. Nishii, A. Onoue and K. Mashima, *Adv. Synth. Catal.*, 2013, 355, 3391; (c) Y. Kita, Y. Nishii, T. Higuchi and K. Mashima, *Angew. Chem., Int. Ed.*, 2012, 51, 5723; (d) E. Balaraman, B. Gnanaprakasam, L. J. W. Shimon and D. Milstein, *J. Am. Chem. Soc.*, 2010, 132, 16756; (e) N. A. Stephenson, J. Zhu, S. H. Gellman and S. S. Stahl, *J. Am. Chem. Soc.*, 2009, 131, 10003; (f) S. G. Koenig, C. P. Vandebosche, H. Zhao, P. Mousaw, S. P. Singh and R. P. Bakale, *Org. Lett.*, 2009, 11, 433 and references therein.
- For selected recent examples of facile cleavage of strained amide bonds, see: (a) J. Aubé, *Angew. Chem., Int. Ed.*, 2012, 51, 3063; (b) M. Hutchby, C. E. Houlden, M. F. Haddow, S. N. G. Tyler, G. Lloyd-Jones and K. I. Booker-Milburn, *Angew. Chem., Int. Ed.*, 2011, 51, 548; (c) K. Tani and B. M. Stoltz, *Nature*, 2006, 441, 731.
- J. March, *Advanced Organic Chemistry: Reactions, Mechanisms and Structures*, John Wiley & Sons, New York, 5th edn, 2001, p. 513, and references cited therein.
- Selected examples of hydrazinolysis of unactivated amides: (a) M. J. G. Stewart, V. K. Jothivasan, A. S. Rowan, J. Wagg and C. J. Hamilton, *Org. Biomol. Chem.*, 2008, 6, 385; (b) K. Fukatsu, O. Uchikawa, M. Kawada, T. Yamano, M. Yamashita, K. Kato, K. Hirai, S. Hinuma, M. Miyamoto and S. Ohkawa, *J. Med. Chem.*, 2002, 45, 4212; (c) L. H. Jones, L. J. Altobelli III, M. T. MacDonald, N. A. Boyle, P. Wentworth, Jr., R. A. Lerner and K. D. Janda, *Angew. Chem., Int. Ed.*, 2002, 41, 4241; (d) S. Knapp, S. Gonzalez, D. S. Myers, L. L. Eckman and C. A. Bewley, *Org. Lett.*, 2002, 4, 4337.
- (a) D. D. Keith, J. A. Tortora and R. Yang, *J. Org. Chem.*, 1978, 43, 3711; (b) R. Geiger and W. Siedel, *Chem. Ber.*, 1968, 101, 3386; (c) M. Miyamoto, Y. Kawamatsu, M. Shinohara and Y. Ueno, *Yakugaku Zasshi*, 1961, 81, 439.
- Y. Shimizu, H. Morimoto, M. Zhang and T. Ohshima, *Angew. Chem., Int. Ed.*, 2012, 51, 8564.
- Although addition of hydrazinium sulfate have been reported for the analysis of peptide sequences, these reaction conditions require *anhydrous* hydrazine and not widely utilized in the field of organic synthesis; see: J. H. Bradbury, *Nature*, 1956, 178, 912.
- See ESI† for detail.
- Safety testing consisting of DSC testing of reaction mixtures was performed. No significant barriers were found to progression of the reaction ($50\text{--}70^\circ\text{C}$) and an exothermic process started at about 160°C (calorific value: $<100\text{ J g}^{-1}$, activation energy: $ca. 700\text{ kJ mol}^{-1}$).
- (a) Our preliminary results for the application of hydrazinolysis of amide bonds to the removal of directing groups were presented: Y. Shimizu, M. Noshita, Y. Mukai, H. Morimoto and T. Ohshima, 104th Symposium on Organic Synthesis, Japan, O-12, November 6–7, 2013; (b) The presented method was applied and cited in the very recent publication: K. S. Kanyiva, Y. Kuninobu and M. Kanai, *Org. Lett.*, 2014, 16, 1968.