



## Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

### BENZYLTRIPHENYLPHOSPHONIUM DICHROMATE AS A MILD REAGENT FOR THE OXIDATION OF ORGANIC COMPOUNDS

Abdol Reza Hajipour <sup>a</sup>, Iraj Mohammadpoor-Baltork <sup>b</sup> & Kurosh Niknam <sup>a</sup>

<sup>a</sup> College of Chemistry, Isfahan University of Technology, Isfahan, 84156, Iran

<sup>b</sup> Department of Chemistry, Isfahan University, Isfahan, 81744, Iran  
Version of record first published: 18 Feb 2009.

To cite this article: Abdol Reza Hajipour, Iraj Mohammadpoor-Baltork & Kurosh Niknam (1999): BENZYLTRIPHENYLPHOSPHONIUM DICHROMATE AS A MILD REAGENT FOR THE OXIDATION OF ORGANIC COMPOUNDS, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 31:3, 335-341

To link to this article: <http://dx.doi.org/10.1080/00304949909458330>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

8.0Hz), 6.28 (1H, d  $J = 5.9\text{Hz}$ ), 5.90 (1H, d,  $J = 5.9\text{Hz}$ ), 4.70 (1H, s), 4.40-4.50 (1H, m), 4.10 (2H, dd,  $J = 12\text{Hz}$ ), 3.90 (1H, m), 3.70 (3H, s), 2.55 (1H, m), 2.10 (1H, m), 1.60 (4H, m), 1.40 (3H, s), 1.10 (2H, m). IR (film): 2980, 1786, 1412, 1261, 798  $\text{cm}^{-1}$ . MS( $m/z$ ): 512 ( $M^+$ , 40), 497(10), 403(25), 327(35), 342(10), 101(100).  $[\alpha]_D^{20} = -25.3^\circ$  (c 0.13, acetone).

\*\*\*\*\*


## BENZYLTRIPHENYLPHOSPHONIUM DICHROMATE AS A MILD REAGENT FOR THE OXIDATION OF ORGANIC COMPOUNDS

Submitted by Abdol Reza Hajipour<sup>††</sup>, Iraj Mohammadpoor-Baltork<sup>††</sup> and Kurosh Niknam (04/15/99)

<sup>†</sup> College of Chemistry, Isfahan University of Technology, Isfahan 84156, IRAN

<sup>††</sup> Department of Chemistry, Isfahan University, Isfahan 81744, IRAN

This paper describes the oxidation of organic compounds under non-aqueous and aprotic conditions using benzyltriphenylphosphonium dichromate (**1**,  $\text{PhCH}_2\text{PPh}_3)_2 \text{Cr}_2\text{O}_7$ ) which is very easily prepared by mixing an aqueous solution of benzyltriphenylphosphonium chloride with  $\text{CrO}_3$  in 3 N HCl at room temperature. This reagent, a stable orange powder which may be stored for month without loss of activity, is soluble in acetonitrile, chloroform and dichloromethane and slightly soluble in carbon tetrachloride, ether and hexane. The oxidation of organic compounds with **1** proceeds well in acetonitrile reflux. Benzylic and allylic alcohols **2** are oxidized to the corresponding carbonyl compounds in high yields; benzoin was converted to benzil in excellent yield (Table 1). In contrast, the oxidation of allylic alcohols with manganese dioxide require a large excess of this reagent and long reaction times.<sup>1</sup> Because of the low reactivity of aliphatic alcohols, only benzylic and allylic alcohols could be converted into the corresponding carbonyl compounds.



We also found that the oxidation of **1** with oximes (**4**) and substituted hydrazones (**5**) previously accomplished by a number of reagents,<sup>2,3,5</sup> in refluxing acetonitrile gave the corresponding carbonyl compounds (Scheme 1). No further oxidation to the carboxylic acids was observed (Tables 2 and 3). The mechanism of the product reaction is not readily apparent at this time.

A noteworthy advantage of this reagent lies in its ability to selectively oxidize oximes in the presence of other oxidizable functions such as alcohols and double bonds. When we retreated an equimolar amount of oxime (**4h** or **4l**) was treated with **1** in the presence of benzyl alcohol, the oxime was




Downloaded by [University of Illinois Chicago] at 06:58 18 April 2013

Downloaded by [University of Illinois Chicago] at 06:58 18 April 2013




Downloaded by [University of Illinois Chicago] at 06:58 18 April 2013

Downloaded by [University of Illinois Chicago] at 06:58 18 April 2013

selectively oxidized (Eq. 2); the hydroxyl group of  $\alpha,\beta$ -unsaturated alcohols and the C=NOH group



$\alpha,\beta$ -unsaturated oximes were oxidized to the corresponding carbonyl compounds; the double bonds remained intact (Table 2, oxime **3d** and Table 1, alcohols **2r-2u**). In order to evaluate the selectivity of reagent **1**, the competitive reactions shown in Eqs. 2-5 were carried out. In the presence of an



**Table 2.** Conversion of **4** to Carbonyl Compounds **3**

Starting Material	Product <sup>a</sup>	Reaction Time (min)	Yield <sup>b</sup> (%)	mp. °C or bp. °C/torr (lit. <sup>2,3,5</sup> )
<b>4a</b>	<b>3a</b>	15	95	127-130/760 (127-130)
<b>4b</b>	<b>3b</b>	15	95	154-156/760 (154-156)
<b>4c</b>	<b>3c</b>	15	98	177/760 (179)
<b>4d</b>	<b>3d</b>	20	97	240/760 (240)
<b>4e</b>	<b>3e</b>	15	99	42-44 (41-44)
<b>4f</b>	<b>3f</b>	15	100	204/760 (203)
<b>4g</b>	<b>3g</b>	15	100	49-52 (50-52)
<b>4h</b>	<b>3h</b>	15	97	55-57 (55-57)
<b>4i</b>	<b>3i</b>	30	95	58-60 (58-60)
<b>4j</b>	<b>3j</b>	15	98	80-83 (80-83)
<b>4k</b>	<b>3k</b>	20	95	232/760 (234)
<b>4l</b>	<b>3l</b>	15	98	117-119 (117-118)
<b>4m</b>	<b>3m</b>	25	94	126/760 (125-128)

a) Confirmed by comparison with authentic sample (IR, TLC and NMR). b) Yield of isolated pure product after chromatography or distillation.

equimolar amount of acetophenone oxime or of benzyl alcohol only 2-mercaptopyridine was selectively oxidized (Eqs. 3 and 4). Treatment of benzyl alcohol with **1** in the presence of thioanisole, led to exclusive oxidation of benzyl alcohol (Eq. 5).

In conclusion, we report here an efficient, rapid and inexpensive method for the conversion of oximes, hydrazones, semicarbazones, alcohols and thiols to the corresponding carbonyl compounds and disulfides which is superior to previously reported methods<sup>1-38</sup> in terms of selectivity, high yields, purity of products and facile work-up.

**Table 3.** Conversion of **5** to Carbonyl Compounds **3**<sup>a</sup>

Starting Material	Product <sup>a</sup>	Reaction Time (min)	Yield <sup>b</sup> (%)	mp. °C or bp. °C/torr (lit. <sup>2,3,5</sup> )
<b>5a</b>	<b>3a</b>	30	95	48 (47-49)
<b>5b</b>	<b>3b</b>	15	98	82-85/760 (83-85)
<b>5c</b>	<b>3c</b>	30	95	48 (47-50)
<b>5d</b>	<b>3d</b>	25	98	118 (117-119)
<b>5e</b>	<b>3e</b>	30	96	42 (41-43)
<b>5f</b>	<b>3f</b>	25	96	78/760 (76-79)
<b>5g</b>	<b>3g</b>	25	96	38 (37-39)
<b>5h</b>	<b>3h</b>	25	91	244-249/760 (245-248)
<b>5i</b>	<b>3i</b>	30	85	49 (47-49)
<b>5j</b>	<b>3j</b>	30	85	120/760 (118-121)
<b>5k</b>	<b>3k</b>	40	90	232/760 (232)
<b>5l</b>	<b>3l</b>	40	85	118-120 (117-119)
<b>5m</b>	<b>3m</b>	20	91	48 (47-49)
<b>5n</b>	<b>3n</b>	20	95	84/760 (83-85)
<b>5o</b>	<b>3o</b>	25	100	48-49 (47-50)
<b>5p</b>	<b>3p</b>	35	98	37-39 (37-39)
<b>5q</b>	<b>3q</b>	40	90	178-190/760 (177-179)
<b>5r</b>	<b>3r</b>	40	80	84-85 (83-85)
<b>5s</b>	<b>3s</b>	40	85	38 (37-39)
<b>5t</b>	<b>3t</b>	40	70	47-49 (47-50)

a) Confirmed by comparison with authentic sample (IR, TLC and NMR). b) Yield of isolated pure product after chromatography or distillation.

## EXPERIMENTAL SECTION

All yields refer to isolated products after purification. Products were characterized by comparison with authentic samples (IR and NMR spectrum, thin layer chromatography, melting and boiling point).<sup>2-5</sup> All reactions were carried out in acetonitrile; all <sup>1</sup>H NMR spectra were recorded at 90 and 250 MHz in CDCl<sub>3</sub> and CCl<sub>4</sub> relative to TMS (0.00 ppm). Elemental analysis was performed by the Research Institute of Petroleum Industry, Tehran, I. R. Iran.

**Preparation of Benzyltriphenylphosphonium Dichromate (1).**- To an aqueous solution of benzyltriphenylphosphonium chloride (8.55 g, 22 mmol, 75 mL H<sub>2</sub>O), was added a solution of chromium (VI) oxide (11 g, 11 mmol) in HCl 3 N (220 mL). The reaction mixture was stirred at room temperature for 15 min. The resulting orange solid product was collected, washed with water (20 mL) and dried in a desiccator under vacuum over calcium chloride, to yield 9.54 g (94%) of orange solid product, mp. 210-212°. <sup>1</sup>H NMR: δ 7.93-6.87 (m, 20 H), 4.7 (d, *J* = 25.6 Hz, CH<sub>2</sub>-P). <sup>13</sup>C NMR: δ 133.50, 133.20, 130.20, 129.60, 129.40, 128.10, 127.70, 127.2, 117.30 (d, *J* = 85.5 Hz, P-CH<sub>2</sub>). IR (KBr): 1298, 1269, 1098, 1060, 700, 658, 590, 546 cm<sup>-1</sup>.

*Anal.* Calcd for C<sub>50</sub>H<sub>44</sub>Cr<sub>2</sub>O<sub>7</sub>: C, 69.70; H, 5.15; Cr, 12.08. Found; C, 69.60; H, 5.02; Cr, 11.95

**Oxidation of 2, 4 and 5 to 3. General Procedure.**- The alcohol 2 (oxime 4, hydrazone or semicarbazone 5) (1 mmol) was added to a stirred solution of the oxidant 1 (1 mmol, 0.92 g) in acetonitrile (10 mL). The mixture was heated at reflux until TLC showed complete disappearance of starting material, which required 5-40 min depending on substrate (Tables 1-3). The mixture was cooled and 2 g of silica gel was added to the reaction mixture. It was stirred for 5 min. The solid was then separated by suction filtration through Celite and washed with acetonitrile (2x10 mL). Evaporation of the solvent gave the carbonyl compounds 3. The products were purified by short-path distillation or column chromatography on silica gel using a mixture of ethyl acetate and hexane as eluent (90:10).

**Competitive Oxidation. Typical Procedure.**- A mixture of benzyl alcohol (1 mmol, 0.11 g) and acetophenone oxime (1 mmol, 0.14 g) was added to a stirred solution of the oxidant 1 (1 mmol, 0.92 g) in acetonitrile (20 mL). The mixture was heated at reflux until TLC showed complete disappearance of acetophenone oxime (15 min). The other competitive reactions for Eqs. 2-5 are the same as above.

**Acknowledgement.**- The authors are thankful of the Isfahan University of Technology, I. R. Iran for financial support.

## REFERENCES

1. H. O. House, "Modern Synthetic Reactions," 2nd ed. Benjamin press, New York (1972), pp 265-267.
2. H. Firouzabadi, A. R. Sardarian and H. Badparva, *Bull. Chem. Soc. Jpn*, **69**, 685 (1996).
3. a) I. Mohammadpoor-Baltork, A. R. Hajipour and H. Mohammadi, *ibid.*, **71**, 1649 (1998); b) A. R. Hajipour and N. Mahboobkhah, *Synth. Commun.*, **28**, 3143 (1998).
4. A. R. Hajipour and N. Mahboobkhah, *Indian J. Chem. Sect. B.*, **378**, 285 (1998).
5. A. R. Hajipour and N. Mahboubghah, *J. Chem. Research (S)*, 123 (1998).
6. L. F. Fieser and M. Fieser, *Reagents for Organic Synthesis*, Vol 1, Wiley, New York, 1967.
7. H. Firouzabadi, A. R. Sardarian, M. Nader and B. Vesal, *Tetrahedron*, **23**, 5001 (1984).

8. H. Firuozabadi and I. M. Baltork, *Bull. Chem. Soc. Jpn*, **65**, 1185 (1992).
9. H. Firuozabadi and Z. Mostavipoor, *ibid.*, **56**, 914 (1983).
10. H. Firuozabadi and Sardarian, *Synth. Commun.*, **13**, 863 (1983).
11. A. Clericl and O. Pona, *J. Chem. Soc., Perkin Trans. 2*, 1234, (1988).
12. W. E. Fristad and J. R. Peterson, *Tetrahedron Lett.*, 4547 (1983).
13. P. T. Perumal, *Synth. Commun.*, **20**, 1353 (1990).
14. T. Itahara, Y. Fujii and M. Toda, *J. Org. Chem.*, **53**, 3421 (1988).
15. T. Itahara, R. Ebihara, Y. Fujii and M. Toda, *Chemistry Lett.*, 1319 (1986).
16. S. Yamasaki and Y. Yamasaki, *Bull. Chem. Soc. Jpn*, **63**, 301 (1990).
17. S. Yamasaki, *Chemistry Lett.*, 823 (1992).
18. S. Yamasaki and Y. Yamasaki, *ibid.*, 571 (1990).
19. J. C. Juung, H. C. Choi and Y. H. Kim, *Tetrahedron Lett.*, **34**, 3581 (1993).
20. H. C. Choi and Y. H. Kim, *Synth. Commun.*, **24**, 2307 (1994).
21. F. Chen, J. Yang, H. Zhang, C. Guan and J. Wan, *ibid.*, **25**, 3163 (1995).
22. H. C. Choi, K. Cho, and Y. H. Kim, *Synlett*, 207 (1995).
23. F. Chen, J. Wan, C. Guan, J. Yang and H. Zhang, *Synth. Commun.*, **26**, 253 (1996).
24. I. Mohammadpoor-Baltork and Sh. Pouranshirvani, *ibid.*, **26**, 1, (1996).
25. I. Mohammadpoor-Baltork and Sh. Pouranshirvani, *Synthesis*, 756 (1997).
26. I. Mohammadpoor-Baltork, M. M. Sadeghi, N. Mahmoodi and B. Kharamesh, *Indian J. Chem. Sect. B*, **36B**, 438 (1997).
27. R. S. Varma and H. M. Meshram, *Tetrahedron Lett.*, **38**, 7973 (1997).
28. R. S. Varma, R. K. Saini and R. Dahia, *ibid.*, **38**, 7823 (1997).
29. R. S. Varma, R. Dahia and R. K. Saini, *ibid.*, **38**, 7029 (1997).
30. R. S. Varma, R. K. Saini and H. M. Meshram, *ibid.*, **38**, 6525 (1997).
31. A. R. Hajipour and N. Mahboobkhah, *Indian J. Chem. Sect. B*, **37B**, 1069 (1998).

32. E. B. Hersberg, *J. Org. Chem.*, **12**, 541 (1948).
33. J. R. Maloney and R. E. Lyle, *Synthesis*, 212 (1978).
34. L. Singh and R. N. Ram, *Synth. Commun.*, **23**, 3139 (1962).
35. J. Drabowicz, *Synthesis*, 125 (1980).
36. C. G. Rao, A. S. Radhakrishna, B. B. Singh and S. P. Bhatnagar, *ibid.*, 805 (1983).
37. S. Satish and N. Kalyanam, *Chem. Ind. (London)*, 809 (1981).
38. S. H. Pines, J. M. Chemerda and M. A. Kozlowski, *J. Org. Chem.*, **31**, 3446 (1966).

\*\*\*\*\*

## PREPARATION AND CHARACTERIZATION OF NEW SUBSTITUTED

### 5- METHOXY-2-STYRYL-4-PYRONES

Submitted by  
(12/24/98)

Zdenka Stiplošek, Marija Šindler-Kulyk and Krešimir Jakopčić\*

*Department of Organic Chemistry*  
Faculty of Chemical Engineering and Technology  
University of Zagreb  
Marulicev trg 20, 10000 Zagreb, CROATIA

During the last several decades, many 4-pyrones or compounds containing 4-pyrone moieties has been found to be biologically active; bactericidal, insecticidal, herbicidal, fungicidal, antiallergenic, cytotoxic and potential anticancer activity has been reported.<sup>1</sup> Some 4-pyrones with the styryl group possess anticancer activity<sup>2</sup> and 5-hydroxy-2-styryl-4-pyrone has been used in the formulation of skin-lightening cosmetics,<sup>3</sup> and the use of such pyrones in the synthesis of polycondensed heterocyclics has been described.<sup>4</sup> Previous papers of this series described some reactions of 5-hydroxy-4-pyrones.<sup>5</sup> The transformation to corresponding *N*-substituted-5-hydroxy-4-pyridones (useful as chelating agents),<sup>5a-d</sup> photochemical isomerizations<sup>5e</sup> and ring-contraction reactions<sup>5f</sup> have been studied. Our continuing interest in the photochemistry of 4-pyrones, especially in regard to the difference between reactions of 5-hydroxy and its methylated analogues, prompted us to study styryl-substituted 4-pyrones.

Herewith we report the synthesis of several aryl-substituted 5-methoxy-2-ethenyl-4-pyrones (**4a-4f**) presumably capable of exhibiting various photochemical reactions. Several 5-hydroxy-2-