## Synthesis of the Ribosomal P-Site Substrate CCA-pcb

## Minghong Zhong<sup>†</sup> and Scott A. Strobel<sup>\*,†,‡</sup>

Department of Molecular Biophysics and Biochemistry, and Department of Chemistry, Yale University, New Haven, Connecticut 06520-8114

strobel@csb.yale.edu

Received October 13, 2005

## ORGANIC LETTERS 2006 Vol. 8, No. 1 55-58

## ABSTRACT



CCA-pcb (cytidylyl-(3'5')-cytidylyl-(3'5')-3'(2')-O-(N-(6-D-(+)-biotinoylaminohexanoyl)-L-phenylalanyl)adenosine), a ribosomal P-site substrate, was synthesized by phosphoramidite chemistry in 26 steps with an overall yield of 18%, starting from biotin. The synthesis relies on the judicious selection of orthogonal silyl protecting groups for the 5'-hydroxyls and acid-labile protecting groups (DMTr, AcE, and MeE) at other reactive sites to ensure the intactness of the labile ester. Both 3'-esterification and nucleotide coupling were accomplished by in situ activation with imidazolium ions.

The catalytic mechanism of peptide bond formation by the ribosome is an area of ongoing research. Proposed catalytic strategies include general acid—base catalysis,<sup>1</sup> substrate-assisted catalysis,<sup>2</sup> and catalysis derived solely from substrate alignment.<sup>3</sup> It is also possible that the reaction may follow a pathway different from the stepwise mechanism involving a tetrahedral intermediate that is observed for aminolysis reactions in aqueous solution.<sup>4</sup> Improved chemical and enzymatic tools are needed to differentiate between these mechanistic possibilities.

In addition to the reaction with full-sized tRNAs, the ribosome can also catalyze peptide bond formation between minimal A-site and P-site substrates. We previously reported a modified fragment reaction that used the A-site substrate, C-puromycin, and a P-site substrate with an extended peptidyl-like chain, CCA-pcb.<sup>5</sup> These substrates have played a valuable role in the structural and biochemical characterization of the peptidyl transfer (PT) reaction.<sup>5,6</sup> Studies have included the use of N<sup>15</sup>-labeled C-puromycin for kinetic isotope effect (KIE) analysis,<sup>6e</sup> a method that can be used to characterize the transition-state structure of a chemical reaction. Our goal is to determine intrinsic KIEs at several

<sup>&</sup>lt;sup>†</sup> Department of Molecular Biophysics and Biochemistry.

<sup>&</sup>lt;sup>‡</sup> Department of Chemistry.

 <sup>(1) (</sup>a) Muth, G. W.; Ortoleva-Donnelly, L.; Strobel, S. A. Science 2000, 289, 947–950.
 (b) Nissen, P.; Hansen, J.; Ban, N.; Moore, P. B.; Steiz, T. A. Science 2000, 289, 920–930.

<sup>(2) (</sup>a) Dorner, S.; Panuschka, C.; Schmid, W.; Barta, A. *Nucleic Acids Res.* **2003**, *31*, 6536–6541. (b) Weinger, J. S.; Parnell, K. M.; Dorner, S.; Green, R.; Strobel, S. A. *Nat. Struct. Mol. Biol.* **2004**, *11*, 1101–1106.

<sup>(3)</sup> Sievers, A.; Beringer, M.; Rodina, M. V.; Wolfenden, R. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 7897–7901.

 <sup>(4) (</sup>a) Blackburn, G. M.; Jencks, W. P. J. Am. Chem. Soc. 1968, 90, 2638–2645. (b) Satterthwait, A. C.; Jencks, W. P. J. Am. Chem. Soc. 1974, 96, 7018–7031.

<sup>(5)</sup> Schmeing, T. M.; Seila, A. C.; Hansen, J. L.; Freeborn, B. S.; Juliane, K.; Scaringe, S. A.; Strobel, S. A.; Moore, P. B.; Steitz, T. A. *Nat. Struct. Biol.* **2002**, *9*, 225–230.

<sup>(6) (</sup>a) Bobkova, E. V.; Yan, Y. P.; Jordan, D. B.; Kurilla, M. G.; Pompliano, D. L. J. Biol. Chem. **2003**, 278, 9802–9807 and references therein. (b) Okuda, K.; Seila, A. C.; Strobel, S. A. Biochemistry **2005**, 44, 6675–6684. (c) Schmeing, T. M.; Huang, K. S.; Strobel, S. A.; Steitz, T. A. Nat. In press. (d) Hansen, J. L.; Schmeing, T. M.; Moore, P. B.; Steitz, T. A. Proc. Natl. Acad. Sci. U.S.A. **2004**, 99, 11670–11675. (e) Seila, A. C.; Okuda, K.; Strobel, S. A. Biochemistry **2005**, 44, 4018–4027.

atomic positions in both the A-site and P-site substrates in order to explore the transition state of the PT reaction. This necessitates that isotopic substitutions be introduced into CCA-pcb, but an efficient chemical synthesis of CCA-pcb has not been reported.

The synthesis of CCA-pcb offers a synthetic challenge because the ester linkage of aminoacylribonucleotide derivatives is an activated, high energy bond with a free energy of hydrolysis comparable to that of ATP.<sup>7</sup> Fast and reversible migration of the amino acyl occurs between the 2'- and 3'hydroxyl of the ribofuranose. This results in a mixture of 2'- and 3'-O-aminoacylated isomers. Most recent syntheses of 2'/3'-O-aminoacyloligonucleotides utilized the cyanomethyl ester of protected amino acids.<sup>8</sup> The only synthetic procedure reported for CCA-pcb was by direct coupling between CpCpA and the cyanomethyl ester of pcb, which is expected to provide low yields due to a mixture of mono-, di-, and multiple aminoacylated byproducts.<sup>6a</sup> Here we report an alternative route that provides an efficient synthesis of CCA-pcb.

It was observed that the absence of a neighboring hydroxyl stabilizes the ester bond of 2'/3'-aminoacylnucleotides against hydrolysis and that the hydrolysis rate decreases dramatically at low pH.<sup>9</sup> A half-life of 250 h at pH 2.5 was observed for a 2'/3'-O-L-phenylalanyladenosine methyl phosphate derivative.<sup>10</sup> Therefore, we reasoned that the 2'-OH and the amino groups should be protected by acid-labile protecting groups such as ketal or ortho ester and trityl group, respectively. In our synthesis, CCA-pcb was prepared in two different schemes by phosphoramidite chemistry using acid-labile 2'-bisacetoxyethoxymethyl (AcE), 2'-bismethoxyethoxymethyl (MeE), and 4/6-(4,4'-dimethoxytrityl) (DMTr) as protecting groups.

Synthesis of A-pcb Fragment 19. Biotin 1 is insoluble in most organic solvents. Its solubility was enhanced by 1-Ntritylation. Esterification of biotin by treatment with SOCl<sub>2</sub> in methanol and tritylation with 4,4'-dimethoxytrityl chloride (DMTrCl) in pyridine in the presence of Et<sub>3</sub>N,<sup>11</sup> followed by hydrolysis of methyl ester in 1 M NaOH/MeOH, gave the tritylated biotin 2 (52%). This biotin derivative is highly soluble even in DCM. DCC-mediated coupling between compound 2 and methyl 6-aminohexanoate in pyridine and hydrolysis of the methyl ester in 1 M NaOH/MeOH produced compound 4 (77%) (Scheme 1). Significant loss of 2 during purification was observed, and a crude separation followed by amide formation gave 3 in much higher yield (83%).

Adenosine **5** was converted into 3',5'-O-(tetraisopropyldisiloxane-1,3-diyl)adenosine **6** (93%) via reaction with 1,3dichloro-1,1,3,3-tetraisopropyl-disiloxane in pyridine. The



6-amino group of adenosine was protected with DMTr by reaction with DMTrCl to give quantitatively compound **7**.<sup>12</sup> MeE was introduced into the 2'-position by reflux of compound **7**, with trismethoxyethoxy orthoformate, 4-*tert*-butyldimethyl-siloxy-3-penten-2-one, and PPTS in DCM. The derived syrup after chromatography was treated with HF–TEMED/CH<sub>3</sub>CN to provide compound **9** (71%).<sup>13</sup> Compound **9** was further protected by a sterically hindered silyl group at the 5'-position by treatment with bis(trimethylsiloxy)cyclododecyloxylsilyl (DOD) chloride and imidazole in cold THF to give **10** (64%) (Scheme 2). Compound **10** is



unstable, and significant migration of the 2'-O-MeE to the 3'-position occurred during storage; thus, it was freshly prepared and purified prior to the subsequent acylation reaction.

<sup>(7)</sup> Preiss, J.; Berg, P.; Ofengand, E. J.; Bergman, F. H.; Dieckmann, M. Proc. Natl. Acad. Sci. U.S.A. **1959**, 45, 319–328.

<sup>(8)</sup> Robertson, S. A.; Ellamn, J. A.; Schultz, P. J. Am. Chem. Soc. 1991, 113, 2722–2729.

<sup>(9) (</sup>a) Chlàdek, S.; Spinzl, M. Angew. Chem., Int. Ed. Engl. 1985, 24, 371–391. (b) Chlàdek, S. In Chemistry of Nucleosides and Nucleotides; Townsend, L. B., Ed.; Plenum Press: New York, 1994.

<sup>(10)</sup> Stutz, A.; Hobartner, C.; Pitsch, S. Helv. Chem. Acta 2000, 83, 2477-2503.

<sup>(11)</sup> Alves, A. M.; Holland, D.; Edge, M. D. Tetrahedron Lett. 1989, 30, 3089–3092.

<sup>(12)</sup> Kawana, M.; Takeuchi, K.; Ohba, T.; Kuzuhara, H. Bull. Chem. Soc. Jpn. **1988**, 61, 2437–2442.

Aminoacylation of compound 10 was first attempted via either the cyanomethyl ester<sup>8</sup> or the acyl fluoride<sup>14</sup> of N-protected L-phenylalanine. Both gave low to moderate yields probably due to significant steric hindrance caused by DOD and MeE. Aminoacylation was then performed with a stronger promoter. Compound 10 was treated with N-FMOC-L-phenylalanine in the presence of mesitylenesulfonyl tetrazole (MST)<sup>15,16</sup> and excess N-methylimidazole in DCM at ambient temperature to produce quantitatively the fully protected derivative 16. The reaction involves the imidazolium cation, as shown by <sup>1</sup>H NMR, which was nucleophilically displaced as a neutral leaving group by the 3'-hydroxyl of 10. Removal of FMOC by treatment of 16 with 20% piperidine in CH<sub>3</sub>CN resulted in 17 (97%). DCC mediated amide formation with 4 (99%), followed by desilylation by treatment with HF-TEMED in CH<sub>3</sub>CN, gave 19 in a yield of 84% (Schemes 2 and 3).



Synthesis of C(3'5')C Phosphoramidite. Cytidine was treated with 1,3-dichloro-1,1,3,3-tetraisopropyldisiloxane in pyridine to afford 3',5'-O-(tetraisopropyldisiloxane-1,3-diyl)cytidine (92%). Compound 12 was selectively tritylated at the N4 to produce 13 (91%), which was further protected at the 2'-position by reflux with a mixture of trisacetoxyethoxy orthoformate, 4-*tert*-butyldimethylsiloxy-3-penten-2-one, and PPTS in DCM. The syrup that resulted after chromatography was treated with HF–TEMED in CH<sub>3</sub>CN to provide compound **14** (68%), which was further protected by DOD (89%) (Scheme 2). The cytidylyl phosphoramidite **20** was obtained by treatment of compound **15** with methyl tetra-isopropylphosphorodiamidite (POMe) in the presence of tetrazole (Tet) in DCM overnight (81%). Compound **20** was a mixture of two diastereomers due to the chirality of the phosphoramidite as shown by <sup>1</sup>H NMR and <sup>31</sup>P NMR.

Compound **20** was directly coupled to compound **14** in the presence of PhImOTf<sup>17</sup> and molecular sieves in CH<sub>3</sub>-CN/DCM (1:1), followed by oxidation with *t*-BuOOH in toluene. The reaction selectively gave **21** in good yield (81%) after chromatography. The diribonucleotide was transformed into its phosphoramidite **22** as before (73%) (Scheme 4).



Compound **22** is a mixture of four diastereomers due to the chirality of phosphorus triester and phosphoramidite. Synthesis of the dinucleotide with tetrazole and S-Et-tetrazole only gave moderate yields, which might be caused by the bulky silyl protecting group.

Synthesis of Protected CCA-pcb and Deprotection. Diribonucleotide phosphoramidite 22 was coupled to 1 molar equiv of compound 19 using PhImOTf and molecular sieves as the promotor in CH<sub>3</sub>CN/DCM (1:1) to form the fully protected CCA-pcb 23 in good yield (72%). Reactions with tetrazole and S-Et-tetrazole gave only traces of products. The success of the coupling reaction involves imidazolium cation, which was displaced by the 5'-OH to release a neutral leaving group.<sup>17</sup> Desilylation of compound 23 by treatment with HF– TEMED in CH<sub>3</sub>CN (73%) and cleavage of methyl phosphate ester with S<sub>2</sub>Na<sub>2</sub> (88%), followed by complete removal of DMTr and orthoformate ester in 0.5 M HCOOH in MeOH–

<sup>(14)</sup> Oliver, J. S.; Oyelere, A. J. Org. Chem. 1996, 61, 4168–4171.
(15) Stawinski, J.; Hozumi, T.; Narang, S. A.; Bahl, C. P.; Wu, R. Nucleic Acids Res. 1977, 4, 353–371.

<sup>(16) (</sup>a) Kumar, G.; Celewicz, L.; Chlàdek, S. J. Org. Chem. **1982**, 47, 634–644. (b) Happ, E.; Scalfi-Happ, C.; Chlàdek, S. J. Org. Chem. **1987**, 52, 5387–5391. (c) Hagen, M.; Chlàdek, S. J. Org. Chem. **1989**, 54, 3189–3195. (d) Hagen, M.; Scalfi-Happ, C.; Happ, E.; Chlàdek, S. J. Org. Chem. **1988**, 53, 5040–5045.

<sup>(17) (</sup>a) Kawai, R.; Sugimoto, J.; Kataoka, M.; Hayakawa, Y. *Nucleosides Nucleotides Nucleic Acids* **2001**, *20*, 1047–1048. (b) Hayakawa, Y.; Kawai, R.; Hirata, A.; Sugimoto, J.; Kataoka, M.; Sakakura, A.; Hirose, M.; Noyori, R. J. Am. Chem. Soc. **2001**, *123*, 8165–8176.



DCM (1:1) at 55 °C for 5 h gave compound **24** (80%) as a mixture of two regiosiomers (2'/3'-ester) (Scheme 5). The overall yield from biotin was 18%.

Compound 23 was also synthesized by sequential addition of cytidylyl phosphoramidite 20 to A-pcb derivative 19 (Scheme 6). Coupling of 19 with 20 in the presence of PhImOTf and molecular sieves gave fully protected compound CpA-pcb, 25 (80%). Compound 25 was then desilylated (60%) and coupled to 20 a second time to afford a white foam (72%), which was deprotected as described above to produce a compound spectroscopically identical to compound 24 in an overall yield of 9%.

In conclusion, we have accomplished the first efficient synthesis of CCA-pcb, which relies on the judicious selection



of orthogonal silyl protecting groups for 5'-hydroxyls and acid-labile protecting groups (DMTr, ACE, and MeE) at other reactive sites to ensure the intactness of the labile ester. Both 3'-esterification and nucleotide coupling were accomplished by in-situ activation with imidazolium ions. This efficient synthesis will make it possible to prepare a series of isotopically substituted P-site substrates for KIE analysis of the PT reaction.

Acknowledgment. We thank Stephen Scaringe and Amy Seila for helpful suggestions. This research was supported by an American Cancer Society Beginning Investigator Grant to S.A.S.

**Supporting Information Available:** Experimental procedures and spectroscopic data. This material is available free of charge via the Internet at http://pubs.acs.org.

OL052484F