

Short Research Article

The syntheses of (^{14}C) and ($^{13}\text{C}_4$)pyromellitic acid[†]

NICK SHIPLEY and KENNETH W. M. LAWRIE*


GlaxoSmithKline, Isotope Chemistry, Medicines Research Centre, Gunnels Wood Road, Stevenage SG1 2NY, UK

Received 17 August 2006; Revised 2 November 2006; Accepted 22 November 2006

Keywords: carbon-14; carbon-13; cyanation; oxidation

Introduction

In the drug development process appropriate salt selection can be critical. For a recent project a pyromellitic acid (1,2,4,5-benzene tetracarboxylic acid), **1**, salt proved optimal.



We required carbon-14 and stable labelled versions of pyromellitic acid: the radio-isotopomer at 50–60 mCi/mmol for traditional ADME work and the stable label as an internal standard in a LC/MS/MS assay. In general an effective internal standard requires an increase of at least 3 a.m.u. over the test substance and a very high degree of isotopic incorporation (typically <0.1% unlabelled). We chose to fulfil these criteria by mass labelling each of the carboxyl groups with carbon-13.

Results and discussion

1,2,4,5-Tetracyanobenzene is readily saponified to pyromellitic acid. We therefore chose this as our initial


target. We envisaged that a route common to both isotopomers could be developed by cyanation of tetrabromo or iodo benzene with either ^{13}CN or ^{14}CN . However, under a variety of conditions (CuCN , KCN / CuI , $\text{Zn}(\text{CN})_2/\text{Pd}(\text{Ph}_3\text{P})_4$) no useful product was isolated. An alternative related approach suggested itself at this point. Oxidation of aryl methyl groups to carboxylic acids is a well known and straightforward process¹. We found that both 1,2,4,5-tetramethyl benzene and 2,4,5-trimethylbenzoic acid could be oxidized by alkaline permanganate to pyromellitic acid in good yield (70–80%). This discovery proved crucial.

Cyanation of 2,4,5-trimethyl bromobenzene with $\text{Cu}[^{14}\text{C}]\text{CN}$ in NMP at 175°C followed by saponification of the crude $[^{14}\text{C}]\text{nitrile}$ (NaOH , EtOH , reflux) gave 2,4,5-trimethyl[carboxyl- ^{14}C] benzoic acid in 84% yield from copper cyanide. Oxidation by slow addition of aliquots of 5% w/v KMnO_4 in water to a solution of the radiolabelled acid in 1.5M KOH at 90°C completed the synthesis. The crude product was triturated with ether/hexane and crystallized from hot water (60% yield, radiochemical purity >98%, specific activity 53 mCi/mmol).

Our initial plan was to prepare $[^{13}\text{C}_4]\text{tetramethylbenzene}$ by exhaustive metalation and methylation of tetrabromobenzene and subsequent oxidation. We anticipated that a large excess of MeI would be required, but $[^{13}\text{C}]\text{MeI}$ is inexpensive and readily available so this was not a major concern. Unfortunately, we were unable to drive the reaction to completion and isolated $[methyl-^{13}\text{C}_3]2,4,5\text{-trimethylbromobenzene}$ as the major product. However, we had already demonstrated in the carbon-14 synthesis that this could be readily converted to pyromellitic acid by a cyanation, saponification and oxidation sequence. Simply substituting $\text{Cu}[^{13}\text{C}]\text{CN}$ for $\text{Cu}[^{14}\text{C}]\text{CN}$ gave the desired product with a chemical purity of >98% and no detectable unlabelled content by MS.

*Correspondence to: Kenneth W. M. Lawrie, GlaxoSmithKline, Isotope Chemistry, Medicines Research Centre, Gunnels Wood Road, Stevenage SG1 2NY, UK. E-mail: ken.w.lawrie@gsk.com

[†]Proceedings of the Ninth International Symposium on the Synthesis and Applications of Isotopically Labelled Compounds, Edinburgh, 16–20 July 2006.



REFERENCE

1. Bromby NG, Peters AT, Rowe M. *J Chem Soc* 1943; 144.