



Asymmetric Catalysis Hot Paper

 How to cite: *Angew. Chem. Int. Ed.* **2021**, 60, 6425–6429

International Edition: doi.org/10.1002/anie.202016771

German Edition: doi.org/10.1002/ange.202016771

Interrupted Pyridine Hydrogenation: Asymmetric Synthesis of δ -Lactams

 Tobias Wagener⁺, Lukas Lückemeier⁺, Constantin G. Daniliuc, and Frank Glorius*

 Dedicated to Professor David A. Evans on the occasion of his 80th birthday

Abstract: Metal-catalyzed hydrogenation is an effective method to transform readily available arenes into saturated motifs, however, current hydrogenation strategies are limited to the formation of C–H and N–H bonds. The stepwise addition of hydrogen yields reactive unsaturated intermediates that are rapidly reduced. In contrast, the interruption of complete hydrogenation by further functionalization of unsaturated intermediates offers great potential for increasing chemical complexity in a single reaction step. Overcoming the tenet of full reduction in arene hydrogenation has been seldom demonstrated. In this work we report the synthesis of sought-after, enantioenriched δ -lactams from oxazolidinone-substituted pyridines and water by an interrupted hydrogenation mechanism.

Metal-catalyzed hydrogenation is known to be a simple and powerful method to increase molecular complexity.^[1] In particular, the hydrogenation of easily accessible N-heteroarenes such as pyridines offers access to important saturated azacycles.^[2] This established reaction is limited solely to the formation of new C–H and N–H bonds; additional synthetic manipulations are required to introduce further chemical functionality.^[3] The stepwise transfer of molecular hydrogen in arene hydrogenation yields intermediates with double bonds remaining, which in principle offer the possibility of further functionalization (Figure 1).^[4] Nevertheless, harnessing unsaturated intermediates in arene hydrogenation for other C–X bond forming events is unknown in the literature, although the partial hydrogenation of certain substrates like benzene or phenol has been achieved.^[5] Recently, Donohoe and co-workers reported a pioneering combination of C–H and C–C bond formation in a transfer hydrogenation of pyridinium salts.^[6] Protected tetrahydropyridines were thus




Figure 1. Mechanistic pathways of unsaturated intermediates in arene hydrogenation and this work. Aux = chiral auxiliary.

synthesized using iridium-catalyzed hydride transfer followed by hydroxymethylation. This precedent underlines the huge potential of an interrupted hydrogenation strategy to rapidly build up chemical complexity, yet, further method development is necessary to elevate this approach to a general synthetic strategy. We now report a pyridine hydrogenation interrupted by nucleophilic substitution of an unsaturated intermediate. Our procedure enables the synthesis of valuable δ -lactams by Pd-catalyzed hydrogenation of oxazolidinone-substituted pyridines in the presence of water.^[7]

Lactams, such as 2-piperidones, are important building blocks and represent core motifs in many pharmaceuticals and natural products.^[8] Despite interest in this moiety, synthetic access to enantioenriched δ -lactams remains challenging.^[9] In this regard, we envision metal-catalyzed hydrogenation as a powerful strategy to address this problem. However, the synthesis of 2-piperidones via direct asymmetric hydrogenation^[10] of pyridone precursors is hampered by amide resonance and, to the best of our knowledge, has never been achieved.^[11] In contrast, our procedure starts with oxazolidinone-substituted pyridines, which are readily available from cheap and abundant 2-halogenated pyridines in a single step.^[12] The oxazolidinone is cleaved within the reaction, thus acting as a traceless chiral auxiliary which can be fully recycled.

[*] T. Wagener,^[†] L. Lückemeier,^[†] Dr. C. G. Daniliuc, Prof. Dr. F. Glorius
 Organisch-Chemisches Institut
 Westfälische Wilhelms-Universität Münster
 Corrensstraße 40, 48149 Münster (Germany)
 E-mail: glorius@uni-muenster.de

[†] These authors contributed equally to this work.

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202016771>.

© 2021 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.


























After our initial discovery, we studied the influence of the reaction parameters in more detail (see Supporting Information for details). The presence of strong Brønsted acids like hydrochloric acid is essential.^[13] Using weak Brønsted acids like formic acid resulted in the exclusive formation of piperidine product (Table S1, Entry 7). Furthermore, different solvents were tested, with a combination of THF and H₂O giving the best result (Table S1, Entries 1–6). The nature of the catalyst is also decisive for the reaction outcome. While various Pd catalysts generally gave the desired δ -lactam in high yields and enantioselectivities, other catalysts such as PtO₂, Ru/C or Rh/C resulted in little or no yield (Table S2, Entries 2–9). Various oxazolidinones from the chiral pool provided the product in a high enantiomeric ratio while a benzyl substituent gave the best result (Table S2, Entries 10–12).^[14] Lowering the hydrogen pressure to 25 bar results in a decrease of the enantiomeric ratio (Table S2, Entry 13). To investigate the reproducibility of our new method, we performed a reaction-condition-based sensitivity screen (Table 1; see Supporting Information for details).^[15]

With optimized conditions in hand, we investigated the reaction scope of this new transformation. A series of alkylated δ -lactams (**1–4**) was synthesized in high yields and excellent enantiomeric ratios. Furthermore, our method was capable of forming multiple stereocenters in a highly selective fashion. Dimethylated lactam **5** was isolated in 66% yield

with a 98:2 enantiomeric ratio and only one diastereomer observed. Additional functional groups such as basic azacycles (**7**) or alkyl ethers (**8**) were tolerated without significant decrease in yield or enantiomeric ratio. Although hydrogenation of fluorinated arenes is challenging,^[16] several fluorinated δ -lactams (**9–13**) were prepared containing CF, CF₃, and CF₂H groups in moderate to good yields. Interestingly, a chemoselective hydrogenation of the pyridine core over a phenyl substituent was observed for our catalytic system to give various substituted phenyl-piperidones (**14–17**). Moreover, δ -lactams carrying important carbonyl functionalities such as esters and amides (**18–23**) were prepared in moderate to high yields and high enantiomeric ratios. Disubstituted δ -lactams **24** and **25** were synthesized highly selectively with only one diastereomer observed in 97:3 and 89:11 enantiomeric ratios, respectively. Substitution in the pyridine 3-position however decreased product formation and resulted in an almost racemic mixture (**6**, see Supporting Information for details).

The absolute configurations of δ -lactams **1** and **10** were unambiguously determined by X-ray analysis and are in accordance with our previous observations (Figure 2, see Supporting Information for details).^[17,7a,b] Based on the Horvut–Polanyi mechanism^[18] for heterogeneous hydrogenation, the absolute configuration of the products with substituents in 4- and 5- position are assigned analogously (see Scheme S1).

Table 1: Reaction scope for the palladium-catalyzed synthesis of 2-piperidones. See Supporting Information for full experimental details.

			
	1 , ^[a] 89% 98:2 e.r.		2 , 90% 98:2 e.r.
	3 , ^[a] 87% 95:5 e.r.		4 , ^[b] 84% 99:1 e.r.
	5 , 66% >95:5 d.r. 98:2 e.r.		6 , ^[a] 40% 62:38 e.r.
	7 , ^[a] 65% 97:3 e.r.		8 , ^[a] 99% 95:5 e.r.
	9 , ^{[a],[c]} 64% 97:3 e.r.		10 , 32% 97:3 e.r.
	11 , ^[a] 66% 96:4 e.r.		12 , 66% 93:7 e.r.
	13 , 73% 99:1 e.r.		14 , R=H, 74%, 90:10 e.r.
	15 , R=Me, ^[a] 65%, 88:12 e.r.		16 , R=CF ₃ , ^[a] 87%, 95:5 e.r.
	17 , R=F, ^[a] 76%, 92:8 e.r.		18 , 89% 96:4 e.r.
	19 , 53% 93:7 e.r.		20 , 86% 93:7 e.r.
	21 , 62% 98:2 e.r.		22 , 83% 95:5 e.r.
	23 , ^[a] 63% 92:8 e.r.		24 , 80% >95:5 d.r. 97:3 e.r.
	25 , 30% >95:5 d.r. 89:11 e.r.		

[a] 4-Isopropylloxazolidinone as chiral auxiliary. [b] 4-*tert*-Butyloxazolidinone as chiral auxiliary. [c] Epimeric (*R*)-isopropylloxazolidinone as chiral auxiliary.





Figure 2. Absolute configuration of δ -lactams **1** and **10**.

In agreement with this model, a deuteration experiment confirms the addition of deuterium to one diastereotopic face of the pyridine exclusively (see Supporting Information for details).

Subsequently, we investigated the scalability of our procedure and the possibility of recycling the auxiliary. The hydrogenation was performed on a 2 g scale without significant loss in yield or enantiomeric ratio (Scheme 1, a). Lactam **ent-4** was obtained in 78 % yield and 97:3 enantiomeric ratio and oxazolidinone **26** was recovered in 92 % yield and >99:1 e.r. thus outlining its good recyclability.




Scheme 1. Gram-scale reaction and ring opening.


As further product application, δ -lactam **ent-4** was hydrolyzed to give the δ -amino acid methyl ester **27** in 99 % yield and 97:3 e.r. after heating in hydrochloric acid for 24 h and treatment with methanol (Scheme 1, b).

Intrigued by these observations, we investigated the reaction mechanism. To identify potential intermediates, the reaction was stopped after 2 h and the mass of an intermediate with one remaining double bond was detected by GC-MS (Scheme 2, a). Although unsaturated intermediates are very short-lived in arene hydrogenation and are rapidly further hydrogenated, imine **28** was isolated in 5 % yield (Scheme 2, b). Conversion of imine **28** in water/THF with hydrochloric acid in the absence of Pd catalyst and H_2 provided the δ -lactam product in almost quantitative yield (Scheme 2, c). Thus, imine **28** is an intermediate species in our procedure and can undergo nucleophilic substitution with water. It is worth mentioning that only imine **28** as one of six potential intermediates in this reaction is suitable to hydrolyze to yield the desired 2-piperidone (Scheme 2, a). Based on these experiments we postulate the following reaction mechanism (Scheme 2, d): By addition of a Brønsted acid, the basic nitrogen of the oxazolidinone-substituted pyridine is proton-


a) potential tetrahydropyridine intermediates




b) isolation of intermediate



c) hydrolysis of intermediate



d) proposed mechanism



Scheme 2. Mechanistic investigation and proposal.

ated and forms a stable conformer through hydrogen bonding to the oxazolidinone. Owing to the steric repulsion of the oxazolidinone substituent of this rigid conformation, a diastereoselective hydrogenation of the unhindered face of the pyridine proceeds to set the new stereocenter. After the addition of two equivalents of H_2 , oxazolidinone-substituted imine **I** is formed (see Scheme S1). In contrast to a classic arene hydrogenation, this intermediate, following our new protocol, does not undergo another catalytic reduction (see product **III**) but reacts with water in a nucleophilic substitution. In this substitution, the oxazolidinone is released and therefore serves as a recyclable, traceless auxiliary. The resulting iminium **II** tautomerizes rapidly to yield the final δ -lactam product.

In summary, we have disclosed a heteroarene hydrogenation interrupted by nucleophilic substitution. Hydrogenation of oxazolidinone-substituted pyridines gives access

to enantioenriched δ -lactams by applying oxazolidinones as traceless chiral auxiliaries. Our procedure was used to access a plethora of sought-after δ -lactams in high yields and excellent enantiomeric ratios. Harnessing unsaturated intermediates in arene hydrogenation is an underexplored research area and overcomes the inherent limitation of forming solely C–H and N–H bonds. We see huge potential in this research direction since it tremendously increases molecular complexity in a single step.

Acknowledgements

Generous financial support by the European Research Council (ERC Advanced Grant Agreement No. 788558), the Deutsche Forschungsgemeinschaft (Leibniz Award), and the Alfried Krupp von Bohlen und Halbach Foundation are gratefully acknowledged. The authors thank Cornelia Weitkamp and Lisa Grabinski for experimental support as well as Torny Dalton, Arne Heusler, Daniel Moock, Dr. J. Luca Schwarz, and Marco Wollenburg for many helpful discussions. Open access funding enabled and organized by Projekt DEAL.

Conflict of interest

The authors declare no conflict of interest.

Keywords: asymmetric catalysis · heterogeneous catalysis · hydrogenation · lactams · nitrogen heterocycles

- [1] a) M. P. Wiesenfeldt, Z. Nairoukh, T. Dalton, F. Glorius, *Angew. Chem. Int. Ed.* **2019**, 58, 10460; *Angew. Chem.* **2019**, 131, 10570; b) W. C. Wertjes, E. H. Southgate, D. Sarlah, *Chem. Soc. Rev.* **2018**, 47, 7996; c) S.-C. Qi, X.-Y. Wei, Z.-M. Zong, Y.-K. Wang, *RSC Adv.* **2013**, 3, 14219; d) F. Lovering, *MedChemComm* **2013**, 4, 515; e) F. Lovering, J. Bikker, C. Humblet, *J. Med. Chem.* **2009**, 52, 6752; f) *The Handbook of Homogeneous Hydrogenation* (Eds.: H. G. de Vries, C. J. Elsevier), Wiley-VCH, Weinheim, **2006**; g) *Handbook of Heterogeneous Catalytic Hydrogenation for Organic Synthesis* (Ed.: S. Nishimura), Wiley, New York, **2001**.
- [2] For selected examples, see: a) S. Kim, F. Loose, M. J. Bezdek, X. Wang, P. J. Chirik, *J. Am. Chem. Soc.* **2019**, 141, 17900; b) Y.-N. Duan, X. Du, Z. Cui, Y. Zeng, Y. Liu, T. Yang, J. Wen, X. Zhang, *J. Am. Chem. Soc.* **2019**, 141, 20424; c) R. Adam, J. R. Cabrero-Antonino, A. Spannenberg, K. Junge, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2017**, 56, 3216; *Angew. Chem.* **2017**, 129, 3264; d) W.-X. Huang, B. Wu, X. Gao, M.-W. Chen, B. Wang, Y.-G. Zhou, *Org. Lett.* **2015**, 17, 1640; e) Y. Liu, H. Du, *J. Am. Chem. Soc.* **2013**, 135, 12968; f) G. E. Dobereiner, A. Nova, N. D. Schley, N. Hazari, S. J. Miller, O. Eisenstein, R. H. Crabtree, *J. Am. Chem. Soc.* **2011**, 133, 7547; g) J. M. Farrell, J. A. Hatnean, D. W. Stephan, *J. Am. Chem. Soc.* **2012**, 134, 15728; h) X.-B. Wang, W. Zeng, Y.-G. Zhou, *Tetrahedron Lett.* **2008**, 49, 4922; i) C. Y. Legault, A. B. Charette, *J. Am. Chem. Soc.* **2005**, 127, 8966; j) N. Douja, M. Besson, P. Gallezot, C. Pinel, *J. Mol. Catal. A: Chem.* **2002**, 186, 145; k) L. Hegedűs, V. Hada, A. Tungler, T. Máthé, L. Szepeszy, *Appl. Catal. A* **2000**, 201, 107.
- [3] For selected examples of chemoselective hydrogenations, see: a) D. Moock, M. P. Wiesenfeldt, M. Freitag, S. Muratsugu, S. Ikemoto, R. Knitsch, J. Schneidewind, W. Baumann, A. H. Schäfer, A. Timmer, M. Tada, M. R. Hansen, F. Glorius, *ACS Catal.* **2020**, 10, 6309; b) M. Wollenburg, D. Moock, F. Glorius, *Angew. Chem. Int. Ed.* **2019**, 58, 6549; *Angew. Chem.* **2019**, 131, 6621; c) L. Ling, Y. He, X. Zhang, M. Luo, X. Zeng, *Angew. Chem. Int. Ed.* **2019**, 58, 6554; *Angew. Chem.* **2019**, 131, 6626; d) M. P. Wiesenfeldt, T. Knecht, C. Schlepphorst, F. Glorius, *Angew. Chem. Int. Ed.* **2018**, 57, 8297; *Angew. Chem.* **2018**, 130, 8429; e) B. L. Tran, J. L. Fulton, J. C. Linehan, J. A. Lercher, R. M. Bullock, *ACS Catal.* **2018**, 8, 8441; f) Y. Wei, B. Rao, X. Cong, X. Zeng, *J. Am. Chem. Soc.* **2015**, 137, 9250.
- [4] For a general combination of hydrogenation and further functionalization, see: a) L. T. Brechmann, J. F. Teichert, *Synthesis* **2020**, 52, 2483; b) S. W. Kim, W. Zhang, M. J. Krische, *Acc. Chem. Res.* **2017**, 50, 2371; c) T. Miao, Z.-Y. Tian, Y.-M. He, F. Chen, Y. Chen, Z.-X. Yu, Q.-H. Fan, *Angew. Chem. Int. Ed.* **2017**, 56, 4135; *Angew. Chem.* **2017**, 129, 4199; d) J. R. Zbieg, E. Yamaguchi, E. L. McInturff, M. J. Krische, *Science* **2012**, 336, 324; e) B. Breit, *Acc. Chem. Res.* **2003**, 36, 264; f) H.-Y. Jang, R. R. Huddleston, M. J. Krische, *J. Am. Chem. Soc.* **2002**, 124, 15156.
- [5] For selected examples, see: a) M. Wollenburg, A. Heusler, K. Bergander, F. Glorius, *ACS Catal.* **2020**, 10, 11365; b) L. Foppa, J. Dupont, *Chem. Soc. Rev.* **2015**, 44, 1886; c) Y. Wang, J. Yao, H. Li, D. Su, M. Antonietti, *J. Am. Chem. Soc.* **2011**, 133, 2362; d) H. Liu, T. Jiang, B. Han, S. Liang, Y. Zhou, *Science* **2009**, 326, 1250; e) H. Nagahara, M. Ono, M. Konishi, Y. Fukuoka, *Appl. Surf. Sci.* **1997**, 121/122, 448.
- [6] A. Grozavu, H. B. Hepburn, P. J. Smith, H. K. Potukuchi, P. J. Lindsay-Scott, T. J. Donohoe, *Nat. Chem.* **2019**, 11, 242.
- [7] a) M. Heitbaum, R. Fröhlich, F. Glorius, *Adv. Synth. Catal.* **2010**, 352, 357; b) F. Glorius, N. Spielkamp, S. Holle, R. Goddard, C. W. Lehmann, *Angew. Chem. Int. Ed.* **2004**, 43, 2850; *Angew. Chem.* **2004**, 116, 2910. For the use of achiral oxazolidinones in asymmetric hydrogenation, see: c) D. Fan, J. Zhang, Y. Hu, Z. Zhang, I. D. Gridnev, W. Zhang, *ACS Catal.* **2020**, 10, 3232.
- [8] a) E. Vitaku, D. T. Smith, J. T. Njardarson, *J. Med. Chem.* **2014**, 57, 10257; b) R. D. Taylor, M. MacCoss, A. D. G. Lawson, *J. Med. Chem.* **2014**, 57, 5845; c) D. O'Hagan, *Nat. Prod. Rep.* **2000**, 17, 435.
- [9] For selected examples, see: a) C. Yin, T. Yang, Y. Pan, J. Wen, X. Zhang, *Org. Lett.* **2020**, 22, 920; b) I. S. Hassan, A. N. Ta, M. W. Danneman, N. Semakul, M. Burns, C. H. Basch, V. N. Dippon, B. R. McNaughton, T. Rovis, *J. Am. Chem. Soc.* **2019**, 141, 4815; c) S. Jayakumar, K. Louven, C. Strohmann, K. Kumar, *Angew. Chem. Int. Ed.* **2017**, 56, 15945; *Angew. Chem.* **2017**, 129, 16161; d) H. R. Campello, J. Parker, M. Perry, P. Ryberg, T. Gallagher, *Org. Lett.* **2016**, 18, 4124; e) D. T. Khong, V. S. Pamarthy, T. Gallagher, Z. M. A. Judeh, *Eur. J. Org. Chem.* **2016**, 3084; f) Z.-T. He, Y.-B. Wei, H.-J. Yu, C.-Y. Sun, C.-G. Feng, P. Tian, G.-Q. Lin, *Tetrahedron* **2012**, 68, 9186; g) F. Zhang, Z.-J. Liu, J.-T. Liu, *Tetrahedron* **2010**, 66, 6864; h) H. K. Lee, J. S. Chun, C. S. Pak, *J. Org. Chem.* **2003**, 68, 2471; i) S. D. Koulocheri, P. Magiatis, S. A. Haroutounian, *J. Org. Chem.* **2001**, 66, 7915.
- [10] a) L. Shi, Y.-G. Zhou *Catalytic, Enantioselective Hydrogenation of Heteroaromatic Compounds in Organic Reactions*, Wiley, New York, **2018**, pp. 1–228; b) K. Mashima, H. Nagae, A. Iimuro, K. Higashida, *Heterocycles* **2017**, 95, 63; c) Z. Zhang, N. A. Butt, W. Zhang, *Chem. Rev.* **2016**, 116, 14769; d) D. Zhao, L. Candish, D. Paul, F. Glorius, *ACS Catal.* **2016**, 6, 5978; e) Z.-P. Chen, Y.-G. Zhou, *Synthesis* **2016**, 48, 1769; f) Y.-M. He, F.-T. Song, Q.-H. Fan, *Top. Curr. Chem.* **2013**, 343, 145; g) D.-S. Wang, Q.-A. Chen, S.-M. Lu, Y.-G. Zhou, *Chem. Rev.* **2012**, 112, 2557; h) F. Glorius, *Org. Biomol. Chem.* **2005**, 3, 4171.
- [11] For an asymmetric hydrogenation of N-methylated pyridones, see: J. Wysocki, C. Schlepphorst, F. Glorius, *Synlett* **2015**, 26, 1557.

- [12] A. Klapars, X. Huang, S. L. Buchwald, *J. Am. Chem. Soc.* **2002**, *124*, 7421.
- [13] Z. Yu, W. Jin, Q. Jiang, *Angew. Chem. Int. Ed.* **2012**, *51*, 6060; *Angew. Chem.* **2012**, *124*, 6164.
- [14] a) G. Diaz-Muñoz, I. L. Miranda, S. K. Sartori, D. C. de Rezende, M. A. Nogueira Diaz, *Chirality* **2019**, *31*, 776; b) D. A. Evans, G. Helmchen, M. Rüping *Chiral Auxiliaries in Asymmetric Synthesis in Asymmetric Synthesis—The Essentials* (Eds.: M. Christmann, S. Bräse), Wiley-VCH, Weinheim, **2007**, pp. 3–9; c) Y. Gnas, F. Glorius, *Synthesis* **2006**, 1899.
- [15] a) L. Pitzer, F. Schäfers, F. Glorius, *Angew. Chem. Int. Ed.* **2019**, *58*, 8572; *Angew. Chem.* **2019**, *131*, 8660; b) T. Gensch, F. Glorius, *Science* **2016**, *352*, 294–295.
- [16] a) T. Wagener, A. Heusler, Z. Nairoukh, K. Bergander, C. G. Daniliuc, F. Glorius, *ACS Catal.* **2020**, *10*, 12052; b) Z. Nairoukh, M. Wollenburg, C. Schlepphorst, K. Bergander, F. Glorius, *Nat. Chem.* **2019**, *11*, 264; c) X. Zhang, L. Ling, M. Luo, X. Zeng, *Angew. Chem. Int. Ed.* **2019**, *58*, 16785; *Angew. Chem.* **2019**, *131*, 16941; d) M. P. Wiesenfeldt, Z. Nairoukh, W. Li, F. Glorius, *Science* **2017**, *357*, 908; e) M.-W. Chen, Y. Ji, J. Wang, Q.-A. Chen, L. Shi, Y.-G. Zhou, *Org. Lett.* **2017**, *19*, 4988; f) R.-N. Guo, X.-F. Cai, L. Shi, Z.-S. Ye, M.-W. Chen, Y.-G. Zhou, *Chem. Commun.* **2013**, *49*, 8537.
- [17] Deposition numbers 2043968 (**1**) and 2043969 (**10**) contain(s) the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service.
- [18] a) B. Mattson, W. Foster, J. Greimann, T. Hoette, N. Le, A. Mirich, S. Wankum, A. Cabri, C. Reichenbacher, E. Schwanke, *J. Chem. Educ.* **2013**, *90*, 613; b) I. Horiuti, M. Polanyi, *Trans. Faraday Soc.* **1934**, *30*, 1164.

Manuscript received: December 17, 2020

Accepted manuscript online: January 18, 2021

Version of record online: February 12, 2021