

Synthesis of 3,3,6,6-Tetraaryl-1,2-Dioxanes via TiO₂-catalyzed Photooxygenation of 1,1-Diarylethenes in the Presence of Mg(ClO₄)₂

Hajime Maeda, Hisayuki Miyamoto, and Kazuhiko Mizuno*

Department of Applied Chemistry, Graduate School of Engineering, Osaka Prefecture University, Sakai, Osaka 599-8531

(Received January 20, 2004; CL-040078)

Photooxygenation of methoxy-substituted 1,1-diarylethenes and 1,1,8,8-tetraaryl-1,7-octadienes catalyzed by titanium dioxide proceeded via photoinduced electron transfer to give 3,3,6,6-tetraaryl-1,2-dioxanes in high yields. The photooxygenation was remarkably accelerated by the addition of Mg(ClO₄)₂.

Semiconductor-catalyzed photoreactions of organic molecules are of current interest.¹ Semiconductors such as TiO₂, CdS, ZnS are utilized as redox type heterogeneous photocatalysts for various types of photoinduced electron transfer reactions using photoexcitation from their valence bands to conduction bands.² Hitherto various alkenes are exposed to the semiconductor-catalyzed photoreactions in the presence of oxygen, and it is already known that the alkenes are converted to ketones, aldehydes, epoxides, hydroperoxides, and so on.³ However, the efficiency, selectivity, and chemical yields are not necessarily high in these photoreactions. To explore the synthetic utility for such reactions, we have investigated the photooxygenation of methoxy-substituted 1,1-diarylethenes and found that 3,3,6,6-tetraaryl-1,2-dioxanes were obtained in high yields.

Photoirradiation of an acetonitrile solution containing 1,1-bis(*p*-methoxyphenyl)ethene (**1a**) in the presence of suspended TiO₂ and Mg(ClO₄)₂ under a constant stream of dioxygen through a Pyrex filter by a 300 W-high pressure mercury lamp for 40 min afforded 3,3,6,6-tetrakis(*p*-methoxyphenyl)-1,2-dioxane (**2a**) in a 97% isolated yield (Scheme 1, Table 1). In the absence of Mg(ClO₄)₂, **2a** was slowly formed accompanying methoxy-substituted benzophenone **3a**. Photoreaction of *p*-chloro derivative **1b** gave a small amount of benzophenone derivative **3b** without any formation of **2b**. Photooxygenation of 1,1,8,8-tetrakis(*p*-methoxyphenyl)-1,7-octadiene (**4a**) in the presence of Mg(ClO₄)₂ proceeded intramolecularly to give


Table 1. Photooxygenation of 1,1-Diarylethenes Catalyzed by Titanium Dioxide^a

Substrate	Additive	Irradiation time/min	Products and yields ^b	Recovery of substrate ^b
1a	none	40	2a (6%), 3a (10%)	56%
1a	Mg(ClO ₄) ₂	40	2a (97%)	0%
1b	none	40	3b (12%) ^c	88% ^c
1b	Mg(ClO ₄) ₂	40	3b (8%)	73%
4a	none	40	5a (9%) ^c	91% ^c
4a	none	960	5a (75%) ^c , 3a (25%) ^c	0%
4a	Mg(ClO ₄) ₂	40	5a (71%)	0%
4a	KClO ₄	40	5a (34%) ^c	66% ^c
4a	NaClO ₄	40	5a (25%)	28%
4a	LiClO ₄	40	5a (41%)	44%
4a	LiBF ₄	40	5a (51%)	40%
4b	none	40	—	97%
4b	Mg(ClO ₄) ₂	40	—	87%


^a Conditions: Substrate (0.25 mmol), TiO₂ (10 mg), additive (0.125 mmol), CH₃CN (15 mL), oxygen bubbling, rt. ^b Isolated yields. ^c Determined by ¹H NMR.

trans-fused 1,2-dioxane derivative **5a** in a 71% isolated yield. In the absence of Mg(ClO₄)₂, prolonged photoirradiation was required to obtain **5a**. Addition of other inorganic salts such as KClO₄, NaClO₄, LiClO₄, and LiBF₄ also enhanced the rate of the photooxygenation, but their efficiency was lower than that of Mg(ClO₄)₂. Photoreaction of unsubstituted 1,1,8,8-tetraphenyl-1,7-octadiene (**4b**) resulted in almost recovery of **4b**.

Irradiation of TiO₂ in acetonitrile causes the promotion of an electron into the conduction band (−0.8 V vs SCE) to generate a hole in the valence band (+2.4 V vs SCE).^{1,4} One electron transfer from **1a** (*E*_{ox} = 1.32 V vs SCE in CH₃CN)⁵ and **4a** (*E*_{ox} = 0.95 V vs Ag/Ag⁺ in CH₃CN)⁶ to the valence band should take place as an exothermic process to produce the radical cations of these alkenes, because their oxidation potentials are sufficiently low.^{1a,b} In addition, electron transfer from the conduction band to O₂ (*E*_{red}⁰ = −0.86 V vs SCE in CH₃CN)⁷ is thermodynamically permissible process to produce O₂^{•−}.^{1a,3a,b,g} From these results, we propose a plausible mechanism for the TiO₂-catalyzed photooxygenation of 1,1-diarylethenes as exemplified in Scheme 2 using the photoreaction of **4a**. The initial step is the excitation of TiO₂ followed by one-electron transfer from **4a** to the valence band of TiO₂ to produce monomer radical cation of **4a**. The monomer radical cation of **4a** cyclizes intramolecularly to give distonic radical cation **6a** via dimer radical cation of **4a**. The attack of triplet dioxygen on the 1,4-radical cation **6a** generates 1,6-radical cation **7a** which gives 1,2-dioxane derivative **5a** by one-electron reduction. The other possible pathway for the photooxygenation is the attack of O₂^{•−} to **6a**. Formation of benzophenone derivative **3a** can be explained by the decomposition of dioxetane **8a**, which is produced by the oxygenation of monomer radical cation. The enhancement of the ef-



Scheme 1.



Scheme 2.

iciency by the addition of $\text{Mg}(\text{ClO}_4)_2$ can be explained by the suppression of back electron transfer from TiO_2^- to $4\text{a}^{+\bullet}$ due to the interaction between Mg^{2+} and ionic species such as TiO_2^- and $\text{O}_2^{\bullet-}$ or the interaction between ClO_4^- and $4\text{a}^{+\bullet}$.^{2r,8-10}

Photooxygenation of **1a** and **4a** to give 1,2-dioxanes under photoinduced electron transfer conditions have already been reported in homogeneous solutions using electron accepting photosensitizers such as 9,10-dicyanoanthracene.^{5,6,11} But the present system using TiO_2 as a heterogeneous photocatalyst has a notable advantage; the photocatalyst can be removed only by the filtration of the reaction mixture. For synthetic use, TiO_2 is a clean, low cost, efficient, and facile tool for the synthesis of 3,3,6,6-tetraaryl-1,2-dioxanes.

In conclusion, it was demonstrated that 1,2-dioxane derivatives can be synthesized by the photooxygenation of electron-rich diarylalkenes by use of TiO_2 as a heterogeneous photocatalyst. Scope and mechanisms are now under investigation.

We would like to thank Prof. M. Anpo, Dr. M. Matsuoka, Dr. M. Takeuchi, Ms. R. Takeuchi, and Mr. H. Nakagawa at Osaka Prefecture University for measurement of diffuse reflectance spectra. We also thank Ishihara Sangyo Co., Ltd., for a gift of TiO_2 (ST-21). This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas (417) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

References and Notes

- 1 a) M. A. Fox, *Acc. Chem. Res.*, **16**, 314 (1983). b) K. Tokumaru, H. Sakuragi, T. Kanno, T. Oguchi, H. Misawa, Y. Shimamura, and Y. Kuriyama, "Organic Phototransformations in Nonhomogeneous Media," ed. by M. A. Fox, ACS Publications, Washington D. C. (1985), p 43. c) M. R. Hoffmann, S. T. Martin, W. Choi, and D. W. Bahnemann, *Chem. Rev.*, **95**, 69 (1995).
- 2 a) J. W. Pavlik and S. Tantayanon, *J. Am. Chem. Soc.*, **103**, 6755

- (1981). b) M. Fujihira, Y. Satoh, and T. Osa, *Bull. Chem. Soc. Jpn.*, **55**, 666 (1982). c) R. A. Barber, P. de Mayo, and K. Okada, *J. Chem. Soc., Chem. Commun.*, **1982**, 1073. d) M. A. Fox, C.-C. Chen, and J. N. N. Younathan, *J. Org. Chem.*, **49**, 1969 (1984). e) A. M. Draper, M. Ilyas, P. de Mayo, and V. Ramamurthy, *J. Am. Chem. Soc.*, **106**, 6222 (1984). f) H. Al-Ekabi and P. de Mayo, *Tetrahedron*, **42**, 6277 (1986). g) M. A. Fox and J. N. Younathan, *Tetrahedron*, **42**, 6285 (1986). h) P. de Mayo and G. Wenska, *Tetrahedron*, **43**, 1661 (1987). i) M. A. Fox and A. A. Abdel-Wahab, *Tetrahedron Lett.*, **31**, 4533 (1990). j) C. Joyce-Pruden, J. K. Pross, and Y. Li, *J. Org. Chem.*, **57**, 5087 (1992). k) B. Ohtani, K. Iwai, H. Kominami, T. Matsuura, Y. Kera, and S. Nishimoto, *Chem. Phys. Lett.*, **242**, 315 (1995). l) B. Ohtani, S. Kusakabe, K. Okada, S. Tsuru, K. Izawa, Y. Amino, and S. Nishimoto, *Tetrahedron Lett.*, **36**, 3189 (1995). m) A. Pace, S. Buscemi, N. Vivona, and T. Caronna, *Heterocycles*, **53**, 183 (2000). n) C. Minero, G. Mariella, V. Maurino, and E. Pelizzetti, *Langmuir*, **16**, 2632 (2000). o) C. Minero, G. Mariella, V. Maurino, D. Vione, and E. Pelizzetti, *Langmuir*, **16**, 8964 (2000). p) B. Ohtani, S. Kusakabe, K. Okada, S. Tsuru, S. Nishimoto, Y. Amino, K. Izawa, Y. Nakato, M. Matsumura, Y. Nakaoka, and Y. Nosaka, *J. Chem. Soc., Perkin Trans. 2*, **2001**, 201. q) S. Vijaikumar, N. Somasundaram, and C. Srinivasan, *Appl. Catal., A*, **223**, 129 (2002). r) H. Maeda, H. Nakagawa, and K. Mizuno, *Photochem. Photobiol. Sci.*, **2**, 1056 (2003).
- 3 a) T. Kanno, T. Oguchi, H. Sakuragi, and K. Tokumaru, *Tetrahedron Lett.*, **21**, 467 (1980). b) M. A. Fox and C.-C. Chen, *J. Am. Chem. Soc.*, **103**, 6757 (1981). c) M. A. Fox, B. Lindig, and C.-C. Chen, *J. Am. Chem. Soc.*, **104**, 5828 (1982). d) M. A. Fox and C.-C. Chen, *Tetrahedron Lett.*, **24**, 547 (1983). e) S. Yanagida, K. Mizumoto, and C. Pac, *J. Am. Chem. Soc.*, **108**, 647 (1986). f) M. A. Fox, D. D. Sackett, and J. N. Younathan, *Tetrahedron*, **43**, 1643 (1987). g) J. L. Muzyka and M. A. Fox, *J. Photochem. Photobiol., A*, **57**, 27 (1991).
- 4 a) S. N. Frank and A. J. Bard, *J. Am. Chem. Soc.*, **97**, 7427 (1975). b) J. M. Bolts and M. S. Wrighton, *J. Phys. Chem.*, **80**, 2641 (1976). c) G. Rothenberger, J. Moser, M. Grätzel, N. Serpone, and D. K. Sharma, *J. Am. Chem. Soc.*, **107**, 8054 (1985). d) N. Serpone, D. Lawless, R. Khairutdinov, and E. Pelizzetti, *J. Phys. Chem.*, **99**, 16655 (1995).
- 5 K. Gollnick, A. Schnatterer, and G. Utschick, *J. Org. Chem.*, **58**, 6049 (1993).
- 6 a) T. Tamai, K. Mizuno, I. Hashida, and Y. Otsuji, *Tetrahedron Lett.*, **34**, 2641 (1993). b) K. Mizuno, T. Tamai, I. Hashida, Y. Otsuji, Y. Kuriyama, K. Tokumaru, *J. Org. Chem.*, **59**, 7329 (1994).
- 7 D. T. Sawyer, T. S. Calderwood, K. Yamaguchi, and C. T. Angelis, *Inorg. Chem.*, **22**, 2577 (1983).
- 8 a) K. Mizuno, N. Ichinose, and Y. Otsuji, *Chem. Lett.*, **1985**, 455. b) K. Mizuno, N. Kamiyama, N. Ichinose, and Y. Otsuji, *Tetrahedron*, **41**, 2207 (1985). c) K. Mizuno and Y. Otsuji, *J. Synth. Org. Chem. Jpn.*, **47**, 916 (1989). d) A. Loupy, B. Tchoubar, and D. Astruc, *Chem. Rev.*, **92**, 1141 (1992). e) K. Mizuno, N. Ichinose, and Y. Otsuji, *J. Org. Chem.*, **57**, 1855 (1992). f) K. Mizuno and Y. Otsuji, "Topics in Current Chemistry," ed. by J. Mattay, Springer-Verlag, Heidelberg (1994), Vol. 169, p 301. g) S. Fukuzumi, *Bull. Chem. Soc. Jpn.*, **70**, 1 (1997). h) K. Mizuno, K. Nire, H. Sugita, and H. Maeda, *Tetrahedron Lett.*, **42**, 2689 (2001). i) S. Fukuzumi, *J. Phys. Org. Chem.*, **15**, 448 (2002). j) S. Fukuzumi, *Pure Appl. Chem.*, **75**, 577 (2003).
- 9 Diffuse reflectance spectrum of suspension of TiO_2 in CH_3CN was not changed by the addition of $\text{Mg}(\text{ClO}_4)_2$.
- 10 The binding energies of metal ions with $\text{O}_2^{\bullet-}$ decrease in the order of $\text{Mg}^{2+} > \text{Li}^+ > \text{Na}^+$. This fact is in good agreement with the effect of metal salts shown in Table I. a) S. Fukuzumi and K. Ohkubo, *Chem.—Eur. J.*, **6**, 4532 (2000). b) K. Ohkubo, S. C. Menon, A. Orita, J. Otera, and S. Fukuzumi, *J. Org. Chem.*, **68**, 4720 (2003).
- 11 a) R. K. Haynes, M. K. S. Probert, and I. D. Wilmot, *Aust. J. Chem.*, **31**, 1737 (1978). b) J. Eriksen and C. S. Foote, *J. Am. Chem. Soc.*, **102**, 6083 (1980). c) K. Gollnick and A. Schnatterer, *Tetrahedron Lett.*, **25**, 185 (1984). d) K. Gollnick and A. Schnatterer, *Tetrahedron Lett.*, **25**, 2735 (1984). e) S. L. Mattes and S. Farid, *J. Am. Chem. Soc.*, **108**, 7356 (1986). f) M. Kojima, A. Ishida, and S. Takamuku, *Chem. Lett.*, **1993**, 979. g) T. Tamai, K. Mizuno, I. Hashida, Y. Otsuji, A. Ishida, and S. Takamuku, *Chem. Lett.*, **1994**, 149.