


Table I. Alkynyl Sulfenylation of Olefins

entry	olefin ^a	alkyne	reagent	solvent ratio ^b Cl THF	temp ^c	time ^c	product ^d	isolated yield ^e
1		$n\text{C}_5\text{H}_{11}\text{C}\equiv\text{CH}$ 7	A	4/1	80 °C	2 h		88%
2		7	B	5/1	rt	45 min		86%
3		7	B	5/1	rt	35 min		86%
4		7	A	4/1	rt	1 h		74%
5		7	A	5/1	0 °C	45 min		83%
6	$(\text{CH}_2)_8\text{CO}_2\text{CH}_3$	7	A	5/1	0 °C	1 h		54%
7		$\text{TBDSOCH}_2\text{C}\equiv\text{CH}$	B	5/1	40 °C	40 min		72%
8		$\text{TBDSOCH}_2\text{C}\equiv\text{CH}$	B	14/1	rt	3 h		53%

^a Normally a 2:1 ratio of the reagent to the olefin is employed. ^b Toluene and hexane derived from the organoaluminum and *n*-butyllithium, respectively, are also present. The given ratio is for the added solvents at the final stage of reaction. ^c Time and temperature at the final stage of reaction, rt = room temperature. ^d All new compounds have been fully characterized. In each case, spectroscopy including ¹³C NMR and chromatography (TLC and GC), indicate a single regio- and/or stereoisomer was present. ^e The products have been purified either by chromatography or distillation. ^f In this case, a byproduct, whose structure has not been deduced, contaminates the major product.





to additions to episulfonium ions, and subsequent chemistry (vide infra). The reaction is completely regioselective (entries 2, 3, 6, and 8) leading to an anti-Markovnikov addition even in the case of a trisubstituted olefin (entry 3) in contrast to other reactions of episulfonium salts. Preliminary results indicate good chemoselectivity too (entries 6–8).

The equivalent of nucleophilic addition to a double bond was explicitly demonstrated in the case of **6** and **8**, which, upon exposure to W-2 or W-5 Raney nickel in refluxing ethanol, gave **12** in 87% and 84% yield, respectively (eq 4). The presence of




a doublet for the ring methyl group in **12b** also establishes the regiochemistry of the addition to 1-methylcyclohexene.

The equivalent of nucleophilic substitution of an olefin was explicitly demonstrated in the case of **6**, **9**, **10**, and **11** (eq 5–8).



Eliminations were accomplished either by oxidation to the sulfoxide followed by heating in a sealed tube at 140–220 °C (CaCO_3 , $\text{C}_2\text{H}_5\text{OCH}=\text{CH}_2$)⁹ [eq 6 and 7 (**14a**:**14b** 2:1)] or by alkylation-elimination [$\text{CF}_3\text{SO}_3\text{CH}_2\text{CO}_2\text{C}_2\text{H}_5$, CH_3CN then DBU¹⁰ eq 5, 7 (**14a**:**14b** 5:1), and 8]. Since both eliminations are known to be *cis*, the elimination products provide chemical evidence for the stereochemistry of the alkynyl sulfenylation.

The versatility of this approach also derives from the versatility of the acetylene. One illustration is the net hydration of the acetylene to a ketone. While mercury-based methods proved less than satisfactory, hydroboration–oxidation¹¹ of **6** and **10** gave the ketones **15** and **16** in 91% and 79% yields, respectively (eq 9).



A short synthesis of the sex pheromone of *Malacosoma disstria* (forest tent caterpillar) **17**¹² from 1-hexene utilizes the adduct **11** (eq 10). Since elimination at the stage of the acetylene gave an *E,Z* mixture (eq 8), an alternative was sought. By an increase in the effective steric bulk of the alkyl group, enhanced stereo-selectivity should result. Indeed, reduction of **11** (H_2 , Lindlar's catalyst, hexane, 99%) followed by elimination (i, $\text{TfOCH}_2\text{CO}_2\text{C}_2\text{H}_5$, CH_3CN ; ii, DBU, 46 °C; iii, $(\text{C}_4\text{H}_9)_4\text{NF}$, THF; 69% overall) proceeds stereoselectively to give the desired diene (eq 10) with high (>98%) stereocontrol.


This direct elaboration of olefins with nucleophiles has several notable merits. The activating sulfur conjunctive reagent, DMTSF, is a readily available, crystalline solid that can be stored indefinitely. The reactions work well with mono-, di-, and tri-

(9) Trost, B. M.; Salzmann, T. N.; Hiroi, K. *J. Am. Chem. Soc.* **1976**, *98*, 4887 and references therein. For reviews, see: Trost, B. M. *Acc. Chem. Res.* **1978**, *11*, 453; *Chem. Rev.* **1978**, 363.

(10) Vedejs, E.; Engler, D. A. *Tetrahedron Lett.* **1976**, 3487.

(11) Brown, H. C.; Zweifel, G. *J. Am. Chem. Soc.* **1960**, *82*, 3222.

(12) Chisholm, M. D.; Underhill, E. W.; Steck, W.; Slessor, K. N.; Grant, G. G. *Environ. Entomol.* **1980**, *39*, 287.



substituted olefins and with high chemo-, regio-, and diastereoselectivity. The anti-Markovnikov orientation even with a tri-substituted double bond is particularly noteworthy and suggests that, in the opening of the presumed episulfonium ion intermediate, bond formation and cleavage occur more nearly to the same extent at the transition state regardless of the olefin substitution, unlike other olefin additions. The importance of vinylacetylenes themselves (e.g., histrionicotoxins) and their use as intermediates for the stereocontrolled synthesis of dienes attaches additional interest to this olefin elaboration method.

Acknowledgment. We thank the National Science Foundation for their generous support of our program.

Registry No. 1, 5799-67-7; 3, 90108-00-2; 4, 34046-61-2; 5, 41578-02-3; 6, 90108-01-3; 7, 628-71-7; 8, 90108-02-4; 9, 90108-03-5; 10, 90108-04-6; 11, 90108-05-7; 12a, 5617-41-4; 12b, 10483-56-4; 13, 90108-06-8; 14a, 90108-07-9; 14b, 90108-09-1; 15, 90108-09-1; 16, 90108-10-4; 17, 72922-18-0; TMSOCH₂C≡CH, 76782-82-6; TBDMSO(CH₂)₄C≡CH, 73448-13-2; Et₂AlCl, 96-10-6; Et₃Al, 97-93-8; cyclohexene, 110-83-8; methylenecyclopentane, 1528-30-9; 1-methyl-1-cyclohexene, 591-49-1; (*E*)-dec-5-ene, 7433-56-9; (*Z*)-dec-5-ene, 7433-78-5; methyl undec-10-enoate, 111-81-9; 1-hexene, 592-41-6; 1-(methylthio)-1-(oct-2-yn-1-yl)cyclopentane, 90108-11-5; methyl 3-(methylthio)undec-5-ynoate, 90108-12-6; *trans*-1-[3-(*tert*-butyldimethylsiloxy)-1-propynyl]-2-(methylthio)cyclohexane, 90108-13-7; 1-(1-heptynyl)cyclohexene, 90108-14-8; (*E*)-1-(*tert*-butyldimethylsiloxy)dodec-7-en-5-yne, 90108-15-9; (*Z*)-1-(*tert*-butyldimethylsiloxy)dodec-7-en-5-yne, 90108-17-1; (*Z*)-1-(*tert*-butyldimethylsiloxy)-8-(methylthio)dodec-5-ene, 90108-16-0; lithium triethylheptynylaluminum, 90108-18-2; lithium diethylheptynylaluminum, 90108-19-3; 1-lithio-1-heptyne, 42017-07-2.

Isolation and Characterization of (PNP)₂Co(CN)₄, an Unusual Square-Planar Cobalt(II) Complex

Stephen J. Carter, Bruce M. Foxman, and Louis S. Stuhl*

Department of Chemistry, Brandeis University
Waltham, Massachusetts 02254

Received February 13, 1984

Cyanide ion is regarded as promoting the formation of higher coordination numbers in transition-metal complexes.¹ Thus it is not surprising that, although good evidence for loss of CN⁻ from Co(CN)₅³⁻ in solution has been presented,^{2,3} no simple monomeric Co(CN)₄²⁻ species has yet been isolated in the solid state. It occurred to us that, since many cyanide complexes are anionic, appropriate variation of counterions would permit stabilization of different geometries or coordination numbers. We now report the successful synthesis and characterization of (PNP)₂Co(CN)₄ (1),⁴ a simple, monomeric, coordinately unsaturated Co(II) complex. This has proven to be a very rare example of low-spin, square-planar Co(II) with unidentate ligands,⁵ in fact to our knowledge, the first with only a *single* type of unidentate ligand.

Anhydrous CoCl₂ reacts with 4 equiv of (PNP)CN in dimethylformamide solution to produce a green solution from which pale blue (PNP)₂Co(CN)₄ can be crystallized by cooling. Com-

Table I. IR Data for Solutions of CoCl₂ in Acetonitrile to Which Increasing Amounts of (PNP)CN Are Added^a


Co:CN ⁻	color	IR, ^b cm ⁻¹
3	deep blue	2131 (w), 2120 (w), 2096 (s), 2089 (m)
4	blue-black	2131 (vw), 2099 (m), 2090 (s), 2071 (w)
5	olive-green	2090 (m), 2072 (s)
6	yellow	2090 (w), 2072 (s)

^a Compare (PNP)₃Co(CN)₅, ν_{CN} 2072 cm⁻¹, K₃Co(CN)₆, ν_{CN} 2131, 2128 cm⁻¹ (mineral oil mulls). ^b IR solutions 1.75 × 10⁻² M.

pound 1 is isolated as a DMF solvate. Drying at 80 °C in vacuo produces solvent-free material as a pale blue, air-sensitive powder. Satisfactory analytical data was obtained on a sample recrystallized from DMF and dried as above, mp 227.5–229 °C (corrected). Anal. (C₇₆H₆₀N₆P₄Co) C, H, N, Co (Pascher Mikroanalytische Laboratorium, Bonn). The infrared spectrum of the unsolvated solid (Nujol mull) shows bands at 2100 (sh, s) and 2095 (s) (ν_{C≡N}) and 397 cm⁻¹ (ν_{Co-C}). The magnetic moment of 1 in the solid state was found to be 2.15 μ_B.⁶ Solution spectral and magnetic data are solvent and concentration dependent; the resultant equilibria are currently under investigation. The magnetic moment of 1 in acetonitrile (1.25 × 10⁻¹ M) is 2.52 μ_B.⁷ Infrared spectral data for varying ratios of cobalt to added cyanide in acetonitrile solution are in Table I; the IR spectrum of 1 in DMF or CH₃CN solution is similar to the 1:4 Co:CN solution. Only one cyanide absorption, at 2096 cm⁻¹, was detected for 1 in CH₂Cl₂ solution.

The only previously reported solid apparently containing the Co(CN)₄²⁻ anion is K₂Co(CN)₄, isolated from liquid ammonia.⁸ The spectroscopic properties of this material clearly indicate its polymeric nature. Monomeric Co(CN)₄²⁻, as well as Co(CN)₃(solvent)⁻, have been observed in solution, but not isolated.^{3,9} Compound 1 has properties distinctly different from those of K₂Co(CN)₄ with solubility and spectral data suggestive of a stable form of monomeric Co(CN)₄²⁻. The magnetic moment of 1 rules out a tetrahedral coordination geometry. Although all known Co(II)L₄ species (L = unidentate ligand) exist in a tetrahedral configuration, the possibility of cyanide inducing a square-planar coordination geometry should not be unexpected since cyanide is a very strong field ligand (cf. Ni(CN)₄²⁻). Square-planar Co(II) complexes with macrocyclic ligands exhibit magnetic moments in the range 2.1–2.8 μ_B,¹⁰ suggesting such a structure for the solid. The cobalt(II) bis(dithioacetylacetonate) complex, Co(SacSac)₂, has a magnetic moment of 2.35 μ_B¹¹ and has been shown to be square planar by X-ray crystallography.¹² The IR data for 1 indicate only the presence of terminal cyanide groups in the solid state, thus precluding the possibility of bridging cyanides bonded to a low-spin octahedral Co(II) center.¹³

The IR data in Table I are consistent with the idea that in a coordinating solvent, such as acetonitrile, the [Co(CN)₄]²⁻ anion participates in a dynamic equilibrium:



Consistent with this, the electronic spectrum of 1 is dominated

(1) Rigo, P.; Turco, A. *Coord. Chem. Rev.* **1974**, *13*, 133–172.

(2) Pratt, J. M.; Williams, R. J. P. *J. Chem. Soc. A* **1967**, 1291–1298.

(3) White, D. A.; Solodar, A. J.; Baizer, M. M., *Inorg. Chem.* **1972**, *11*, 2160–2167.

(4) PNP is the bis(triphenylphosphine)iminium cation.

(5) The compound (Mes)₂Co(PtPh₂)₂ and analogous ortho-substituted aryl cobalt phosphine complexes are square planar. This is attributed to steric hindrance by the ortho substituents. No such steric effect is operative in 1. Owston, P. G.; Rowe, J. M., *J. Chem. Soc.* **1963**, 3411–19. Falvello, L.; Gerloch, M., *Acta Cryst.* **1979**, B35, 2547–50.

(6) Because the PNP cation is so large, an accurate molar diamagnetic correction is necessary. The value used (7.37 × 10⁻⁴) was determined from values for PPh₃ and standard Pascal's constants: Figgis, B. N.; Lewis, J. In "Technique of Inorganic Chemistry"; Jonassen, H. B., Weissberger, A., Eds.; Interscience: New York, 1965; Vol. 4, pp 142–143.

(7) Determined by using the method of: Evans, D. F. *J. Chem. Soc.* **1959**, 2003–2005.

(8) Mosha, D. M. S.; Nicholls, D. *Inorg. Chim. Acta* **1980**, *38*, 127–130.

(9) Weisz, A.; Gutmann, V. *Monatsh. Chem.* **1971**, *101*, 19–26. Gutmann, V.; Wegleitner, K. H., *Ibid.* **99**, 368–79.

(10) Nicholls, D. In "Comprehensive Inorganic Chemistry"; Trotman-Dickenson, A. F., Ed.; Pergamon Press: Oxford, 1973; Vol. 3, p 1092.

(11) Barraclough, C. G.; Martin, R. L.; Stewart, J. M. *Aust. J. Chem.* **1969**, *22*, 891–904.

(12) Beckett, R.; Hoskins, B. F. *J. Chem. Soc., Dalton Trans.* **1974**, 622–625.

(13) Hartman, J. R.; Hints, E. J.; Cooper, S. R. *J. Chem. Soc., Chem. Commun.* **1984**, 386–387.