

Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Synthesis and Biological Evaluation of Some New Thioether-Ester Crown Ethers

Seyed Mohammad Seyedi ^a, Ali Sadeghian ^b, Hamid Sadeghian ^a, Ayla Hazrathoseyni ^a & Mohammad Sadeghian ^c

^a Department of Chemistry, Ferdowsi University of Mashhad, Mashhad, I. R. Iran

^b Department of Microbiology, Ghaem Hospital, Mashhad, I. R. Iran

^c Faculty of Medicine, Medical University of Mashhad, Mashhad, I. R. Iran

Published online: 18 Feb 2007.

To cite this article: Seyed Mohammad Seyedi, Ali Sadeghian, Hamid Sadeghian, Ayla Hazrathoseyni & Mohammad Sadeghian (2007) Synthesis and Biological Evaluation of Some New Thioether-Ester Crown Ethers, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 182:2, 265-272, DOI: [10.1080/10426500600917060](https://doi.org/10.1080/10426500600917060)

To link to this article: <http://dx.doi.org/10.1080/10426500600917060>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and

are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis and Biological Evaluation of Some New Thioether-Ester Crown Ethers

Seyed Mohammad Seyed

Department of Chemistry, Ferdowsi University of Mashhad,
Mashhad, I. R. Iran

Ali Sadeghian

Department of Microbiology, Ghaem Hospital, Mashhad, I. R. Iran

Hamid Sadeghian

Ayla Hazrathoseyni

Department of Chemistry, Ferdowsi University of Mashhad,
Mashhad, I. R. Iran

Mohammad Sadeghian

Faculty of Medicine, Medical University of Mashhad, Mashhad,
I. R. Iran

*New thioether-ester crown ethers have been synthesized starting from oxalyl chloride and different β,β' -dihydroxydithioethers. The synthesized compounds are screened for their antibacterial activity. Among the macrocyclic thioether-esters (**5a–j**), only 5,12-di[(allyloxy)methyl]-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (**5e**) and 5,12-di(isopropoxymethyl)-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (**5f**) were effective inhibitors against *Staphylococcus aureus* methicillin resistance and *Pseudomonas aeruginosa* with an MIC value of 525 and 265 μM . Structures of the synthesized compounds have been confirmed by ^1H NMR, ^{13}C NMR, and MS spectral studies.*

Keywords Dihydroxydithioethers; *Pseudomonas aeruginosa*; *Staphylococcus aureus*; thioether-ester crown ethers

Received May 15, 2006; accepted June 26, 2006.

Address correspondence to Seyed Mohammad Seyed, Ferdowsi University of Mashhad, Department of Chemistry, Faculty of Sciences, Mashhad, 91775 I. R. Iran.
E-mail: smseyedi@yahoo.com

INTRODUCTION

β,β' -dihydroxydithioethers with two α,α' -substitutions have been recently synthesized.¹ The two secondary β,β' -dihydroxy groups make these compounds susceptible reagents to synthesize new thiocrown ethers possessing various sidearms.^{2,3} Biological activities of such sidearmed crownethers have been studied³ but no biological evaluations have been reported on β,β' -dihydroxydithioethers and their corresponding thiocrown ethers. In this article, we describe the synthesis of β,β' -dihydroxydithioether (**3a-j**) and their corresponding thioether-ester crown ethers (**5a-j**). The antibacterial activity of these compounds is also investigated.


RESULTS

Synthesis

β,β' -dihydroxydithioether (**3a**) and the mixture of two diastereomers¹ of its derivatives (**3b-j**) were prepared by the action of two mole equivalents of oxiranes (**1a-j**) with dimercaptoethane (**2**) using benzyltrimethylammonium hydroxide (Triton B) as a catalyst¹ (Scheme 1). All dihydroxy compounds showed one spot on TLC except the (**3d**), which was a mixture and was purified by column chromatography (silica gel 60, 0.04–0.067 mm, hexane-ethylacetate 1:1). Treatment of dihydroxy compounds (**3a-i**) and (**3j**) with oxalyl chloride (**4**) under a basic condition of triethylamine in dried dichloromethane afforded 5,12-disubstituted 1,4-dioxo-7,10-dithiacyclododecane-2,3-diones (**5a-i**) and perhydrodibenzo[*e,k*][1,4,7,10]dioxadithiacyclododecine-6,7-dione (**5j**), respectively⁴ (Scheme 1).

Biological Evaluations

Antibacterial activities of β,β' -dihydroxydithioethers (**3a-j**) and their corresponding thioether-ester crown ether (**5a-j**) were evaluated. MIC values, i.e., the lowest concentration of a drug that prevents growth of a particular pathogen,⁵ of (**3a-j**) and (**5a-j**) against two strains of bacteria *S. aureus* methicilin resistance and *P. aeruginosa* were measured. Ten isolated strains of *S. aureus* and *P. aeruginosa* from different organs of patients at the Microbiological Laboratory of Ghaem Hospital of Medical University of Mashhad–Iran were tested. Oxacillin (for *S. aureus*) and Gentamycin (for *P. aeruginosa*) were used as a positive control in all tests, and their MIC values were expressed in μM . Among the synthesized compounds (**3a-j**) and (**5a-j**), only (**5e**) and (**5f**) showed significant



SCHEME 1

activities against *S. aureus* and *P. aeruginosa* with MIC values of 525 and 265 μM (100 and 200 $\mu\text{g/mL}$). These results were compared with Gentamycin and Oxacillin activity with the standard MIC values of 16 and 32 $\mu\text{g/mL}$, respectively.

DISCUSSION

The ring opening of the starting oxiranes (**1b–i**) was region specific by a nucleophilic attack on the terminal carbon atoms affording secondary diols. This regiospecificity was not observed in the case of (**1d**), which gave a mixture of isomeric derivatives. Compounds (**3b–j**) were obtained as

a mixture of isomeric diastereomers. Their spectral data were perfectly consistent with the literature data.¹

Accordingly, compounds (**5b–j**) were obtained as a mixture of diastereomers, but the two stereogenic centers within the molecule were far from each other so they could not be discerned by the NMR technique. Their ¹H NMR spectrum showed a sharp singlet for the two methylene groups of the dimercaptoethane moiety. This is not the case for compound (**5j**) for which a multiplet was observed for these methylene groups. This may be explained if one assumes that the ring opening of cyclohexene oxide (**1j**) gave exclusively *trans*-diols, which were formed as an equimolar mixture of *meso*- and *threo*-.¹

The *S. aureus* methicillin resistance and *P. aeruginosa* have become a major nosocomial pathogen in communities, long-term-care facilities, and tertiary-care hospitals.^{6–9} Although (**5e**) and (**5f**) exhibited weak antibacterial activities against these pathogens in comparison with Gentamycin and Oxacillin, this study can be a beginning for the synthesis and evaluation of a new generation of antibacterial agents. Although both compounds of (**5e**) and (**5f**) are each a mixture of diastereomers,¹ if one can obtain a diastereomerically pure species of any of these compounds, the MIC value of pure diastereomer might be less or more than what we expected.

CONCLUSION

The aim of this study was to develop an efficient synthetic approach to construct various 5,12-disubstituted thioether-ester crown ethers and to screen for possible antibacterial activities. The efficient synthetic approach disclosed herein that led to quick output of a series of 1,4-dioxo-7,10-dithiacyclododecane-2,3-diones for the evaluation of antibacterial activities of these compounds indicated that only (**5e**) and (**5f**) can be considered the inhibitors for *S. aureus* methicillin resistance and *P. aeruginosa*.

EXPERIMENTAL

¹H NMR (500 MHz) and ¹³C NMR (125 MHz) were obtained by using a Bruker Avance DRX-500 fourier transformer spectrometer. Chemical shifts are reported in ppm (δ) downfield from tetramethylsilane (TMS). Electron impact mass spectra were recorded on a Varian Match 7A spectrometer. All chemicals were purchased from Merck and Fluka Co. and used without further purification.

General Procedure for the Synthesis of 1,4-dioxo-7,10-dithiacyclododecane-2,3-diones (5a-j)

A mixture of β,β' -dihydroxydithioether (3.0 mmoles) and triethylamine (0.65 g, 6.4 mmoles) in dried dichloromethane (10 mL) was cooled to 5–10°C. Oxalyl chloride (0.40 g, 3.1 mmoles) in dried dichloromethane (10 mL) was then added in one portion to the mixture while the mixture was allowed to warm to r. t. After 30 min, the mixture was poured into cold water (100 mL) and acidified with 5% HCl (20 mL). The aqueous solution was extracted with dichloromethane (2×30 mL). The combined organic extracts were washed with 5% NaHCO_3 (2×30 mL) and then brine (2×30 mL), dried with Na_2SO_4 , and concentrated under reduced pressure to provide the desired compounds. The purity of the compounds was checked on TLC (silica gel 60 F₂₅₄, dichloromethane-methanol 9:1).

General Procedure for Minimum Inhibitory Concentration

The Minimum Inhibitory Concentrations (MICs) of (3a-j) and (5a-j) were determined in the dilution tube-test method, which had been introduced by the National Committee for Clinical Laboratory Standards.¹⁰ For both dilution methods, decreasing concentrations of the antimicrobial agents to be tested, usually prepared in serial twofold dilution, were placed in tubes of a broth medium that supported the growth of the test microorganism. After sufficient incubation (usually overnight), the tubes were examined for turbidity, indicating growth of the microorganism. The organism grew in the tube that did not contain enough antimicrobial agent to inhibit growth. The lowest drug concentration of the agent that prevented growth of the test organism, as detected by lack of visual turbidity (matching the negative growth control), was designated the MIC. A serial dilution of the tested compounds (400 to 25 $\mu\text{g/mL}$) were added to the test bacteria in the Mueller-Hinton broth and were incubated at 37°C for 18–20 h. Growth was presented in the medium control and was absent from the inoculum control.¹¹

1,4-Dioxo-7,10-dithiacyclododecane-2,3-dione (5a)

White solid (54%), m.p.: 112°C; ^1H NMR: δ 2.84 (s, 4H, $-\text{SCH}_2\text{CH}_2\text{S}-$), 2.88 (t, $J=8$ Hz, 4H, $-\text{SCH}_2-$), 4.45 (t, $J=8$ Hz, 4H, $-\text{CH}_2\text{OCO}$); ^{13}C NMR: δ 30.8 ($-\text{SCH}_2\text{CH}_2\text{S}-$), 32.6 ($-\text{SCH}_2-$), 68.3 ($-\text{CH}_2\text{OCO}$), 158.6 (C=O); MS m/z : 238 (M^+), 148 (100%).

5,12-Dimethyl-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5b)

White viscous liquid (65%); ^1H NMR: δ 1.45 (d, $J=8$ Hz, 6H, $-\text{CH}_3$), 2.79–2.89 (m, 4H, $-\text{SCH}_2-$), 2.81 (s, 4H, $-\text{SCH}_2\text{CH}_2\text{S}-$), 5.13 (m, 2H,

CH-OCO); ^{13}C NMR: δ 21.9 (-CH₃), 32.3 (-SCH₂CH₂S-), 35.5 (-SCH₂-), 70.3 (CH-OCO), 157.7 (C=O); MS m/z : 264 (M⁺), 118 (100%).

5,12-Diethyl-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5c)

White viscous liquid (69%); ^1H NMR: δ 1.00 (t, J = 8 Hz, 6H, CH₃), 1.84 (m, 4H, -CH₂-), 2.78–2.87 (m, 4H, -SCH₂-), 2.82 (s, 4H, -SCH₂CH₂S-), 5.10 (m, 2H, CH-OCO); ^{13}C NMR: δ 9.8 (CH₃), 28.8 (-CH₂-), 32.3 (-SCH₂CH₂S-), 35.4 (-SCH₂-), 70.7 (CH-OCO), 157.5 (C=O); MS m/z : 292 (M⁺), 132 (100%).

5,12-Diphenyl-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5d)

Yellow solid (70%), m.p.: 71–73°C; ^1H NMR: δ 2.84–2.93 (m, 4H, -SCH₂-), 2.85 (s, 4H, -SCH₂CH₂S-), 5.97 (m, 2H, CH-OCO), 7.14–7.34 (m, 10H, Ar-H); ^{13}C NMR: δ 32.7 (-SCH₂CH₂S-), 38.1 (-SCH₂-), 76.6 (CH-OCO), 127.5, 129.2, 130.2 and 137.7 (aromatic carbons), 157.7 (C=O); MS m/z : 388 (M⁺), 120 (100%).

5,12-Di((allyloxy)methyl)-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5e)

White viscous liquid (68%); ^1H NMR: δ 2.79–2.88 (m, 4H, -SCH₂-), 2.81 (s, 4H, -SCH₂CH₂S-), 3.72 (m, 4H, -OCH₂-), 4.06 (d, J = 6 Hz, 4H, -CH₂O-), 5.07 (m, 2H, CH-OCO), 5.22 (dd, J = 10 Hz, 1.2 Hz, 2H, *cis*), 5.30 (dd, J = 17 Hz, 1.2 Hz, 2H, *trans*), 5.86–5.94 (m, 2H, -CH=); ^{13}C NMR: δ 33.1 (-SCH₂CH₂S-), 36.0 (-SCH₂-), 69.3 (-CH₂O-), 70.1 (-OCH₂), 72.9 (CH-OCO), 118.0 (=CH₂), 134.7 (-CH=), 158.3 (C=O); MS m/z : 376 (M⁺), 174 (100%).

5,12-Di(isopropoxymethyl)-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5f)

White viscous liquid (70%); ^1H NMR: δ 1.21 (d, J = 6 Hz, 6H, -CH₃), 2.80–2.89 (m, 4H, -SCH₂-), 2.84 (s, 4H, -SCH₂CH₂S-), 3.63 (m, 2H, -O-CH), 3.71 (m, 4H, -CH₂O-), 5.20 (m, 2H, CH-OCO); ^{13}C NMR: δ 22.4 (-CH₃), 33.1 (-SCH₂CH₂S-), 36.1 (-SCH₂-), 70.1 (-O-CH), 71.0 (-CH₂O-), 72.7 (CH-OCO), 158.2 (C=O); MS m/z : 388 (M⁺), 120 (100%).

5,12-Di(butoxymethyl)-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5g)

White viscous liquid (51%); ^1H NMR: δ = 0.89 (t, J = 8 Hz, 6H, -CH₃), 1.18–1.69 (m, 8H, -CH₂CH₂-), 2.75–2.91 (m, 4H, -SCH₂-), 2.81 (s, 4H, -SCH₂CH₂S-), 3.47 (t, J = 8 Hz, 4H, -OCH₂-), 3.70 (m, 4H, -CH₂O-), 5.15 (m, 2H, CH-OCO); ^{13}C NMR: δ 14.4 (-CH₃), 19.3 (-CH₂-), 31.2 (-CH₂-),

33.1 (-SCH₂CH₂S-), 36.0 (-SCH₂-), 70.3 (-OCH₂-), 71.1 (-CH₂O-), 72.6 (CH-OCO), 158.1 (C=O); MS *m/z*: 408 (M⁺), 190 (100%).

5,12-Di(tert-butoxymethyl)-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5h)

White viscous liquid (53%); ¹H NMR: δ 1.17 (s, 18H, -CH₃), 2.80–2.90 (m, 4H, -SCH₂-), 2.81 (s, 4H, -SCH₂CH₂S-), 3.67 (m, 4H, -CH₂O-), 5.12 (m, 2H, CH-OCO); ¹³CNMR δ 27.4 (-CH₃), 33.2 (-SCH₂CH₂S-), 36.0 (-SCH₂-), 70.4 (-O-C), 71.3 (-CH₂O-), 72.8 (CH-OCO), 158.2 (C=O); MS *m/z*: 408 (M⁺), 190 (100%).

5,12-Di(phenoxyethyl)-1,4-dioxo-7,10-dithiacyclododecane-2,3-dione (5i)

White solid (63%), m.p.: 155–157°C; ¹H NMR: δ 2.78–2.91 (m, 4H, -SCH₂-), 2.81 (s, 4H, -SCH₂CH₂S-), 4.25 (m, 4H, -CH₂O-), 5.35 (m, 2H, CH-OCO), 6.77–7.30 (m, 10H, Ar-H); ¹³CNMR: δ 33.1 (-SCH₂CH₂S-), 36.3 (-SCH₂-), 71.9 (-CH₂O-), 73.5 (CH-OCO), 158.5 (C=O); MS *m/z*: 448 (M⁺), 210 (100%).

Perhydrodibenzo[e,k][1,4,7,10]dioxadithiacyclododecane-6,7-dione (5j)

White solid (64%), m.p.: 82–84°C; ¹H NMR: δ 1.29–1.62 (m, 8H, -CH₂CH₂-), 1.76 (m, 4H, -CH₂-), 2.14 (m, 4H, -CH₂-), 2.79–2.94 (m, 6H, -SCH- and -SCH₂CH₂S-), 4.88 (m, 2H, CH-OCO), ¹³CNMR: δ 24.8 (-CH₂-), 26.4 (-CH₂-), 30.7 and 31.9 (-CH₂-), 32.5 and 33.7 (-CH₂-), 33.4 and 34.5 (-SCH₂-CH₂S-), 47.4 and 48.0 (-SCH-), 79.4 and 80.0 (CH-OCO), 157.3 and 157.5 (C=O); MS *m/z*: 344 (M⁺), 158 (100%).

REFERENCES

- [1] M. R. Younes, M. M. Chaabouni, and A. Baklouti, *Tetrahedron Letters*, **42**, 3167 (2001).
- [2] V. Guyon, A. Guy, J. Foos, M. Lemaire, and M. Draye, *Tetrahedron*, **51**, 4065 (1995).
- [3] G. W. Gokel, W. M. Leevy, and E. M. Weber, *Chem. Rev.*, **104**, 2723 (2004).
- [4] S. M. Seyedi, M. Shadkam, and A. Ziafati, *Phosphorus, Sulfur, and Silicon*, **180**, 1953 (2005).
- [5] L. M. Prescott, J. P. Harley, and D. A. Klein. In *Microbiology*, 5th ed., pp. 809 (McGraw-Hill, New York, 2002).
- [6] M. P. Jevsons, A. V. Coe, and M. T. Parker, *Lancet*, **1**, 904 (1963).
- [7] M. P. Jevsons and M. T. Parker, *J. Clin. Pathol.*, **17**, 243 (1964).
- [8] T. K. McKinney, V. K. Sharma, W. A. Craig, and G. L. Archer, *J. Bacteriol.*, **183**, 6862 (2001).
- [9] M. E. Mulligan, K. A. Murray-Leisure, B. S. Ribner, H. C. Standiford, J. F. John, J. A. Korvick, C. A. Kauffman, and V. L. Yu, *Am. J. Med.*, **94**, 313 (1993).

- [10] S. M. Finegold and L. Garrod, In *Bailey & Scott's Diagnostic Microbiology*, 8th ed., pp. 171–193 (C.V. Mosby Co., Toronto, ONT, 1995).
- [11] L. Phillips, J. D. Willians, and R. Wise, In *Laboratory Methods in Antimicrobial Chemotherapy*, p. 3 (Churchill Livingstone, Edinburgh, 1978).