


275. Cyclic Conjugated Polyenes. Part IV. Attempted Synthesis of 6-Methyl-4:5-benzazulene.


By J. R. NUNN.

An attempt to prepare 6-methyl-4:5-benzazulene by the dehydrogenation of 6-methyl-1:2:3:8:9:10-hexahydro-4:5-benzazulene is reported. This attempt failed but, instead, the latter hydrocarbon, on dehydrogenation, rearranged to 9-methylphenanthrene.


RECENT attempts (Nunn and Rapson, *J.*, 1949, 1051) to isolate 4:5-benzazulene (I) in a pure state failed owing to instability. However, the molecular complexes of the hydrocarbon with 1:3:5-trinitrobenzene and 2:4:6-trinitrotoluene were isolated as reasonably stable compounds, presumably because of polarisation effects associated with complex formation. In an attempt to assess the factors which may control stability in the 4:5-benzazulene structure, an attempt has been made to prepare 6-methyl-4:5-benzazulene (II).



(I; R = H.)
(II; R = Me.)



(III.)



(IV.)

6-Hydroxy-6-methyloctahydro-4:5-benzazulene (III) was made by the action of methylmagnesium iodide on 6-keto-octahydro-4:5-benzazulene in the usual way, and was dehydrated to 6-methyl-1:2:3:8:9:10-hexahydro-4:5-benzazulene (IV) by heating it with potassium hydrogen sulphate. This was then dehydrogenated in the vapour phase in a vacuum. No conditions were found, however, under which the expected 6-methyl-4:5-benzazulene was produced and in all reactions the isomeric 9-methylphenanthrene was formed.

Substituted 4:5-benzazulenes are therefore not always accessible through dehydrogenation of hydro-derivatives, owing to the ease of rearrangement to yield derivatives of phenanthrene.

EXPERIMENTAL.

6-Methyl-1:2:3:8:9:10-hexahydro-4:5-benzazulene (IV).—Methylmagnesium iodide, from magnesium (0.97 g., 0.04 mol.) and methyl iodide (5.52 g., 0.039 mol.), in anhydrous ether was cooled in ice, and 6-keto-octahydro-4:5-benzazulene (4.0 g., 0.02 mol.) (Nunn and Rapson, *loc. cit.*) in a little anhydrous ether was added slowly with vigorous stirring. After 1 hour's refluxing the solution was set aside overnight. The complex was decomposed with ice-cold dilute sulphuric acid, and the ethereal layer was separated and washed with sodium hydrogen sulphite solution and water. Distillation of the dry neutral extract afforded 6-hydroxy-6-methyloctahydro-4:5-benzazulene as a colourless, very viscous oil (4.1 g., 95%), b. p. 123—125°/0.4 mm.

This carbinol (1.5 g.), mixed thoroughly with finely powdered potassium hydrogen sulphate (1.5 g.), was heated at 180—190° for 1.5 hours. The mixture was extracted with ether, and the ethereal solution washed with sodium carbonate solution (10%) and water. Distillation gave 6-methyl-1:2:3:8:9:10-hexahydro-4:5-benzazulene as a colourless mobile oil, b. p. 92—94°/0.3 mm., n_D^{25} 1.5769, d_4^{25} 1.024, $[R]_D^{25}$ = 64.22 (Calc.: 63.00) (Found: C, 90.4; H, 9.1. C₁₅H₁₆ requires C, 90.9; H, 9.1%).

Dehydrogenation. The hexahydro-derivative was dehydrogenated in the vapour phase at 22 mm. over 30% palladium-charcoal, using an apparatus previously described (Nunn and Rapson, *J.*, 1949, 825). The hydrocarbon (0.80 g.) was passed through the catalyst, which was maintained at 340°, at the rate of 0.01 c.c./min. The dark distillate (0.395 g.) was dissolved in pentane (60 c.c.) and chromatographed on alumina (60 g.; Peter Spence Type H) in a column 25.5 × 1.87 cm. Fractions were collected as indicated in the table.

Fraction.	Eluent.	Eluate, c.c.	Residue, mg.	Remarks.
1—8	Pentane	420	275.9	Colourless mobile oil.
9	Pentane-benzene (9:1)	100	27.9	Colourless viscous oil.
10	"	100	21.3	Colourless viscous oil.
11	Pentane-benzene (4:1)	100	30.9	Colourless crystals. Deep-blue fluorescence in ultra-violet light.
12	"	100	9.6	Green oil.

Fraction 11 was again chromatographed on a column containing 5 g. of alumina with pentane only as the eluent. This afforded 24 mg. of crystalline material which, after 2 crystallisations from methanol, gave colourless prisms. These were sublimed at 85° (block temp.) at 0.01 mm. and then had m. p. 90—91° (Found: C, 93.7; H, 6.7. Calc. for C₁₅H₁₄: C, 93.7; H, 6.3%). The picrate formed orange-yellow needles, m. p. 151—153°, from methanol. Windaus, Jensen, and Schramme (*Ber.*, 1924, 57, 1875) give m. p. 91—92° for 9-methylphenanthrene and m. p. 152—153° for the picrate.

[1950] *A Synthesis of β -Anhydrotrimethylbrazilone.* 1353

Fractions 1—8 proved to be mainly unchanged starting material. Further dehydrogenations were carried out at 270°, 290°, 310°, and 360° without any of the desired product being obtained.

The author thanks Dr. W. S. Rapson for advice during this work and the South African Council for Scientific and Industrial Research for permission to publish this paper.

NATIONAL CHEMICAL RESEARCH LABORATORY,
PRETORIA, SOUTH AFRICA.

[Received, February 25th, 1950.]
