## PALLADIUM-PROMOTED AROMATIC ANNELATION WITH ACETYLENE DICARBOXYLATES

Tsutomu SAKAKIBARA,<sup>\*</sup> Yasuo TANAKA, and Shin-ichi YAMASAKI Institute of Chemistry, College of Liberal Arts, Kagoshima University, Korimoto, Kagoshima 890

Benzene rings of N-alkyltoluidine and benzene (or iodobenzene) in acetic acid were annelated with acetylenedicarboxylates by palladium(II) acetate to give N-alkylindole-2,3-dicarboxylate and naphthalene-1,2,3,4-tetracarboxylate. The formation of the products are explained in terms of toluidine radical cation intermediates.

Recent work elucidates that N,N-dialkylanilines react with Pd(II) compounds to give the radical cation intermediates which may lead to interesting oxidation products.<sup>1)</sup> On the other hand, many palladium-catalyzed intramolecular cyclizations which are usually explained in terms of polar mechanisms in nature, have been reported to convert functionalized anilines into aromatic fused heterocyclic compounds.<sup>2)</sup> We now report novel palladium-promoted annelation of benzene (or iodobenzene) and N-alkylanilines with acetylenedicarboxylates to give naphthalene-1,2,3, 4-tetracarboxylates and 1-alkylindole-2,3-dicarboxylates, respectively. Since onepot formation of naphthalenes by annelation of acetylenes onto benzene rings has been hitherto unknown,<sup>3)</sup> some related reactions were also investigated from mechanistic view-points. The reactions are reasonably explained in terms of homolytic process of toluidine radical cation intermediates.

$$\begin{array}{c} R_{1} & & & R_{2} & R_{1} & R_{2} & R_{2} & R_{1} & R_{2} & R_{2} & R_{2} & R_{2} & R_{2} & R_{2} & R_$$



Table 1. Aromatic Annelation with Acetylene Dicarboxy1	ſable	1. Aromatic	Annelation	with	Acetylene	Dicarboxylat
--	-------	-------------	------------	------	-----------	--------------

Run		Aromatics (mmol)	Alkyne (mmol )	Metals (mmol )		Product	S	
1.	la,	Benzene	2a	Pd(II) <sup>a)</sup>	3a,	24% <sup>b)</sup>	(	24%) <sup>c)</sup>
(]	(10)	(900)	(10)	(10)	<u>4a</u> ,	5.8%	(	5.8%)
					<u>5a</u> ,	24%	(	12%)
2. (	<u>lb</u> ,	Benzene (900)	$(\frac{2a}{10})$	Pd(II) <sup>a)</sup>	<u>3b</u> ,	22%	(	22%)
	(10)			(10)	4b,	26%	(	26%)
					<u>5a</u> ,	2.0%	(	1.0%)
3. (4	<u>la</u> ,	Benzene	$(\frac{2b}{4.8})$	Pd(II) <sup>a)</sup>	3c,	3.2%	(	3.2%)
	$(\overline{4.8})$	(450)		(4.8)	4c,	1.0%	(	1.0%)
					<u>5b</u> ,	39%	(	20%)
4. $(\frac{1a}{10})$	la,	Benzene, Iodobenzene	$(\frac{2a}{5.0})$	$Pd(II)^{a,d}$	Зa,	7.3%	(	73%)
	$(\overline{10})$	(450) (22.5)		(0.5)	4a,	trace	(	1.0%)
					<u>5a</u> ,	43%	(	220%)
5. (	la,	Toluene, Iodobenzene	$(\frac{2a}{10})$	Pd(II) <sup>a,d)</sup>	3a,	3.0%	(	30%)
	$(\overline{10})$	(380) (45)		(1.0)	4a,	9.5%	(	94%)
					<u>5a</u> ,	17%	(	86%)
6.	Benze	ene	2a	Pd(II) <sup>a)</sup>	5a	trace	(1	trace)
	(450	(450)		(5.0)	<u>6a</u> ,	31%	(	10%)
7. E	Benze	Benzene, Iodobenzene (450) (10)		$Pd(II)^{a}$	5a,	4.5%	(	23%)
	(450			(0.5)	6a,	1.4%	(	4.5%)
					<u>7a</u> ,	8.9%	(	30%)
8. (	la,	enzene	$(\frac{2a}{10})$	Pd(II) <sup>a)</sup> /Cu(II) <sup>e)</sup>	3a,	5.3%	(	26%)
	$(\overline{10})$	(1700)		(2.0) (10)	4a,	0.8%	(	4.1%)
					<u>5a</u> ,	4.2%	(	11%)

a)  $Pd(OAc)_2$  and acetic acid (40 ml) as a solvent were used.

b) Yields are based on  $\underline{2}$ .

c) Yields are based on Pd(II).

d) Triphenylphosphine and triethylamine were added.
[Pd(II): P(Ph)<sub>3</sub>:NEt<sub>3</sub>=1:2:7]

e)  $Cu(OAc)_2$  was used.

In a typical experiment, a mixture of N-methyl-p-toluidine (la, 10 mmol), benzene (900 mmol), dimethyl acetylenedicarboxylate (2a, 10 mmol), and Pd(OAc), (10 mmol) in acetic acid (40 ml) was stirred at 60 °C for 5 h under an atmosphere of nitrogen. After usual work-up, 2.35 mmol of dimethyl 1,5-dimethylindole-2,3-dicarboxylate (3a) and 1.20 mmol of tetramethyl naphthalene-1,2,3,4-tetracarboxylate (5a) were isolated by silica gel column-chromatography, together with 0.58 mmol of dimethyl N-methyl-N-(p-tolyl)aminomaleate (4a) and 1.84 mmol of recovered lb. The use of N-ethyl-p-toluidine (1b) or diethyl acetylenedicarboxylate (2b) in place of <u>la</u> or <u>2a</u> gave the corresponding products as were summarized in Table 1. The typical 3 and 5 were characterized as follows; 3a: IR (KBr) 1700, 1740 cm<sup>-1</sup> (-COOCH<sub>3</sub>). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) & 7.82 (s, 1H), 7.45-7.07(m, 2H), 3.90(s, 3H), 3.82(s, 3H), 3.67(s, 3H), 2.37(s, 3H). <sup>13</sup>C-NMR (CDCl<sub>3</sub>, TMS)  $\delta$  164.7(s), 163.3(s), 135.4(s), 134.4(s), 132.3(s), 126.3(d), 125.5(s), 121.8(d), 109.8(d), 107.7(s), 52.9(q), 51.4 (q), 31.4(q), 21.5(q). MS m/e 261(M<sup>+</sup>), 230, 171. Found: C, 64.20; H, 5.77; N, 5.27. Calcd for  $C_{14}H_{15}NO_4$ : C, 64.37; H, 5.75; N, 5.36. <u>5a</u>: IR (KBr) 1740 cm<sup>-1</sup> (-COOCH<sub>3</sub>). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) & 7.57-8.23 (m, 4H), 3.93 and 4.03(s, 6H, resp.). MS  $m/e 360(M^{+})$ , 329. Found: 360.0825. Calcd for  $C_{18}H_{16}O_8$ : 360.0845.

Iodobenzene is known to be an effective reoxidant of metallic palladium.<sup>4)</sup> The aromatic annelation catalytically proceeded in the presence of iodobenzene (runs 4 and 5). Therein, the improved yields of 5a may show that iodobenzene also should be one of reactants leading to 5a. Moreover, the reactions of 2a with benzene or iodobenzene in the absence of <u>1</u> resulted in formation of various cyclized oligomers of 2a, such as  $6a^{5)}$  or 7a, 6 along with relatively poor yields of 5a. The results suggest that 5a might be derived from a common intermediate with that of <u>3</u>. The metallocyclic intermediates shown in olefin annelation by  $2^{7}$  may not be important concerning the formation of 5.

The possibility of intramolecular cyclization of 4 as an intermediate to give 3 was investigated. The reaction of 4a with an equimolar amount of Pd(OAc), at 60° was relatively slow in a mixed solvent of acetic acid and benzene, and only after 24 h gave 48.6 % of 3a (based on consumed 4a; the conversion of 4a, 83%). Furthermore, a similar reaction of 4a with a catalytic amount of Pd(OAc)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> complex [4a:Pd(II)=25:1] at 90 °C for 20  $h^{8}$  formed 44.6% of 1a (conversion of 4a, 89%), together with some oligomers of 4a, but neither 3a nor 5a. Thus, the formation of <u>3a</u> in the present aniline annelation by 2a may not be completely attributed to <u>4a</u>. It has been already elucidated that the reaction of N,N-dialkylanilines with palladium(II) compounds may involve a single electron transfer process.<sup>1</sup>/ A similar SET process between 1 and Pd(II) species might generate the radical cation intermediate of 1 which could react with 2 to form vinyl radicals (8) followed by intramolecular aromatic substitution to produce  $\underline{3}$ . A possible mechanism for the formation of 3 and 5 is tentatively shown in Scheme 1. Therein, the product 5 is explained in terms of homolytic addition and substitution by  $\underline{8}$ , and subsequent thermal electrocyclization of conjugated aromatic diene (10). Copper(II) salts are known to be an effective radical scavenger.<sup>9)</sup> The suppression of the aromatic annelation in the presence of Cu(II) (run 8), in spite of its ability to reoxidize metallic palladium, may support the homolytic process.



This work has been partially supported by the Grant-in-Aid for Scientific Research (No. 59550582) from the Ministry of Education, Science and Culture, Japan.

## References

- 1) T. Sakakibara, S. Karasumaru, and I. Kawano, J. Am. Chem. Soc., <u>107</u>, 6417 (1985).
- M. Mori and Y. Ban, Tetrahedron Lett., <u>1976</u>, 1803, 1807; M. Mori, K. Chiba, and Y. Ban, ibid., <u>1977</u>, 1037; L. S. Hegedus, G. F. Allen, J. J. Bozell, and E. L. Waterman, J. Am. Chem. Soc., <u>100</u>, 5800 (1978); M. O. Terpk and R. F. Heck, ibid., <u>101</u>, 5281 (1979); S. E. Diamond, A, Szalkiewicz, and F. Mares, ibid., <u>490</u> (1979); L. S. Hegedus, G. F. Allen, and D. J. Olsen, ibid., <u>102</u>, 3583 (1980).
- S. G. Davies, "Organotransition Metal Chemistry. Application to Organic Synthesis," Pergamon Press (1982), pp. 255-260.
- 4) B. A. Patel, C. B. Ziegler, N. A. Cortese, J. E. Plevyyak, T. C. Zebovitz,
  M. Terpko, and R. F. Heck, J. Org. Chem., <u>42</u>, 3909 (1977); N. A. Cortese, C.
  B. Ziegler, Jr., B. J. Hornjez, and R. F. Heck, ibid., <u>43</u>, 2952 (1978).
- 5) <u>6a</u>: mp 152-154 °C, IR(KBr) 760, 700 cm<sup>-1</sup> (-Ph), <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 3.48 and 3.90 (s, 6H, -CH<sub>3</sub>, resp.), 3.87 (s, 3H, -CH<sub>3</sub>), 7.03-7.50 (m, 5H, Ar-H), MS m/e 444(M<sup>+</sup>), 413, 381.
- 6) <u>7a</u>: IR(KBr) 1745 cm<sup>-1</sup> (-CO<sub>2</sub>CH<sub>3</sub>), <sup>1</sup>H-MNR (CDCl<sub>3</sub>)  $\delta$  3.83 (s, 18H, -CH<sub>3</sub>), MS m/e 427(M<sup>+</sup>), 396, 395.
- 7) H. Suzuki, K. Ito, Y. Ishii, K. Simon, and J. A. Ibers, J. Am. Chem. Soc., <u>98</u>, 8494 (1976); L. D. Brown, K. Ito, H. Suzuki, K. Hirai, and J. A. Ibers, ibid., <u>100</u>, 8282 (1978).
- 8) Iodobenzene was added (Pd:PhI=1:100).
- 9) N. Kornblum, R. E. Michel, and R. C. Kerber, J. Am. Chem. Soc., <u>88</u>, 5662 (1962). ( Received March 7, 1986 )

800