

Structural Change of Pheophorbide *a* Methyl Ester by Contact with Titanium Oxide Particles

Yoshitaka Saga* and Tae Otono

Department of Chemistry, Faculty of Science and Engineering, Kinki University,
Higashi-Osaka, Osaka 577-8502

(Received December 9, 2011; CL-111181; E-mail: saga@chem.kindai.ac.jp)

Pheophorbide *a* methyl ester, which was derived from natural chlorophyll *a*, was converted to purpurin-18 methyl ester by contact with titanium oxide particles in the presence of oxygen under mild conditions in the dark.

Utilization of solar energy is a promising strategy to overcome energy and environmental crises. From this viewpoint, dye-sensitized solar cells (DSSCs) have attracted much attention recently,^{1–3} since they can be produced with low-cost and low-energy processes. To date, various photosensitizers for DSSCs have been developed.^{4–6} Among them, ruthenium complexes were reported to function as good photosensitizers of DSSCs.^{6–9} Organic dyes have also been employed as DSSC photosensitizers that are free from rare metals.^{6,10–13} However, many organic dyes can hardly absorb near-infrared light, resulting in the limitation of application to DSSCs.

Chlorophylls (Chls), which are abundant photosynthetic pigments in nature, and their derivatives have been regarded as potential photosensitizers in DSSCs,^{4–6,14–19} since these molecules can capture photons in visible and near-infrared regions. In addition to the advantage of such spectral properties, Chls can be easily prepared from higher plants and algae. The molecular structure of Chl *a* is shown in Figure 1A. Chl *a* possesses central magnesium and a phytyl chain at the 17-propionate. Since naturally occurring Chl *a* is unstable, stable Chl derivatives, which are modified from Chl *a*, have been widely examined as photosensitizers of DSSCs.^{14–19} However, little attention has been paid to molecular states of Chl derivatives on TiO₂ electrodes and their potential denaturation. In the present study,

we report a unique structural change of a Chl derivative by contact with TiO₂ particles in the presence of molecular oxygen in the dark.

Pheophorbide *a* methyl ester (**1**) was used as a model compound of Chl derivatives in this study, because of their stability and easy preparation. The molecular structure of **1** is shown in Figure 1A. Structural differences between **1** and natural Chl *a* are central atoms in the chlorin macrocycle and esterifying chains: **1** lacks central magnesium (M = 2H in Figure 1A) and has methyl ester (R = CH₃ in Figure 1A) instead of phytyl ester in Chl *a*. Pheophorbide *a* methyl ester (**1**) was synthesized from Chl *a*, which was extracted from a cyanobacterium *Spirulina geitleri*, through two steps according to previous reports.^{20,21} Anatase-type TiO₂ particles, which were purchased from Wako Pure Chemical Industries, Ltd. (diameter; 5 μm, purity; 99.9%), were added to a CH₂Cl₂ solution of **1**, and stirred for 4 h at room temperature in the dark. Figure 2 depicts spectral changes of supernatants of the CH₂Cl₂ suspension of **1** containing TiO₂ particles. Soret and Q_y absorption bands of **1**




Figure 1. (A) Molecular structures of Chl *a* (M = Mg, R = phytyl) and pheophorbide *a* methyl ester (**1**, M = 2H, R = CH₃). (B) Molecular structure of purpurin-18 methyl ester (**2**).




Figure 2. Visible absorption spectra of supernatants of CH₂Cl₂ suspensions of pheophorbide *a* methyl ester (**1**) by bubbling air for 4 h (solid curve), by bubbling N₂ for 4 h (broken curve), and intact **1** in CH₂Cl₂ (dotted curve). These spectra were normalized at Soret absorption bands. Insert: overlay of three spectra between 680 and 740 nm.




Figure 3. HPLC chromatogram of reaction products in the supernatant of a CH_2Cl_2 suspension of pheophorbide *a* methyl ester (**1**) by bubbling air for 4 h. The products were eluted on 5C₁₈-AR-II (6 mm ϕ $\times$ 250 mm) with methanol at a flow rate of 1.0 mL min⁻¹. The fraction denoted by # was the 700-nm absorbing compound.

were positioned at 413 and 667 nm, respectively, before stirring with TiO_2 particles. When the suspension of **1** and TiO_2 particles was bubbled with air for 4 h, a new Q_y absorption band appeared at 700 nm. In contrast, the absorption band around 700 nm was hardly observed by bubbling nitrogen gas for 4 h. No spectral change of **1** was detected by bubbling air without TiO_2 particles (data not shown). These indicate that the 700-nm absorbing product was generated from **1** in the presence of TiO_2 particles and molecular oxygen.

The supernatant of the CH_2Cl_2 suspension of **1** and TiO_2 particles by bubbling air for 4 h was analyzed by reverse-phase HPLC as shown in Figure 3. Fractions at 16.5 and 17.4 min were assigned to unreacted pheophorbide *a* methyl ester (**1**) and the 13²-stereoisomer of **1**, namely pheophorbide *a'* methyl ester, respectively. A new fraction, which had Soret and Q_y bands at 407 and 697 nm, respectively, in the HPLC eluent, appeared at 28.5 min (denoted by # in Figure 3). Other fractions at 10.5 and 32.3 min exhibited Q_y absorption bands around 665 nm. Therefore, the compound eluted at 28.5 min was the origin of the 700-nm absorption band formed by incubation of **1** with TiO_2 particles.

The 700-nm absorbing compound, which was detected at 28.5 min in the chromatogram in Figure 3, was isolated by reverse-phase HPLC. The purified compound was characterized by means of UV-vis, NMR, ESI-MS, and FT-IR measurements.²² In addition, the spectroscopic data of this product were compared with those of purpurin-18 methyl ester (**2**) (Figure 1B) that was conventionally synthesized under an alkaline condition with molecular oxygen according to previous reports.^{23–25}

The compound purified from the supernatant of **1** after treatment with TiO_2 particles exhibited Soret and Q_y bands at 411 and 700 nm, respectively, in CH_2Cl_2 . The Q_y band was red-shifted by 33 nm compared with **1**, and was characteristic of purpurin-18-type pigments, which had a six-membered anhydride exo-ring instead of an original five-membered exo-ring in **1**. This compound showed the molecular-ion peak at m/z 579.4 in the ESI-MS,²² which was identical to the molecular weight of purpurin-18 methyl ester. In the ¹H NMR spectrum of the




Figure 4. FT-IR spectra of the 700-nm absorbing compound purified from products after incubation of pheophorbide *a* methyl ester (**1**) with TiO_2 particles in CH_2Cl_2 by bubbling air (A), synthesized purpurin-18 methyl ester (**2**) (B), and pheophorbide *a* methyl ester (**1**) (C) in the region between 1500 and 1850 cm⁻¹.

purified compound, two singlet peaks at 6.28 and 3.88 ppm, which were assigned to protons at the 13²-position (1H) and in the 13⁴-methoxycarbonyl group (3H), respectively, of original **1**, disappeared (Figure S1).²² All the signals in ¹H and ¹³C NMR spectra of the purified compound corresponded with those of purpurin-18 methyl ester (**2**) by conventional synthesis.²² Figure 4 shows expanded FT-IR spectra of the purified compound and the synthesized **2** as well as pheophorbide *a* methyl ester (**1**) in the wavenumber region between 1500 and 1850 cm⁻¹. The purified compound exhibited no stretching vibrational peak of the 13¹-carbonyl group, which was clearly detected at 1701 cm⁻¹ in the FT-IR spectrum of **1**.²⁶ New vibrational bands were observed at 1713 and 1730 cm⁻¹ in the IR spectrum of the purified compound instead. These bands were identical to those of the synthesized **2**, and were ascribable to the anhydride group of purpurin-18 type pigments. The vibrational band at 1603 cm⁻¹, which was assigned to a skeletal vibration characteristic of the chlorin macrocycle, of the purified compound was present at the same position as that of the synthesized **2**, and was shifted to lower wavenumber than that of **1** at 1617 cm⁻¹. The vibrational stretching band of 17³-carbonyl group was preserved at 1744 cm⁻¹ in the spectrum of the purified compound after incubation of **1** with TiO_2 particles. All the vibrational bands of the purified compounds were the same as those of the synthesized **2** (Figure S3).²² These characterizations revealed that the new compound formed by treatment of **1** with TiO_2 particles was purpurin-18 methyl ester.

The conversion efficiency from pheophorbide *a* methyl ester (**1**) to purpurin-18 methyl ester (**2**) under the present conditions for 4 h was estimated to be 32% (the average of three independent measurements). This value was smaller than the overall yield from **1** to **2** (50%) reported by Lee et al.²⁵ In the present reaction, CH_2Cl_2 was appropriate compared with other solvents such as methanol, which was used as a cosolvent




Figure 5. Visible absorption spectra of TiO₂ particles isolated from a CH₂Cl₂ suspension of pheophorbide *a* methyl ester (**1**).

in conventional synthesis of purpurin-18.^{23–25} Reuse of TiO₂ particles decreased the conversion efficiency from **1** to **2** in this reaction (data not shown), probably because the active surface of TiO₂ particles was partially covered by adsorbed pigments. The amount of **1** used in this reaction was smaller than that in conventional synthesis.^{23–25} Further studies will be required to increase the yield of **2** and the reaction scale by this method.

TiO₂ particles were collected from a CH₂Cl₂ suspension of **1** and TiO₂ particles, which was incubated for 4 h in the dark, by filtration, and washed with CH₂Cl₂. Figure 5 shows visible absorption spectra of TiO₂ particles after the incubation. The intense absorption band, which was ascribed to the Q_y band of adsorbed **1**, was observed at 672 nm. The Q_y band of **1** on TiO₂ particles was shifted to longer wavelength by 5 nm than **1** in CH₂Cl₂. Additionally, a new absorption band around 710 nm appeared, suggesting that **1** on TiO₂ particles was partially converted to purpurin-18 methyl ester.

To summarize, conversion of pheophorbide *a* methyl ester (**1**) to purpurin-18 methyl ester (**2**) by contact with TiO₂ particles in the presence of molecular oxygen was reported for the first time. The present one-step conversion from **1** to **2** under the mild condition is advantageous in conventional synthesis of purpurin-18 derivatives, in which hydrolysis of the 17-ester group takes place inevitably under strong alkaline conditions.^{23–25} Investigation of a detailed reaction mechanism is now in progress. Such conversion will be useful for construction of DSSCs using Chl derivatives as photosensitizers²⁷ as well as development of facile synthesis of purpurin-18 derivatives, which attract considerable interest as near-infrared absorbing dyes.

We thank Dr. Takashi Okubo and Dr. Toshie Minematsu, Kinki University, for visible absorption spectroscopy of TiO₂ and NMR measurements, respectively. This work was partially supported by a Research Grant from the Asahi Glass Foundation.

References and Notes

- 1 B. O'Regan, M. Grätzel, *Nature* **1991**, *353*, 737.
- 2 A. Hagfeldt, M. Grätzel, *Acc. Chem. Res.* **2000**, *33*, 269.
- 3 M. Grätzel, *Nature* **2001**, *414*, 338.
- 4 H. Imahori, T. Umeyama, S. Ito, *Acc. Chem. Res.* **2009**, *42*, 1809.
- 5 X.-F. Wang, H. Tamiaki, *Energy Environ. Sci.* **2010**, *3*, 94.
- 6 A. Hagfeldt, G. Boschloo, L. Sun, L. Kloo, H. Pettersson, *Chem. Rev.* **2010**, *110*, 6595.
- 7 P. Wang, C. Klein, R. Humphry-Baker, S. M. Zakeeruddin, M. Grätzel, *J. Am. Chem. Soc.* **2005**, *127*, 808.
- 8 Y. Chiba, A. Islam, Y. Watanabe, R. Komiya, N. Koide, L. Han, *Jpn. J. Appl. Phys.* **2006**, *45*, L638.
- 9 F. Gao, Y. Wang, D. Shi, J. Zhang, M. Wang, X. Jing, R. Humphry-Baker, P. Wang, S. M. Zakeeruddin, M. Grätzel, *J. Am. Chem. Soc.* **2008**, *130*, 10720.
- 10 N. Koumura, Z.-S. Wang, S. Mori, M. Miyashita, E. Suzuki, K. Hara, *J. Am. Chem. Soc.* **2006**, *128*, 14256.
- 11 Z.-S. Wang, Y. Cui, Y. Dan-oh, C. Kasada, A. Shinpo, K. Hara, *J. Phys. Chem. C* **2007**, *111*, 7224.
- 12 C. Li, J.-H. Yum, S.-J. Moon, A. Herrmann, F. Eickemeyer, N. G. Pschirer, P. Erk, J. Schöneboom, K. Müllen, M. Grätzel, M. K. Nazeeruddin, *ChemSusChem* **2008**, *1*, 615.
- 13 S. Ito, H. Miura, S. Uchida, M. Takata, K. Sumioka, P. Liska, P. Comte, P. Péchy, M. Grätzel, *Chem. Commun.* **2008**, 5194.
- 14 A. Kay, M. Grätzel, *J. Phys. Chem.* **1993**, *97*, 6272.
- 15 Y. Amao, Y. Yamada, *Biosens. Bioelectron.* **2007**, *22*, 1561.
- 16 M. Ikegami, M. Ozeki, Y. Kijitori, T. Miyasaka, *Electrochemistry* **2008**, *76*, 140.
- 17 X.-F. Wang, Y. Koyama, H. Nagae, Y. Wada, S.-i. Sasaki, H. Tamiaki, *J. Phys. Chem. C* **2008**, *112*, 4418.
- 18 X.-F. Wang, O. Kitao, H. Zhou, H. Tamiaki, S.-i. Sasaki, *Chem. Commun.* **2009**, 1523.
- 19 X.-F. Wang, H. Tamiaki, L. Wang, N. Tamai, O. Kitao, H. Zhou, S.-i. Sasaki, *Langmuir* **2010**, *26*, 6320.
- 20 H. Tamiaki, M. Amakawa, Y. Shimono, R. Tanikaga, A. R. Holzwarth, K. Schaffner, *Photochem. Photobiol.* **1996**, *63*, 92.
- 21 Y. Saga, H. Tamiaki, *J. Photochem. Photobiol., B* **2004**, *73*, 29.
- 22 Supporting Information is available electronically on the CSJ-Journal Web site, <http://www.csj.jp/journals/chem-lett/index.html>.
- 23 J. K. Hooper, T. W. Sery, N. Yamamoto, *Photochem. Photobiol.* **1988**, *48*, 579.
- 24 A. S. Brandis, A. N. Kozyrev, A. F. Mironov, *Tetrahedron* **1992**, *48*, 6485.
- 25 S.-J. H. Lee, N. Jagerovic, K. M. Smith, *J. Chem. Soc., Perkin Trans. 1* **1993**, 2369.
- 26 J. J. Katz, R. C. Dougherty, L. J. Boucher, in *The Chlorophylls*, ed. by L. P. Vernon, G. R. Seely, Academic Press, New York, **1966**, pp. 185–251.
- 27 X.-F. Wang, L. Wang, N. Tamai, O. Kitao, H. Tamiaki, S.-i. Sasaki, *J. Phys. Chem. C* **2011**, *115*, 24394.