

Accepted Article

Title: Conversion of γ - and δ -Keto Esters into Optically Active Lactams. Transaminases in Cascade Processes

Authors: Ángela Mourelle-Insua, Luiz Arthur Zampieri, Iván Lavandera, and Vicente Gotor-Fernández

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Adv. Synth. Catal. 10.1002/adsc.201701304

Link to VoR: http://dx.doi.org/10.1002/adsc.201701304

FULL PAPER

DOI: 10.1002/adsc.201((will be filled in by the editorial staff))

Conversion of γ - and δ -Keto Esters into Optically Active Lactams. Transaminases in Cascade Processes

Ángela Mourelle-Insua,^a Luiz Arthur Zampieri,^{a,b} Iván Lavandera^{a,*} and Vicente Gotor-Fernández,^{a,*}

- ^a Organic and Inorganic Chemistry Department, Biotechnology Institute of Asturias (IUBA), University of Oviedo, Avenida Julián Clavería 8, 33006 Oviedo (Spain).
 Corresponding authors: lavanderaivan@uniovi.es (Phone: +34 98 5103452); vicgotfer@uniovi.es (Phone: +34 98 5103454)
- ^b Actual address: Organic and Inorganic Chemistry Department, Biotechnology and Natural Products Laboratory, Universidade Federal do Ceará, Campus do Pici, Fortaleza, Ceará 60455-970 (Brazil).

Received: ((will be filled in by the editorial staff))

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/adsc.201#######. Extensive transaminase screenings, analytics, copies of HPLC chiral analyses, and ¹H, ¹³C and DEPT NMR spectra for described organic compounds are included.

Dedicated to Prof. Vicente Gotor on occasion of his 70th birthday

Abstract. A one-pot two-step enzymatic strategy has been designed for the production of optically active γ - and δ -lactams in aqueous medium under mild conditions. The approach is based on the biotransamination of ethyl or methyl keto esters bearing different alkyl or aryl substitution patterns at α -position to the ketone functionality. In this manner, the keto esters were transformed into the corresponding amino esters with excellent conversions, which underwent spontaneous cyclisation in the reaction medium without addition of external reagents.

Depending on the transaminase selectivity, both lacta 1 enantiomers can be obtained, so initial enzyme screenings were performed using commercially available and *made n house* enzymes. Reaction conditions were optimised focusing on the substrate concentration, temperature were ratio of amine donor vs acceptor. Thus, ten γ - and δ -lactams were obtained in good to high isolated yields (70-90%) we excellent selectivities (94-99%) after one or two days at 30 or 45 °C.

Keywords: Cascade reaction; Intramolecular cyclisation; Lactams; Stereoselective synthesis; Transaminase

Introduction

Lactams are pivotal compounds in organic chemistry since they are present in numerous bioactive products, also serving as valuable intermediates for more complex structures including synthetic polymers.^[1] Apart from β -lactams (azetidin-2-ones, Figure 1) that constitute the most important class of currently approved antibiotics,^[2] the syntheses of γ -lactams (pyrrolidin-2-ones) and δ -lactams (piperidin-2-ones) have attracted considerable attention in recent years,^[3] due to the presence of these nitrogen containing heterocycles in demanding biologically active compounds.



Figure 1. General structure of β -, γ - and δ -lactams.

Biocatalysis provides access to optically pure products under mild reaction conditions, including the stereoselective synthesis of lactams.^[4] Kinetic resolutions of racemic lactams *via* hydrolase-catalysed ring opening or N-acylation reactions, have gained great attention in the last decades using lactamases, lipases or esterases.^[5] However, most of the examples are focused on the preparation of optically active β lactams, and are limited to 50% yield due to the inherent limitations of classical kinetic resolutions.

Interestingly, lipases have served for the production of lactams through intramolecular cyclisation of carboxylic acid derivatives intermediates, although these approaches have been disclosed in a nonselective fashion.^[6] Transaminase-catalysed processes have also been reported for the production of lactams in a highly selective manner.^[7] This strategy is based on the biotransamination of aldo or keto esters and subsequent intramolecular cyclisation of the soobtained optically active amino esters. In this manner, the chemical equilibrium is shifted to the formation of the desired lactams. In this scenario, the usefulness of commercially available transaminases has been reported for the conversion of ethyl 4-oxo-3phenylbutyrate into (*R*)-4-phenylpyrrolidin-2-one,^[7a] isopropyl 4-(4-bromophenyl)-5-oxopentanoate into (*S*)-5-(4-bromophenyl)piperidin-2-one,^[7c] and ethyl 4acetylbutyrate in both (*S*)- and (*R*)-6-methylpiperidin-2-one.^[7b] More recently a novel reductive aminase from *Aspergillus oryzae* has been engineered and applied in the reductive amination of ethyl levulinate using three aliphatic amines, allowing the asymmetric synthesis of optically active N-substituted lactams.^[8]

Herein, we propose a general strategy for the asymmetric preparation of γ - and δ -lactams based on the biotransamination of easily accessible keto esters, followed by intramolecular cyclisation in the own reaction medium (Scheme 1). Screening and optimisation of the reaction conditions will be disclosed in order to give access to both enantiomers of a series of lactams in a one-pot two-step synthetic approach. The selection of the proper transaminase will be a key issue, affecting the reaction conditions such as pH, substrate and amine donor concentrations and temperature, to develop high-yielding and stereoselective protocols.

Results and Discussion

Levulinic acid derivatives are considered valuable lactam precursors due to their simple preparation from lignocellulosic biomass and in some cases commercial availability.^[9] For that reason ethyl levulinate (**1a**) was selected as model substrate for this biotransamination screening. A 25 mM substrate concentration was initially considered in order to test 26 commercial transaminases, all of them accepting isopropylamine (IPA) as amine donor. Initially, IPA was used in a large excess (1 M concentration), aiming to drive the equilibrium towards the formation of the amino ester **2a**, which will later spontaneously cyclise into the lactam **3a**.

For an exhaustive screening study, more data can be found in Table S3 in the Supporting Information. Based on previous studies,^[10] 2.5% of DMSO was used as cosolvent in combination with phosphate

buffer 100 mM pH 7.5. The reactions were incubated for 24 h at 30 °C, observing the solely formation of active 5-methylpyrrolidin-2-one (**3a**). optically obtained after spontaneous intramolecular cyclisation in the reaction medium of ethyl 4-aminopentanoate (2a). The best results have been summarised in Table 1 (entries 1-5), finding the best conditions for TA-P2-B01 (87% conversion, >99% ee(R), entry 2) and ATA-237 (92% conversion, >99% *ee* (S), entry 3) in the complementary synthesis of both 3a lactam antipodes. In addition, transaminases such the (S)selective ones from Chromobacterium violaceum (Cv-TA)^[11] and a variant from Arthrobacter citreus (ArS-TA),^[12] and the (R)-selective TAs from Arthrobacter species (ArR-TA)^[13] and an evolved mutant (ArRmut11-TA),^[14] were also tested as freeze-dried cell powder of *E. coli* containing the corresponding overexpressed enzyme. Particularly, an excellent activity and selectivity for ArS-TA was achieved towards the formation of (S)-3a (97% conversion, >99% ee, entry 5).

We also studied the effect of the leaving group, performing biotransamination experiments over methyl levulinate (**1b**, **R**= Me), which affords an alternative access to **3a** (entries 6-11). Conversions over 80% were reached leading to both lactam enantiomers with excellent selectivity, finding a much higher reactivity for some enzymes such as TA-P2-A07 when considering the methyl instead of the ethyl ester as starting material (compare entries 1 and 6). For a full comparison of the transaminase activity between substrates **1a** and **1b**, see Tables S3 and S4 in the Supporting Information. In summary, hits for both substrates **1a,b** were found, although in general bette conversions were achieved with the methyl keto ester derivative.

Looking for a further exploitation of the synthetic approach, the study of a substrate with an increased alkyl chain length between both carbonyl groups was considered. Thus, both enantiomers of δ -lactam 6-methylpiperidin-2-one were straightforward obtained from ethyl 5-oxohexanoate (**3c**) in a highly selective manner (>88% conversion and >96% *ee*), using different enzymatic preparations (see entries 12-18 and also Tables S3 and S4 in the Supporting Information).



Scheme 1. General transformation of γ - and δ -keto esters into optically active lactams mediated by transaminases.

Table 1. Stereoselective transformation of keto esters 1a-c into lactams 3a and 3c.

		0 () 25 m	OR OR KPi M 3	• or (<i>R</i>)-Transaminase sopropylamine (1 M) PLP (1 mM) buffer 100 mM pH 7.5 DMSO (2.5% v/v) 0 °C. 24 h. 250 rpm	() _n *		
		1a : n= 1, R= Et 1b : n= 1, R= Me 1c : n= 2, R= Et		3a : n= 1 3c : n= 2			
Entry	Keto ester	n	R	Transaminase	<i>c</i> [%] ^[a]	<i>ee</i> [%] ^[b]	
1	1a	1	Et	TA-P2-A07	41	>99 (<i>R</i>)	
2	1a	1	Et	TA-P2-B01	87	>99 (<i>R</i>)	
3	1a	1	Et	ATA-237	92	>99 (S)	
4	1a	1	Et	TA-P1-B04	89	>99 (<i>S</i>)	
5	1a	1	Et	ArS-TA	97	>99 (<i>S</i>)	
6	1b	1	Me	ATA-412	84	>99 (<i>R</i>)	
7	1b	1	Me	TA-P2-A07	81	>99 (<i>R</i>)	
8	1b	1	Me	TA-P2-B01	80	>99 (<i>R</i>)	
9	1b	1	Me	TA-P1-A06	90	>99 (<i>S</i>)	
10	1b	1	Me	TA-P1-G06	91	>99 (<i>S</i>)	
11	1b	1	Me	ArS-TA	>99	>99 (<i>S</i>)	
12	1c	2	Et	ATA-013	89	96 (<i>R</i>)	
13	1c	2	Et	ATA-033	87	97 (<i>R</i>)	
14	1c	2	Et	TA-P2-B01	86	97 (<i>R</i>)	
15	1c	2	Et	ATA-200	86	>99 (<i>S</i>)	
16	1c	2	Et	TA-P1-A06	88	>99 (<i>S</i>)	
17	1c	2	Et	TA-P1-G06	96	99 (<i>S</i>)	
19	1c	2	Et	ArS-TA	98	>99(S)	

^[a] Conversion values determined by GC (see Supporting Information). ^[b] Determined by HPLC using chiral columns (see Supporting Information). Absolute configurations of the corresponding lactams **3a** and **3c** appear in brackets.

Table 2. Stereoselective transformation of a	v-keto esters 1d-h into	5-arvlpyrrolidin-2-ones 3d-h .
	Reco esters ia in into	

Supporting Information). Absolute configurations of the corresponding lactams 3a and 3c appear in brackets.								
Table 2. S	Stereoselective tra	insformation of γ-keto este	ers 1d-h into 5-aryl	pyrrolidin-2-ones 3d	-h.		\geq	
	R	O O O Me K	S)- or (<i>R</i>)-Transamin Isopropylamine (1 f PLP (1 mM) Pi buffer 100 mM pH DMSO (2.5% v/v) 24 b, 250 rpm	ase M) 17.5 H 3d b	R		eq	
Entry	B (Vato astar)	[Kata astar] [mM]	T [°C]	Transaminasa	a [0/][a]	aa [0/][b]		
	K (Kelo ester)				<i>C</i> [%] ¹³			
1	H (Id)	25	30	ATA-303	59	>99(R)		
2	H (1d)	15	30	ATA-303	/8	>99(R)		
3	H (1d)	15	45	ATA-303	97	>99 (R)		
4	H (1d)	25	30	ATA-237	45	98 (S)		
5	H (1d)	15	30	ATA-237	57	>99 (S)		
6	H (1d)	15	45	ATA-237	57	93 (S)		
7	OMe (1e)	15	45	ATA-025	94	>99 (<i>R</i>)		
8	OMe (1e)	15	45	ATA-033	95	>99 (<i>R</i>)		
9	OMe (1e)	15	45	ATA-234	91	>99 (S)		
10	Me (1f)	15	45	ATA-025	98	>99 (<i>R</i>)		
11	Me (1f)	15	45	ATA-033	98	>99(R)		
12	Me (1f)	15	45	ATA-234	89	>99(S)		
13	F (1g)	15	45	ATA-024	99	>99 (R)		
14	F (1 g)	15	45	ATA-025	98	>99(R)		
15	F(1g)	15	45	ATA-415	99	>99(R)		
16	$F(\mathbf{1g})$	15	45	ATA-234	81	>99 (S)		
17	Et (1h)	15	45	ATA-025	83	>99(R)		
18	$\tilde{\mathbf{Et}}(\mathbf{1h})$	15	45	ATA-033	86	>99(R)		
19	Et (1h)	15	45	ATA-234	82	>99 (S)		

^[a] Conversion values determined by GC (see Supporting Information). ^[b] Determined by HPLC using chiral columns (see Supporting Information). Absolute configurations of the corresponding lactams **3d-h** appear in brackets.

Next, the enzymatic study over methyl 4-aryl-4oxobutanonates was envisaged, firstly starting from commercially available methyl 4-oxo-4phenylbutanoate (1d), and later extending the study to compounds bearing different substitution patterns at the *para*-position of the aromatic ring (OMe, Me, F and Et). These keto esters **1e-h** were obtained through esterification of the corresponding carboxylic acids with refluxing methanol in acidic media (98-99%). Subsequent reductive amination using sodium cyanoborohydride and ammonium acetate at room temperature led to the synthesis of the racemic lactams **3d-h** (59-74%), used as references for the enzymatic transformations.

 γ -Keto ester **1d** was initially studied and the best results are summarised in Table 2. In most of the cases, a lower reactivity was detected in comparison with alkylated esters 1a-c (see Tables S3-S5 in the Supporting Information for additional data). In fact, conversions lower than 60% were attained when working at 25 mM substrate concentration (entries 1 and 4), even with the best candidates. In order to reach higher conversions maintaining the excellent selectivity towards the synthesis of 3d, two reaction parameters were analysed: the substrate concentration (15 mM, entries 2 and 5) and the temperature (45 °C, entries 3 and 6). Remarkably, in the case of ATA-303 it was possible to obtain enantiopure (R)-3d in 97% conversion at 15 mM concentration and 45 °C.

Under these conditions, keto esters **1e-h** (entries 7-19) were transformed, yielding in all cases the lactams in enantiomerically pure form and in 81-99% conversion, depending on the enzyme (see Tables S6-S9 in the Supporting Information for extensive screenings). Interestingly, ATA-025 and ATA-033 were found as very robust enzymes for the formation of the (*R*)-enantiomers, while ATA-234 allowed the formation of their antipodes in a complete selective

manner, and much higher conversions in comparison with the one achieved with the non-substituted keto ester **1d** (56% conversion and 97% *ee*, see Table S5 in the Supporting Information).

As a continuation, 5-aryl-5-oxopentanoates 1i and 1j were selected in order to obtain the corresponding 6-arylpiperidin-2-ones **3i** and **3j**. Firstly, the chemical syntheses of both keto esters and the corresponding lactams were carried out, following identical procedures than the ones described for 1e-h and 3d-h. In general, biotransamination experiments led to lower conversion values in comparison with the results obtained with the homologue butanoates 1d and 1g (Table 3 and see additional data in Tables S10 and S11 in the Supporting Information). An increase in the temperature led to a conversion improvement for the (R)-selective ATA-033 and ATA-025 (entries 1-2 and 7-8), although this failed for ATA-415, which led to the best conversion value towards enantiopure (R)-3: at 30 °C (entries 3 and 4). (S)-Selective ATA-237 displayed the highest activities (94-96% conversion) acting with a high stereocontrol (93-95% ee, entries 6 and 10) at 45 °C.

Driving the equilibrium towards the formation of amine products is a key issue in transaminasecatalysed reactions, as the reductive amination process is thermodynamically hampered; therefore, a large molar excess of the amine donor is required. In the last years, the search for novel smart co-substrates or the *in situ* removal of the formed co-products have been applied.^[15] The advantage of the present strategy is the formation of an amine intermediate that cyclises in the same reaction medium forming the final lactam, so the reversibility of the equilibrium should be dramatically diminished. For that reason, we envisaged the use of lower contents of isopropylamine and the results havbeen summarised in Figure 2 (for numerical data see Table S12).

Table 3. Stereoselective transformation of δ -keto esters **1i**, **j** into 6-arylpiperidin-2-ones **3i**, **j**.

	R	0 1i,j (15 mM)	O (S)- Iso OMe KPi	or (<i>R</i>)-Transaminase opropylamine (1 M) PLP (1 mM) buffer 100 mM pH 7.5 DMSO (2.5% v/v) 24 h, 250 rpm	O N H 3i,j	
Entry	Keto ester	R	T [°C]	Transaminase	<i>c</i> [%] ^[a]	<i>ee</i> [%] ^[b]
1	1i	Н	30	ATA-033	20	>99 (<i>R</i>)
2	1i	Н	45	ATA-033	28	>99 (<i>R</i>)
3	1i	Н	30	ATA-415	71	>99 (<i>R</i>)
4	1i	Н	45	ATA-415	47	>99 (<i>R</i>)
5	1i	Н	30	ATA-237	94	94 (<i>S</i>)
6	1i	Н	45	ATA-237	94	95 (<i>S</i>)
7	1j	F	30	ATA-025	43	>99 (<i>R</i>)
8	1j	F	45	ATA-025	58	>99 (<i>R</i>)
9	1j	F	30	ATA-237	90	92 (<i>S</i>)
10	1i	F	45	ATA-237	96	93 (S)

^[a] Conversion values determined by GC (see Supporting Information). ^[b] Determined by HPLC using chiral columns (see Supporting Information). Absolute configurations of the corresponding lactams **3i** and **3j** appear in brackets.



Figure 2. Influence of the isopropylamine concentration in the TA-catalysed transformation of keto esters **1a-j** into lactams **3a,c-j**. Blue colour denotes the production of (*S*)-amines (89->99% *ee*), while red bars are used to indicate (*R*)-amines (>99% *ee*). **Figure 2A (left):** Biotransamination of keto ester **1b** into lactam **3a** using TA-P1-G06 under the following conditions: 25 mM **1b**, 25-1000 mM ¹PrNH₂, 30 °C, 24 h and 250 rpm. **Figure 2B (right):** Biotransamination of keto esters **1a-j** into lactams **3a,c-j** under the following conditions: for **1a-c**, 25 mM keto ester, 0.1 or 1 M ¹PrNH₂, 30 °C, 24 h and 250 rpm; for **1d-j**, 15 mM keto ester, 0.1 or 1 M ¹PrNH₂, 45 °C, 24 h and 250 rpm.

Methyl levulinate (**1b**, 25 mM) was selected as model substrate, and the biotransformations with TA-P1-G06 at 30 °C were carried out using from an equimolar amount to a 40-molar excess of IPA (Figure 2A). After 24 h, similar conversions (89-91%) were found between 0.1-1 M IPA concentrations, observing a significant decrease at 50 mM IPA concentration (80%). Even at equimolar amount, a high conversion was still observed (71%). Then, the transaminasecatalysed reactions with all the previously tested substrates (Figure 2B) were carried out at 0.1 and 1 M IPA concentrations, finding in general slightly lower conversions at lower IPA contents, although a clear negative effect was only remarkable for the case of **1h**.

Finally, preparative biotransformations (up to 100mg scale) were carried out under the optimised reaction conditions using 1 M of IPA in order to assure very high conversions (Table 4), yielding in all cases the desired lactams with good to high isolated yields (66-89%) and in general excellent selectivities (94->99% *ee*). Cosolvents with a lower boiling point than DMSO were used, such as MeCN for commercially available TAs, and MTBE for ArS-TA, searching for high-yielding processes towards the final products with easier work-up protocols. Optical rotation values were measured for all the isolated lactam products, and their values compared with reported data in order to assign their absolute configurations.^[16-20]

Keto esters **1a-c** bearing an alkyl substitution attached to the ketone functionality were used at 25 mM substrate concentration and 30 °C, while for aromatic compounds **1d-j** the substrate concentration was reduced to 15 mM, and a higher temperature (4. °C) was required to reach conversions over 70%. As previously observed in Figure 2, the use of a lowe amount of IPA (100 mM) could also lead to practical processes (entries 6 and 12). Thus, (*S*)- and (*R*) selective TAs were chosen for preparative biotransformations and the amount of IPA was 10-fold reduced, obtaining high conversions (89-93%) and good to excellent selectivities (91->99% *ee*) for substrates **1e** and **1j**.

Conclusions

A simple and straightforward one-pot two-step process has been described for the transformation of different γ - and δ -keto esters into the corresponding optically active lactams. The strategy is based on a selective biotransamination reaction that allows the formation of the amino ester intermediates, which undergo spontaneous intramolecular cyclisation in the own aqueous medium without the addition of external reagents. Overall, this is a general process as it was extended to 10 substrates with different substitution patterns. Amino ester intermediates were not detected in any case, facilitating the isolation of the final optically active lactam products. Table 4. Stereoselective transformations of keto esters 1b-j into lactams 3a,c-j under preparative scale.

		0		(S)- or (R)-Transa Isopropylamine (0	minase .1-1 M)	(
				PLP (1 mM	<u>)</u> ∩≊			
		R' (~)n	Ĭ	KPi buffer 100 mM	1 pH 7.5	H N		
		1 a_i (15_2	0 25 mM)	Cosolvent (2.5%	% v/v)	3a c₋i		
		Id-j (10-2	.5 (1111)	250 rpm		Ja,c-j		
Entry	[Keto ester]	[^{<i>i</i>} PrNH ₂] [M]	T [°C]	Transaminase	Cosolvent	t [h]	<i>c</i> [%] ^[a]	<i>ee</i> [%] ^[b]
1	25 (1a)	1	30	ATA-237	MeCN	24	97 (75)	>99 (S)
2	25 (1b)	1	30	TA-P1-G06	MeCN	48	88 (80)	>99 (S)
3	25 (1c)	1	30	ArS-TA	MTBE	24	98 (89)	>99 (S)
4	15 (1d)	1	45	ATA-303	MeCN	24	97 (88)	>99 (<i>R</i>)
5	15 (1e)	1	45	ATA-025	MeCN	48	82 (74)	>99 (<i>R</i>)
6	15 (1e)	0.1	45	ATA-025	MeCN	48	89 (80)	>99 (<i>R</i>)
7	15 (1f)	1	45	ATA-025	MeCN	48	91 (86)	>99 (<i>R</i>)
8	15 (1g)	1	45	ATA-025	MeCN	48	80 (75)	>99 (<i>R</i>)
9	15 (1h)	1	45	ATA-025	MeCN	48	89 (81)	>99 (<i>R</i>)
10	15 (1i)	1	45	ATA-237	MeCN	24	77 (70)	94 (S)
11	15 (1j)	1	45	ATA-237	MeCN	24	70 (66)	96 (<i>S</i>)
12	15 (1j)	0.1	45	ATA-237	MeCN	24	93 (90)	91 (S)

^[a] Conversion values determined by GC (see Supporting Information). Isolated yields appear in brackets. ^[b] Determined by HPLC using chiral columns (see Supporting Information). Absolute configurations of the corresponding lactams **3a,c-j** appear in brackets.

Chiral 6-substituted piperidin-2-ones and 5substituted pyrrolidin-2-ones were obtained with high optical purity (94->99% *ee*) and generally in good to very high yields (66-90%) for selected preparative biotransformations, the choice of the transaminase allowing the preparation of the desired lactam enantiomer under mild reaction conditions. Substrate concentration and temperature resulted to be key issues for driving the reactions to high conversions, being possible the reduction of the amine donor equivalents in selected cases without a significant drop of the conversion values due to the thermodynamically favoured intramolecular cyclisation process.

Experimental Section

General methods

Codex Transaminase ATA Screening Kit (ATASK-000250) and pyridoxal 5'-phosphate (PLP) were purchased from Codexis. Transaminases from *Chromobacterium violaceum* (Cv-TA, internal plasmid number pET20) and *Arthrobacter* sp. [ArR-TA (pEG23), ArS-TA (pEG29) and ArRmut11-TA (pEG90)] overexpressed on *E. coli cells* were provided by Prof. Wolfgang Kroutil (University of Graz).^[13c] All other reagents were obtained from commercial sources (Sigma–Aldrich, Acros, and Fluka) and used as received except dry methanol that was previously distilled under nitrogen using calcium hydride as desiccant.

NMR spectra were recorded on a Bruker AV300 MHz spectrometer. All chemical shifts (δ) are given in parts per million (ppm) and referenced to the residual solvent signal as internal standard. IR spectra were recorded on a Bruker ALPHA spectrophotometer on NaCl pellets or KBr pellets. v_{max} values are given for the main absorption bands. High resolution mass spectra (HRMS) were obtained in a Micro Tof Q spectrometer using ESI⁺ or ESI⁻. Measurement of the optical rotation values was carried out at 590 nm on a PerkinElmer 241 polarimeter.

Gas chromatography (GC) and high performance liquid chromatography (HPLC) analyses were performed for conversion and enantiomeric excess measurements (see the Supporting Information). GC analyses were performed on an Agilent HP7820 GC chromatograph equipped with a FID detector. HPLC analyses were carried out in a Hewlett Packard 1100 chromatograph UV detector at 210 nm. Thin layer chromatography (TLC) was conducted with Merck Silica Gel 60 F254 precoated plates and visualised with UV and potassium permanganate stain. Column chromatography was performed using Merck Silica Gel 60 (230-400 mesh).

General procedure for the chemical synthesis of γ -keto esters 1e-h and δ -keto esters 1i and 1j. To a solution of the corresponding keto acid 4e-j (1.6 mmol) in MeOH (8 mL, 0.2 M), a few drops of concentrated sulfuric acid were added at room temperature. The mixture was stirred and heated at 68 °C overnight. After this time, H₂O (10 mL) was added. The mixture was neutralised with an aqueous NaOH 2 M solution and the product was extracted with CH₂Cl₂ (3 x 15 mL). The organic layers were combined, dried over Na₂SO₄ and concentrated under reduced pressure, isolating the corresponding keto esters 1e-h,i,j in excellent purity, which were later used without further purification (91-99%).

Methyl 4-(4-methoxyphenyl)-4-oxobutanoate (1e). Yellowish solid (323 mg, 99% yield). $R_{\rm f}$ (50% EtOAc/Hexane): 0.4. Mp: 46-47 °C. IR (KBr): v 3058, 2954, 2918, 2844, 1737, 1602, 1439, 1318, 1030, 737 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 2.73 (t, ³*J*_{HH} = 6.7 Hz, 2H) 3.25 (t, ³*J*_{HH} = 6.7 Hz, 2H), 3.68 (s, 3H), 3.84 (s, 3H) 6.92 (d, ³*J*_{HH} = 8.9 Hz, 2H), 7.94 (d, ³*J*_{HH} = 8.9 Hz, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 28.0 (CH₂), 32.9 (CH₂), 51.7 (CH₃), 55.4 (CH₃), 113.7 (2 CH), 129.5 (C), 130.2 (2 CH), 163.5 (C), 173.4 (C), 196.5 (C). HRMS (ESI⁺, *m*/z): calcd for (C₁₂H₁₄NaO₄)⁺ (M+Na)⁺ 245.0784; found 245.0780. **Methyl 4-(4-methylphenyl)-4-oxobutanoate (1f)**

calcd for $(C_{12}H_{14}NaO_3)^+$ (M+Na)⁺ 229.0835; found 229.0831.

Methyl 4-(4-fluorophenyl)-4-oxobutanoate (1g). **Methyl** 4-(4-fluorophenyl)-4-oxobutanoate (1g). White solid (210 mg, 98% yield). R_f (50% EtOAc/Hexane): 0.37. Mp: 51-52 °C. IR (KBr): v 3064, 2954, 2918, 2850, 1738, 1599, 1439, 1412, 1268, 1157, 737 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 2.76 (t, ³J_{HH} = 6.6 Hz, 2H), 3.29 (t, ³J_{HH} = 6.6 Hz, 2H), 3.70 (s, 3H), 7.13 (t, ³J_{HH} = 8.6 Hz, 2H), 8.01 (dd, ³J_{HH} = 8.9 Hz, ³J_{HF} = 5.4 Hz, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 27.9 (CH₂), 33.2 (CH₂), 51.8 (CH₃), 115.6 (d, ²J_{CF} = 21.9 Hz, 2 CH), 130.6 (d, ³J_{CF} = 9.3 Hz, 2 CH), 132.9 (d, ⁴J_{CF} = 2.8 Hz, C), 165 (d, ¹J_{CF} = 254.8 Hz, C), 173.2 (C), 196.4 (C). HRMS (ESI⁺, m/z): calcd for (C₁₁H₁₁FNaO₃)⁺ (M+Na)⁺ 233.0584; found 233.0583. Methyl 4-(4-ethylphenyl)-4-oxobutanoate (1h).

4-(4-ethylphenyl)-4-oxobutanoate **Methyl** 4-(4-ethylphenyl)-4-oxobutanoate (1h). Yellowish oil (528 mg, 98% yield). $R_{\rm f}$ (50% EtOAc/Hexane): 0.5. IR (NaCl): v 3057, 2969, 2920, 2877, 2850, 1737, 1607, 1439, 1415, 1266, 1169, 738 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 1.26 (t, ³J_{HH} = 7.6 Hz, 3H), 2.67-2.78 (m, 4H), 3.30 (t, ³J_{HH} = 6.7 Hz, 2H), 3.70 (s, 3H), 7.28 (d, ³J_{HH} = 8.3 Hz, 2H), 7.91 (d, ³J_{HH} = 8.3 Hz, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 15.1 (CH₃), 28.0 (CH₂), 28.9 (CH₂), 33.2 (CH₂), 51.7 (CH₃), 128.0 (2 CH), 128.2 (2 CH), 134.2 (C), 150.1 (C), 173.4 (C), 197.6 (C). HRMS (ESI⁺, *m*/z): calcd for (C₁₃H₁₆NaO₃)⁺ (M+Na)⁺ 243.0992; found 243.0990. Methyl (**1h**). 243.0990

Methyl 5-oxo-phenylpentanoate (1i). Yellowish oil **Methyl 5-oxo-phenylpentanoate (1i).** Yellowish oil (489 mg, 92% yield). $R_{\rm f}$ (50% EtOAc/Hexane): 0.57. IR (KBr): v 3057, 2986, 2953, 2848, 1735, 1449, 1419, 1372, 1266, 1179, 704, 692 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 2.08 (quin, ³*J*_{HH} = 7.1 Hz, 2H), 2.44 (t, ³*J*_{HH} = 7.2 Hz, 2H), 3.04 (t, ³*J*_{HH} = 7.1 Hz, 2H), 3.66 (s, 3H), 7.44 (m, 2H), 7.54 (m, 1H), 7.95 (dd, ³*J*_{HH} = 8.4 Hz, ⁴*J*_{HH} = 1.4 Hz, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 19.3 (CH₂), 33.0 (CH₂), 37.4 (CH₂), 51.5 (CH₃), 127.9 (2 CH), 128.5 (2 CH), 133.0 (CH), 136.7 (C), 173.6 (C), 199.3 (C). HRMS (ESI⁺, *m/z*): calcd for (C₁₂H₁₄NaO₃)⁺ (M+Na)⁺ 229.0835; found 229.0835. **Methyl 5-(4-fluorophenyl)-5-oxopentanoate (1i).**

5-(4-fluorophenyl)-5-oxopentanoate **Methyl 5-(4-fluorophenyl)-5-oxopentanoate (1j).** White solid (504 mg, 91% yield). R_f (50% EtOAc/Hexane): 0.33. Mp: 46-47 °C. IR (KBr): v 3059, 2953, 2849, 1735, 1598, 1410, 1371, 1267, 1157, 738 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 2.06 (quin, ³J_{HH} = 7.1 Hz, 2H), 2.46 (t, ³J_{HH} = 7.1 Hz, 2H), 3.04 (t, ³J_{HH} = 7.1 Hz, 2H), 3.60 (s, 3H), 7.14 (t, ³J_{HH} = 8.6 Hz, 2H), 8.00 (dd, ³J_{HH} = 8.9 Hz, ³J_{HF} = 5.4 Hz, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 19.3 (CH₂), 33.0 (CH₂), 37.3 (CH₂), 51.6 (CH₃), 115.6 (d, ²J_{CF} = 21.8 Hz, 2 CH), 130.6 (d, ³J_{CF} = 9.4 Hz, 2 CH), 133.2 (d, ⁴J_{CF} = 3.3 Hz, C), 165.0 (d, ¹J_{CF} = 254.9 Hz, C), 173.6 (C), 197.7 (C). HRMS (ESI⁺, *m*/z): calcd for (C₁₂H₁₃FNaO₃)⁺ (M+Na)⁺ 247.0741; found 247.0742. Methyl (1i).

General procedure for the chemical synthesis of racemic γ -lactams 3d-h and δ -lactams 3i and 3j. In order to develop robust analytical methods for monitoring the biotransaminations, the synthesis of racemic lactams was previously chemically performed in the following manner: Ammonium acetate (200 mg, 2.6 mmol) and sodium cyanoborohydride (33 mg, 0.52 mmol) were successively added to a solution of the corresponding keto ester **1d-j** (0.26 mmol) in dry MeOH (1.0 mL) under inert atmosphere. The mixture was stirred at room temperature during 16 h and, after this time, H_2O (5 mL) was added to quench the reaction. The solution was basified until pH around 11 by adding a saturated aqueous solution of Na₂CO₃. Then the mixture was extracted with Et_2O (3 x 10 mL) and the organic layers were combined, dried over Na_2SO_4 and concentrated under reduced pressure. The product was purified by chromatography column on silica gel (2% MeOH/CH₂Cl₂), affording the racemic lactams **3d-j** (48-74%

5-Phenylpyrrolidin-2-one (3d). White solid (29 mg, **5-Phenylpyrrolidin-2-one (3d).** White solid (29 mg, 74% yield). $R_{\rm f}$ (5% MeOH/CH₂Cl₂): 0.38. Mp: 107-108 °C. IR (KBr): v 3055, 2959, 2926, 2872, 2858, 1687, 1266, 1157, 740, 705 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 1.94-2.06 (m, 1H), 2.38-2.52 (m, 2H), 2.54-2.66 (m, 1H), 4.78 (t, ³J_{HH} = 7.1 Hz, 1H), 6.01 (br s, 1H), 7.28-7.42 (m, 5H). ¹³C NMR (300.13 MHz, CDCl₃): δ 30.2 (CH₂), 31.4 (CH₂), 58.0 (CH), 125.6 (2 CH), 127.9 (CH), 128.9 (2 CH), 142.3 (C), 178.4 (C). HRMS (ESI⁺, m/z): calcd for (C₁₀H₁₁NNaO)⁺ (M+Na)⁺ 184 0733: found 184 0735 184.0733; found 184.0735.

5-(4-Methoxyphenyl)pyrrolidin-2-one (3e). Yellowish solid (26 mg, 60% yield). $R_{\rm f}$ (5% MeOH/CH₂Cl₂): 0.43. Mp: 115-116 °C. IR (KBr): v 3426, 3055, 2986, 1696, 1265, 1035, 739 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 1.96 (m, 1H), 2.39-2.56 (m, 3H), 3.80 (s, 3H), 4.71 (t, ³J_{HH} = 7.0 Hz, 1H), 6.44 (br s, 1H), 6.89 (d, ³J_{HH} = 8.7 Hz, 2H), 7.21 (d, ³J_{HH} = 8.7 Hz, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 30.4 $C(H_2)$, 31.5 (CH₂), 55.3 (CH₃), 57.6 (CH), 114.2 (2 CH), 126.8 (2 CH), 134.4 (C), 159.2 (C), 178.4 (C). HRMS (ESI⁺, m/z): calcd for (C₁₁H₁₃NNaO₂)⁺ (M+Na)⁺ 214.0838; found 214.0836

5-(4-Methylphenyl)pyrrolidin-2-one (3f). White solid **5-(4-Methylphenyl)pyrrolidin-2-one (3f).** White solid (31 mg, 73% yield). $R_{\rm f}$ (5% MeOH/CH₂Cl₂): 0.5. Mp: 89-90 °C. IR (KBr): v 3425, 3054, 2985, 2922, 2852, 1695, 1266, 1156, 738, 705 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 1.93-2.06 (m, 1H), 2.38 (s, 3H), 2.45 (m, 2H), 2.51-2.62 (m, 1H), 4.73 (t, ³*J*_{HH} = 7.1 Hz, 1H), 6.23 (br s, 1H), 7.14-7.28 (m, 4H). ¹³C NMR (300.13 MHz, CDCl₃): δ 21.0 (CH₃), 30.3 (CH₂), 31.4 (CH₂), 57.8 (CH), 126.2 (2 CH), 129.2 (2 CH), 137.7 (C), 139.4 (C), 178.4 (C). HRMS (ESI⁺, *m*/*z*): calcd for (C₁₁H₁₃NNaO)⁺ (M+Na)⁺ 198.0889; found 198.0888 198.088

calcd for $(C_{11}H_{13}NNaO)^{+}$ $(M+Na)^{+}$ 198.0889; found 198.0888. **5-(4-Fluorophenyl)pyrrolidin-2-one (3g).** White solid (25 mg, 59% yield). R_f (5% MeOH/CH₂Cl₂): 0.5. Mp: 138-139 °C. IR (KBr): v 3425, 3055, 2986, 2924, 1698, 1606, 1512, 1422, 1265, 738 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 1.88-2.00 (m, 1H), 2.41-2.52 (m, 2H), 2.58 (m, 1H), 4.78 (t, ³J_{HH} = 7.1 Hz, 1H), 7.05 (t, ³J_{HH} = 8.6 Hz, 2H), 7.08 (s, 1H), 7.29 (dd, ³J_{HH} = 8.7 Hz, ⁴J_{HF} = 5.2 Hz, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 30.3 (CH₂), 31.4 (CH₂), 57.4 (CH), 115.7 (d, ²J_{CF} = 21.6 Hz, 2 CH), 127.3 (d, ³J_{CF} = 8.1 Hz, 2 CH), 130.7 (d, ⁴J_{CF} = 3.1 Hz, C), 162.0 (d, ¹J_{CF} = 246.1 Hz, C), 178.3 (C). HRMS (ESI⁺, m/z): calcd for (C₁₀H₁₀FNNaO)⁺ (M+Na)⁺ 202.0639; found 202.0635. **5-(4-Ethylphenyl)pyrrolidin-2-one** (**3h**). Yellowish solid (30 mg, 69% yield). R_f (5% MeOH/CH₂Cl₂): 0.55. Mp: 82-83 °C. IR (KBr): v 3425, 3054, 2984, 2970, 2931, 1696, 1.24 (t, ³J_{HH} = 7.6 Hz, 3H), 1.99 (m, 1H), 2.37-2.47 (m, 2H), 2.48-2.60 (m, 1H), 2.66 (q, ³J_{HH} = 7.6 Hz, 2H), 4.74 (t, ³J_{HH} = 7.1 Hz, 1H), 6.05 (br s, 1H), 7.24 (m, 4H). ¹³C NMK (300.13 MHz, CDCl₃): δ 15.5 (CH₃), 28.4 (CH₂), 30.3 (CH₂), 31.4 (CH₂), 57.9 (CH), 125.6 (2 CH), 128.3 (2 CH), 139.7 (C), 144.1 (C), 178.3 (C). HRMS (ESI⁺, m/z): calcd for (C₁₂H₁₅NNaO)⁺ (M+Na)⁺ 212.1046; found 212.1044. **6-Phenylpiperidin-2-one** (**3i**). White solid (12 mg, 48%, vield). R_t (5% MeOH/CH₂Cl₂): 0.46, Mp; 112-113 °C. IR

(C₁₂H₁₅NNaO)⁺ (M+Na)⁺ 212.1046; found 212.1044. **6-Phenylpiperidin-2-one (3i).** White solid (12 mg, 48%, yield). $R_{\rm f}$ (5% MeOH/CH₂Cl₂): 0.46. Mp: 112-113 °C. IR (KBr): v 3389, 3054, 2985, 2957, 293, 2853, 1657, 1265, 738, 704 cm^{-1.} ¹H NMR (300.13 MHz, CDCl₃): δ 1.63-1.74 (m, 1H), 1.76-1.86 (m, 1H), 1.87-1.99 (m, 1H), 2.03-2.17 (m, 1H), 2.37-2.52 (m, 2H), 4.57 (dd, ³J_{HH} = 8.8, 4.6 Hz, 1H), 6.02 (s, 1H), 7.28-7.43 (m, 5H). ¹³C NMR (300.13 MHz, CDCl₃): δ 19.7 (CH₂), 31.3 (CH₂), 32.2 (CH₂), 57.8 (CH), 126.0 (2 CH), 127.9 (CH), 128.8 (2 CH), 142.4 (C), 172.3 (C). HRMS (ESI⁺, m/z): calcd for (C₁₁H₁₃NNaO)⁺ (172.3 (C). HRMS (ESI⁺, m/z): calcd for (C₁₁H₁₃NNaO)⁺ (M+Na)⁺ 198.0889; found 198.0887.

(M+Na)⁺ 198.0889; found 198.0887. **6-(4-Fluorophenyl)piperidin-2-one (3j).** White solid (11 mg, 50% yield). $R_{\rm f}$ (5% MeOH/CH₂Cl₂): 0.54. Mp: 99-100 °C. IR (KBr): v 3389, 3055, 2986, 2930, 1659, 1265, 739 cm^{-1.} ¹H NMR (300.13 MHz, CDCl₃): δ 1.59-1.70 (m, 1H), 1.71-1.86 (m, 1H), 1.87-1.98 (m, 1H), 2.11 (m, 1H), 2.37-2.56 (m, 2H), 4.56 (dd, ³J_{HH} = 8.9, 4.5 Hz, 1H), 6.02 (br s, 1H), 7.08 (t, ³J_{HH} = 8.6 Hz, 2H), 7.29 (m, 2H). ¹³C NMR (300.13 MHz, CDCl₃): δ 19.8 (CH₂), 31.2 (CH₂), 32.3 (CH₂), 57.2 (CH), 115.7 (d, ²J_{CF} = 21.6 Hz, 2 CH), 127.7 (d, ³J_{CF} = 8.2 Hz, 2 CH), 138.2 (d, ⁴J_{CF} = 2.9 Hz, C), 164.0 (d, ¹J_{CF} = 246.8 Hz, C), 172.2 (C). HRMS (ESI⁺, *m/z*): calcd for (C₁₁H₁₂FNNaO)⁺ (M+Na)⁺ 216.0795; found 216.0793.

General protocol for the transformation of γ - and δ -keto esters 1a-j into optically active lactams using commercially available transaminases. In a 1.5 mL Eppendorf tube, transaminase (2 mg) and the corresponding keto ester (**1a-j**, 15 or 25 mM) were added in a phosphate buffer 100 mM pH 7.5 (500 μ L, 1 mM PLP, 1 M isopropylamine), using DMSO (12.5 μ L) as cosolvent. The reaction was shaken at 30 or 45 °C and 250 rpm for 24 h and stopped by the addition of an aqueous Na₂CO₂ saturated stopped by the addition of an aqueous Na₂CO₃ saturated solution (200 μ L). Then the mixture was extracted with EtOAc (2 x 500 μ L), the organic layers separated by

General method for the transformation of γ - and δ -keto esters 1a-j into optically active lactams using transaminases overexpressed in *E. coli*. In a 1.5 mL Eppendorf tube, dry cells of *E. coli* overexpressing the transaminase (10 mg) and γ -keto ester (**1a-j**, 25 mM) were added in a phosphate buffer 100 mM pH 7.5 (500 μ L, 1 mM PLP, 1 M isopropylamine) using DMSO (12.5 μ L) as cosolvent. The reaction was shaken at 30 °C and 250 rpm for 24 h and stopped by the addition of an aqueous Na_2CO_3 saturated solution (200 μ L). Then the mixture was extracted with EtOAc (2 x 500 μ L), the organic layers separated by centrifugation (2 min, 13000 rpm), combined and finally dried over Na₂SO₄. Conversions into lactams **3a-j** were determined by GC and *ee* values measured by HPLC.

Preparative biotransformation of γ -keto esters 1a-c into optically active lactams. In an Erlenmeyer flask, the transaminase (30 mg, ATA-237 for 1a, TA-P1-G06 for 1b or ArS-TA for 1c) and keto ester (1a-c, 30 mg, 25 mM) were added in a phosphate buffer 100 mM pH 7.5 (1 mM PLP, 1 M isopropylamine) and cosolvent (2.5% v/v MeCN for 1a-b and 2.5% v/v MTPE for 1a). The reaction was abelien at **b** and 2.5% v/v MTBE for **1c**). The reaction was shaken at 30 °C and 250 rpm for 24 h (**1a** and **1c**) or 48 h (**1b**) and then stopped by the addition of an aqueous Na₂CO₃ saturated solution until pH 10-11. Then, the mixture was extracted with EtOAc (5 x 15 mL), the organic layer separated by centrifugation (5 min, 4900 rpm), combined and finally dried over Na₂SO₄. The reaction crude was (MeOH/CH₂Cl₂ purified by column chromatography (MeOH/CH₂Cl₂ mixtures), isolating the enantiopure lactams (*S*)-**3a**,**c** in good yields (75-89%, respectively).

mixtures), isolating the enalthopine factants (3)-**5a**, **c** in good yields (75-89%, respectively). (*S*)-**5-Methylpyrrolidin-2-one** [(*S*)-**3a**]: Yellowish solid (19 mg, 80% yield). R_f (5% MeOH/CH₂Cl₂): 0.57. Mp: 43-44 °C. IR (NaCl): v 3227, 3054, 2971, 2930, 1694, 1462, 1265 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 1.17 (d, ³*J*_{HH} = 6.3 Hz, 3H), 1.46-1.74 (m, 1H), 2.14-2.25 (m, 1H), 2.25-2.34 (m, 2H), 3.56-3.90 (m, 1H), 7.37 (s, 1H). ¹³C NMR (300.13 MHz, CDCl₃): δ 19.8 (CH₂), 22.7 (CH₃), 30.4 (CH₂), 30.9 (CH₂), 48.7 (CH), 172.5 (C). HRMS (ESI⁺, m/z): calcd for for (C₅H₉NNaO)⁺ (M+Na)⁺ 122.0576; found 122.0574. (*S*)-**6-Methylpiperidin-2-one** [(*S*)-**3c**]: White solid (19 mg, 89% yield). R_f (5% MeOH/CH₂Cl₂): 0.51. Mp: 87-88 °C. IR (NaCl): v 3392, 3054, 2972, 2935, 1659, 1468, 1265 cm⁻¹. ¹H NMR (300.13 MHz, CDCl₃): δ 1.18 (d, ³*J*_{HH} = 6.4 Hz, 3H), 1.26-1.36 (m, 1H), 1.52-1.79 (m, 1H), 1.75-1.98 (m, 2H), 2.09-2.45 (m, 2H), 3.44-3.55 (m, 1H), 6.69 (s, 1H). ¹³C NMR (300.13 MHz, CDCl₃): δ 22.1 (CH₃), 29.0 (CH₂), 30.7 (CH₂), 50.2 (CH), 178.7 (C). HRMS (ESI⁺, m/z): calcd for (C₆H₁₁NNaO)⁺ (M+Na)⁺ 136.0732; found 136.0730.

Preparative biotransformation of y-keto esters 1d and 1e and δ -keto esters 1i and 1j into optically active lactams. In an Erlenmeyer flask, the TA (30 mg, ATA-303 for keto ester 1d, ATA-025 for keto ester 1e and ATA-237 for keto ester **1u**, A1A-2257 for keto ester **1e** and A1A-257 for keto esters **1i** and **1j**) and keto ester (**1d**, **1e**, **1i** and **1j**, 30 mg, 15 mM) were added in phosphate buffer 100 mM pH 7.5 (1 mM PLP, 0.1 or 1 M isopropylamine) and MeCN (2.5% v/v). The reaction was shaken at 45 °C and 250 rpm for 24 h (**1d**, 1i and 1j) or 48 h (1e) and then stopped by adding an aqueous saturated solution of Na_2CO_3 until pH 10-11. Then the mixture was extracted with ethyl acetate (5 x 15 mL), the organic layer separated by centrifugation (5 min, 4900 rpm), combined and finally dried over Na_2SO_4 . The reaction (MeOH/CH₂Cl₂ mixtures), yielding the lactams (R)-3d and 3e and (S)-3i and 3j in moderate to good yields (66-90%) and good to excellent enantiomeric excess (91->99%).

Preparative biotransformation of y-keto esters 1f-h into optically active lactams. In an Erlenmeyer flask, ATA-025 (100 mg) and keto ester (1f-h, 100 mg, 15 mM) were added in phosphate buffer 100 mM pH 7.5 (1 mM PLP, 0.1 or 1 M isopropylamine) and MeCN (2.5% v/v). The reaction was shaken at 45 °C for 48 h and then stopped by adding a

saturated solution of Na₂CO₃ until pH 10-11. Then, the mixture was extracted with ethyl acetate (5 x 15 mL), the organic layer separated by centrifugation (5 min, 4900 rpm), combined and finally dried over Na₂SO₄. The reaction crude was purified by column chromatography (MeOH/CH₂Cl₂ was purfied by column chromatography (MeOH/CH₂Cl₂ mixtures), yielding lactams (*R*)-**3f**-**h** in good yields (75-86%) and excellent enantiomeric excess (>99%). Optical rotation values for (*R*)-lactams **3f**-**h**: $[\alpha]_D^{25} = +24.5$ (c = 1.0, EtOH) for (*R*)-**3f** in >99% *ee*; $[\alpha]_D^{25} = +21.1$ (c = 1.0, EtOH) for (*R*)-**3g** in >99% *ee*; $[\alpha]_D^{25} = +44.1$ (c = 1.0, EtOH) for (*R*)-**3h** in >99% *ee*.

Acknowledgements

Financial support from the Spanish Ministry of Economy and (MEC, Projects CTQ2013-44153-P Competitiveness and CTQ2016-75752-R) and the Government of the Principado de Asturias (Project FC-15-GRUPIN14-002) is Asturias (Project FC-15-GRUPIN14-002) is gratefully acknowledged. A.M.-I thanks MEC for a predoctoral fellowship inside the FPI program, while L.A.Z thanks the national Brazilian agency CAPES for a postdoctoral fellowship agency CAPES for a postdoctoral fellowship.

References

- [1] M. A. Ogliaruso, J. F. Wolfe, Synthesis of Lactones and Lactams, John Wiley & Sons, Inc., West Sussex, 1993.
- [2] a) C. R. Pitts, T. Lectka, Chem. Rev. 2014, 114, 7930-7953; b) G. S. Singh, S. Sudheesh, Arkivoc 2014, i, 227-385; c) Y. Kumar, P. Singh, G. Bhargava, RSC Adv. 2016, 6, 99220-99250.
- [3] a) L.-W. Ye, C. Shu, F. Gagosz, Org. Biomol. Chem. 2014, 12, 1833-1845; b) J. Caruano, G. G. Muccioli, R. Robiette, Org. Biomol. Chem. 2016, 14, 10134-10156.
- [4] J. Albarrán-Velo, D. González-Martínez, V. Gotor Fernández, Biocatal. Biotransf. 2018, 36, 112-140.
- [5] E. Busto, V. Gotor-Fernández, V. Gotor, Chem. Rev. 2011, 111, 3998-4035.
- [6] See, for instance: a) A. L. Gutman, E. Meyer, X. Yue, C. Abell, Tetrahedron Lett. 1992, 33, 3943-3946; b) E. Stavila, K. Loos, Tetrahedron Lett. 2013, 54, 370-372; c) C.-J. Aurell, S. Karlsson, F. Pontén, S. M. Andersen, Org. Process Res. Devel. 2014, 18, 1116-1119.
- [7] a) D. Koszelewski, D. Clay, K. Faber, W. Kroutil, J. Mol. Catal. B: Enzym. 2009, 60, 191-194; b) M. D. Truppo, J. D. Rozzell, N. J. Turner, Org. Process Res. Devel. 2010, 14, 234-237; c) C. K. Chung, P. G. Bulger, B. Kosjek, K. M. Belyk, N. Rivera, M. E. Scott, G. R. Humphrey, J. Limanto, D. C. Bachert, K. M. Emerson, Org. Process Res. Devel. 2014, 18, 215-227.
- [8] G. A. Aleku, S. P. France, H. Man, J. Mangas-Sanchez, S. L. Montgomery, M. Sharma, F. Leipold, S. Hussain, G. Grogan, N. J. Turner, Nat. Chem. 2017, 9, 961-969.
- [9] a) A. S. Touchy, S. M. A. Hakim Siddiki, K. Kon, K.-i. Shimizu, ACS Catal. 2014, 4, 3045-3050; b) J. D. Vidal, M. J. Climent, P. Concepcion, A. Corma, S. Iborra, M. J. Sabater, ACS Catal. 2015, 5, 5812-5821.
- [10] C. E. Paul, M. Rodríguez-Mata, E. Busto, I. Lavandera, V. Gotor-Fernández, V. Gotor, S. García-Cerrada, J. Mendiola, O. de Frutos, I. Collado, Org. Process Res. Devel. 2014, 18, 788-792.

- [11] U. Kaulman, K. Smithies, M. E. B. Smith, H. C. Hailes, J. M. Ward, *Enzyme Microb. Technol.* **2007**, *41*, 628-637.
- [12] S. Pannuri, S. V. Kamat, A. R. M. Garcia, WO 2006/063336 A2 20060615, Cambrex North Brunswick Inc., 2006.
- [13] a) Y. Yamada, A. Iwasaki, N. Kizaki, K. Matsumoto, Y. Ikenaka, M. Ogura, J. Hasegawa, (Kaneka Corporation), PCT Int. Appl. WO 9848030A1 19981029, **1998**; b) A. Iwasaki, Y. Yamada, N. Kizaki, Y. Ikenaka, J. Hasegawa, *Appl. Microbiol. Biotechnol.* **2006**, 69, 499-505; c) F. G. Mutti, C. S. Fuchs, D. Pressnitz, J. H. Sattler, W. Kroutil, *Adv. Synth. Catal.* **2011**, *353*, 3227-3233.
- [14] C. K. Savile, J. M. Janey, E. M. Mundorff, J. C. Moore, S. Tam, W. R. Jarvis, J. C. Colbeck, A. Krebber, F. J. Fleitz, J. Brands, P. N. Devine, G. W. Huisman, G. J. Georges, *Science* **2010**, *329*, 305-309.
- [15] F. Guo, P. Berglund, Green Chem. 2017, 19, 333-360.
- [16] (*S*)-5-Methylpyrrolidin-2-one (**3a**): $[\alpha]_D{}^{25} = -10.0$ (c = 1.0, EtOH). Described in the literature: $[\alpha]_D{}^{25} = -21.3$ (c = 0.9, water) for the (*S*)-enantiomer. J. M. García, M. A. Maestro, M. Oiarbide, J. M. Odriozola, J. Razkin, C. Palomo, *Org. Lett.* **2009**, *11*, 3826-3829.

- [17] (*S*)-6-Methylpiperidin-2-one (**3c**): $[\alpha]_D^{22} = +23.7$ (c = 1.0, EtOH). Described in the literature: $[\alpha]_D^{20} = -24.0$ (c = 0.4, CH₂Cl₂) for the (*R*)-enantiomer. M. Amat, N. Llor, J. Hidalgo, C. Escolano, J. Bosch, *J. Org. Chem.* **2003**, 68, 1919-1928.
- [18] (*R*)-5-Phenylpyrrolidin-2-one (**3d**): $[\alpha]_D{}^{20} = +42.2$ (c = 1.0, EtOH). Described in the literature: $[\alpha]_D{}^{20} = +41.0$ (c = 0.4, CH₂Cl₂) for the (*R*)-enantiomer. M. N. Cheemala, P. Knochel, *Org. Lett.* **2007**, *9*, 3089-3092.
- [19] (*R*)-5-(4-Methoxyphenyl)pyrrolidin-2-one (**3e**): $[\alpha]_D^{20}$ = +28.1 (c = 1.0, EtOH). Described in the literature: $[\alpha]_D^{20}$ = + 40.5 (c = 0.7, CH₂Cl₂) for the (*R*)-enantiomer. D. Guijarro, O. Pablo, M. Yus, *J. Org. Chem.* **2013**, 78, 3647-3654.
- [20] (*S*)-6-Phenylpiperidin-2-one (**3i**): $[\alpha]_D{}^{20} = -48.6$ (c = 1.0, EtOH). Described in the literature: $[\alpha]_D{}^{20} = -58.0$ (c = 0.54, CHCl₃) for the (*S*)-enantiomer. (*S*)-6-(4-Fluorophenyl)piperidin-2-one (**3j**): $[\alpha]_D{}^{20} = -35.4$ (c = 0.6, EtOH). Described in the literature: $[\alpha]_D{}^{20} = -47.2$ (c = 0.60, CHCl₃) for the (*S*)-enantiomer. R. Sallio, S. Lebrun, F. Agbossou-Niedercorn, C. Michon, E. Deniau, *Tetrahedron: Asymmetry* **2012**, *23*, 998-1004.

FULL PAPER

Conversion of γ - and δ -Keto Esters into Optically Active Lactams. Transaminases in Cascade Processes

Adv. Synth. Catal. Year, Volume, Page – Page

Ángela Mourelle-Insua, Luiz Arthur Zampieri, Iván Lavandera* and Vicente Gotor-Fernández*

