


This article was downloaded by: [North Dakota State University]

On: 07 September 2013, At: 12:45

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcar20>

Synthesis of the C-Glycoside of Methyl α -d-Altropyranosyl-(1 \rightarrow 4)- α -d-glucopyranoside

Richard W. Denton^a & David R. Mootoo^a

^a Department of Chemistry, Hunter College, 695 Park Avenue, New York, New York, 10021, USA

Published online: 23 Feb 2007.

To cite this article: Richard W. Denton & David R. Mootoo (2003) Synthesis of the C-Glycoside of Methyl α -d-Altropyranosyl-(1 \rightarrow 4)- α -d-glucopyranoside, Journal of Carbohydrate Chemistry, 22:7-8, 671-683, DOI: [10.1081/CAR-120026467](https://doi.org/10.1081/CAR-120026467)

To link to this article: <http://dx.doi.org/10.1081/CAR-120026467>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis of the C-Glycoside of Methyl α -D-Altropyranosyl-(1 \rightarrow 4)- α -D-glucopyranoside[†]

Richard W. Denton and David R. Mootoo*

Department of Chemistry, Hunter College, New York, New York, USA

ABSTRACT

The C-glycoside of methyl α -D-altropyranosyl-(1 \rightarrow 4)- α -D-glucopyranoside **2** was prepared in a convergent fashion, from readily available precursors, 4-*O*-*tert*-butyl-diphenylsilyl-1,2-*O*-isopropylidene-D-*erythro*-5-phenyl monothiohemiacetal **13** (five steps from D-ribose) and the known acid, methyl 2,3,6-tri-*O*-benzyl-4-*C*-(carboxymethyl)-4-deoxy- α -D-glucopyranoside **17** (seven steps from methyl α -D-glucopyranoside). The key reactions in the synthesis are the oxocarbenium ion cyclization of thioacetal-enol ether **19** to a C1 substituted glycal **20**, and the stereoselective hydroboration of **20** to the α -C-altroside **21**.

Key Words: C-glycoside; Disaccharide; Glycomimetic; Altrose; Allose; Conformation.

INTRODUCTION

Interest in disaccharide analogues as specific inhibitors of glycosidases and glycosyl transferases is based on the notion that they may act as bisubstrate-type transition state mimetics.^[1,2] This represents a somewhat oversimplified model because other aspects of the transition state structure such as the involvement of active site residues

[†]This paper is dedicated to Professor Gérard Decotes on the occasion of his 70th birthday.

*Correspondence: David R. Mootoo, Department of Chemistry, Hunter College, 695 Park Avenue, New York, NY 10021, USA; Fax: +1-212-772-5332; E-mail: dmootoo@hunter.cuny.edu.






Figure 1.

and sugar nucleotide components, are not incorporated.^{[3–8]ab} Notwithstanding this issue, simple disaccharide probes continue to be popular because of their potential to provide information on the minimal requirements for recognition. In this regard, disaccharide analogues^{[9–13]c} with “unnatural” glycone and aglycone substitutes are of interest. C-glycosides have additional appeal because of their hydrolytic stability, and their subtle conformational differences compared with O-glycosides.^[12–21] These principles are embodied in a recent investigation on the C-disaccharide of α -D-mannopyranosyl-(1 \rightarrow 3)-*N*-acetylgalactosamine **1**.^{[9]c} Compound **1** was found to be a good bisubstrate inhibitor of human α -(1,3)-fucosyltransferase VI (which catalyzes the transfer of GDP-fucose to *N*-acetylactosamine). Presumably, the α -L-fucopyranosyl and the D-GlcNAc binding sites of the enzyme are homogeneous with the respective binding regions for the C- α -D-mannopyranosyl and the D-GalNAc residues of **1** (Figure 1).


Against this backdrop, we have shown that 1-thio-1,2-*O*-isopropylidene acetals (TIAs) are precursors to β -C-galactodisaccharides.^[22,23] Herein, in the preparation of the C-glycoside of methyl α -D-altropyranosyl-(1 \rightarrow 4)- α -D-glucopyranoside **2**, we illustrate the suitability of this methodology to the synthesis of C-disaccharides with unusual glycone segments. This investigation is also pertinent to the altromycins, a naturally occurring class of aryl- α -C-altrosides.^[24]

Our synthesis of the β -C-galactoside **3** centered on the methyl triflate promoted cyclization of the thioacetal-enol ether **6** to the C1-substituted glycal **4**. This reaction presumably proceeds via a dioxolenium ion intermediate. Notably, the other double bond regioisomer or the acetal stereoisomer was not observed. Hydroboration of **4** led to a single C-glycoside of β -galacto stereochemistry. The cyclization precursor **6** was assembled from the TIA alcohol **7** and the requisite acid **8** over two steps, esterification of **7** and **8** followed by Tebbe olefination of ester **5**. This sequence is convergent from the glycone and aglycone segments and is general for β -C-galactosides. We anticipated that extension of these concepts to diastereomeric TIA alcohol **13**, should lead to the corresponding structures **10** and **9**, thereby constituting a general route to α -C-altrosides (Scheme 1).

^aFor transferase inhibitors containing glycosyl donor and sugar nucleotide residues, see: Refs. ^[3–5].

^bFor tricomponent, bisubstrate analogues, see: Refs. ^[6] and ^[7].

^cFor a more complex, two component bisubstrate inhibitor of an α -1,3-fucosyltransferase, see: Ref. ^[10].




Scheme 1.

RESULTS AND DISCUSSION


The TIA segment **13** for the C-altroside synthesis was obtained through the identical methodology that was used for the TIA in the C-galactoside series.^[23] Thus 2,3-*O*-isopropylidene-D-ribofuranose **14**^[25] was converted to the silyl ether **15**, and this material treated with diacetoxyiodosobenzene and iodine, according to the Suarez protocol.^[26] The resulting 1-*O*-acetyl-1,2-*O*-isopropylidene **16** was exposed to thiophenol and $\text{BF}_3 \cdot \text{OEt}_2$ at -78°C , and the crude reaction mixture subjected to basic hydrolysis of the formate ester. This sequence provided TIA alcohol **13** in 81% yield from **16**. The configuration of the acetal carbon in **13** was not determined (Scheme 2).

The aglycone segment **17** for the title C-altroside, was obtained by hydrolysis of the known ethyl ester derivative.^[27,28] DCC mediated esterification of alcohol **13** and acid **17** afforded ester **18**, which was transformed to enol ether **19** by treatment with the Tebbe reagent (Scheme 3). The key cyclization step on **19** was promoted by methyl triflate in the presence of 2,6-di-*tert*-butyl-4-methylpyridine, and led to C1 substituted glycal **20** in 70% yield. Hydroboration of **20** provided an approximately 5:1 ratio of two stereoisomers **21** and **22**, in a combined yield of 84%. Chromatography of this mixture




Scheme 2. (a) TBDPSCI, DMF, imidazole, 87%; (b) diacetoxyiodoso-benzene, I_2 , cyclohexane, 77%; (c) (i) PhSH , CH_2Cl_2 , $\text{BF}_3 \cdot \text{OEt}_2$, -78°C ; (ii) NaOMe , MeOH , 88%.





Scheme 3. (a) **13**, DCC, DMAP, PhH, 85%; (b) Tebbe, 60%; (c) MeOTf, DTBMP, CH₂Cl₂, 70%; (d) BH₃ · DMS then Na₂O₂, 63%; (e) HCl, MeOH; (f) Bu₄NF, THF; (g) Ac₂O, DMAP, EtOAc.

provided partial separation, giving a sample of pure **21** and an unseparated mixture of **21** and **22**. The stereochemistry of **21** was determined by ¹H NMR analysis of tetraacetate **26**. Compound **26** was obtained by sequential removal of isopropylidene and silyl ether protecting groups in **21**, followed by acetylation of the derived tetraol **25**. The *J* values (*J*_{2',3'} = *J*_{3',4'} = 8.5, *J*_{4',5'} = 2.5, *J*_{5',6'} < 2.0 Hz) for **26** were in agreement with an α-C-altroside in primarily the ¹C₄ conformation.^[29] The minor product **22**, was characterized as the triacetate **27**. Thus, acid hydrolysis of the mixture of hydroboration products **21** and **22**, provided an easily separable mixture of the corresponding triols **23** and **24**. Acetylation of **24** provided **27**. The data for **27** (*J*_{2',3'} = 9.5, *J*_{3',4'} = *J*_{4',5'} = 2.0, *J*_{5',6'} = 9.9 Hz) supported a β-alloside in the ⁴C₁ conformation,^[30] a result that is consistent with the stereochemistry of the hydroboration reaction.



Scheme 4. (a) H₂, Pd/C, MeOH, 80%.

The targeted *altro*- α -C-disaccharide **2** was obtained by the straightforward processing of the major component **21** from the mixture of hydroboration products (Scheme 4). Thus, hydrogenolysis of the aforementioned tetraol **25**, provided **2**. The structure of **2** was confirmed by ^1H and ^{13}C NMR and HRMS data. The stereochemical integrity of **2** resides in the assignment of the aforementioned tetraacetate derivative **26**. Unfortunately, poor signal resolution in the ^1H NMR of **2** did not allow for conformational analysis of the altropyranoside residue.

In summary, the synthesis of **2**, taken together with earlier studies, illustrates that the TIA C-glycoside methodology may be used for the stereoselective synthesis of β -C-*galacto*- and α -C-*altro*-disaccharides. An attractive feature is the convergent nature of this strategy, which allows for incorporation of different aglycone subunits. Elaboration of the intermediate disaccharide glycal intermediates provides additional possibilities for variation of the glycone segment. These attributes are especially relevant to the synthesis of libraries of C-disaccharides with unusual substitution patterns.

EXPERIMENTAL

General methods. Unless otherwise stated, all reactions were carried out under a nitrogen atmosphere in oven-dried glassware using standard syringe and septa technique. Diethyl ether and THF were distilled from sodium or potassium/benzophenone ketyl under N_2 immediately prior to use. Dry CH_2Cl_2 was obtained by distilling over P_2O_5 . Dry DMF was distilled from CaSO_4 under reduced pressure. Anhydrous benzene and toluene were obtained by azeotropic removal of water.

^1H and ^{13}C NMR spectra were obtained using a GE QE 300 (300 MHz) or Varian Unity Plus 500 (500 MHz) spectrometer. Chemical shifts are relative to the deuterated solvent peak or the tetramethylsilane (TMS) peak at (δ 0.00) and are in parts per million (ppm). The ^1H and ^{13}C NMR spectra were recorded at either 300 or 500 MHz and 75 MHz, respectively. Assignments for selected nuclei were determined from ^1H COSY experiments. The assignments for signals that are marked with an asterisk (*) are interchangeable.

High resolution mass spectroscopy was performed at the Mass Spectrometry Facility at University of Illinois, Urbana. Infrared (IR) spectra were recorded for neat films or CHCl_3 solutions on a Perkin-Elmer 710B spectrometer and are reported in cm^{-1} . Thin-layer chromatography (TLC) was done on 0.25 mm thick precoated silica gel HF₂₅₄ (Whatman) aluminum sheets. The chromatograms were observed under UV light and/or were visualized by heating plates that were dipped in ammonium molybdate solution. Flash column chromatography (FCC) was performed using silica gel 60 (230–400 mesh) and employed a stepwise solvent polarity gradient, correlated with TLC mobility.

2,3-O-Isopropylidene-5-O-*tert*-butyldiphenylsilyl- α/β -D-ribofuranose (15). A solution of 2,3-O-isopropylidene-D-ribofuranose **14**^[25] (7.86 g, 41.0 mmol), TBDPSCl (11.4 mL, 41.0 mmol), and imidazole (5.64 g, 85.0 mmol) in anhydrous DMF (100 mL) was stirred at 50°C for 1.5 h. The reaction mixture was then diluted with water and extracted with ether. The combined organic phase was washed with brine, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by FCC to give an inseparable mixture **15 α/β** (15.2 g, 87%) as a colorless oil; R_f = 0.42 (20% EtOAc/petroleum ether); IR (neat) 3423 cm^{-1} . Compound **15 β** had ^1H



NMR (300 MHz, CDCl_3) δ 1.06 [s, 9H, $(\text{CH}_3)_3\text{CSi}$], 1.33, 1.48 [both s, 3H ea, $\text{C}(\text{CH}_3)_2$], 3.66 (dd, $J = 2.6, 11.4$ Hz, 1H, H-5a), 3.83 (dd, $J = 2.6, 11.4$ Hz, 1H, H-5b), 4.29 (bs, 1H, H-4), 4.54 (d, $J = 10.6$ Hz, D_2O ex, 1H, OH), 4.61 (d, $J = 6.2$ Hz, 1H, H-3), 4.72 (d, $J = 5.9$ Hz, 1H, H-2), 5.38 (d, $J = 10.3$ Hz, 1H, H-1), 7.43–7.80 (m, 10H, Ph). ^{13}C NMR (75 MHz, CDCl_3) δ 25.3, 26.8 [$\text{C}(\text{CH}_3)_2$], 27.2 ($(\text{CH}_3)_3\text{CSi}$), 65.8, 82.0, 87.4, 87.5 (4C, C-2, 3, 4, 5), 103.6 (C-1), 112.3 [$\text{C}(\text{CH}_3)_2$], 128.0–135.9 (Ph). Compound **15 α** had ^1H NMR (300 MHz, CDCl_3) δ (selected signals) 1.06 [s, 9H, $(\text{CH}_3)_3\text{CSi}$], 1.41, 1.56 [both s, 3H ea, $\text{C}(\text{CH}_3)_2$], 3.94 (d, $J = 11.4$ Hz, 1H, D_2O ex, OH), 4.16 (bs, 1H, H-4), 5.67 (dd, $J = 4.0, 11.4$ Hz, 1H, H-1). ^{13}C NMR δ 25.1, 26.5 [$\text{C}(\text{CH}_3)_2$], 27.2 ($(\text{CH}_3)_3\text{CSi}$), 66.2, 79.8, 81.6, 82.2 (4C, C-2, 3, 4, 5), 98.2 (C-1), 113.3 [$\text{C}(\text{CH}_3)_2$]. Mixture **15 α/β** had HRMS(EI) m/z calcd for $\text{C}_{23}\text{H}_{29}\text{O}_5\text{Si}$ (M- CH_3) 413.1784. Found 413.1777.

1-*O*-Acetyl-4-*O*-*tert*-butyldiphenylsilyl-3-*O*-formyl-1,2-*O*-isopropylidene-D-erythro-hemiacetal (16). A solution of compound **15** (5.82 g, 13.6 mmol) in anhydrous cyclohexane (90 mL) containing diacetoxyiodosobenzene (4.39 g, 15.2 mmol) and iodine (3.45 g, 13.5 mmol) was stirred under an atmosphere of argon at rt for 4 h. The reaction mixture was then diluted with water and extracted with ether. The organic phase was washed with aq $\text{Na}_2\text{S}_2\text{O}_3$ and brine, then dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was separated by FCC to give major (4.00 g, 61%) and minor (1.08 g, 16%) isomers of **16**. The major isomer was isolated as a white solid; $R_f = 0.23$ (5% EtOAc/petroleum ether); IR (film) 1732, 1752 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.09 (s, 9H, $(\text{CH}_3)_3\text{CSi}$), 1.41, 1.50 [both s, 3H ea, $\text{C}(\text{CH}_3)_2$], 2.04 (s, 3H, CH_3CO), 3.91 (m, 2H, H-4a, 4b), 4.52 (dd, $J = 1.8, 5.9$ Hz, 1H, H-2), 5.18 (q, $J = 4.8$ Hz, 1H, H-3), 6.27 (d, $J = 1.8$ Hz, 1H, H-1), 7.38–7.70 (m 10H, Ph), 8.02 (s, 1H, HCO); ^{13}C NMR (75 MHz, CDCl_3) δ 21.9 (CH_3CO), 27.5, $(\text{CH}_3)_3\text{CSi}$, $\text{C}(\text{CH}_3)_2, 1\text{C}$, 28.1 [$\text{C}(\text{CH}_3)_2, 1\text{C}$], 62.9, 73.4, 81.2 (3C, C-2, 3, 4), 96.8 (C-1), 113.6 [$\text{C}(\text{CH}_3)_2$], 128.4, 130.4, 133.6, 136.2 (Ph), 160.3 (HCO), 170.2 (CH_3CO). HRMS (EI) m/z calcd for $\text{C}_{24}\text{H}_{31}\text{O}_5\text{Si}$ (M- $\text{C}_2\text{H}_3\text{O}_2$) 427.1941. Found 427.1932.

The minor isomer was isolated as a colorless oil; $R_f = 0.28$ (5% EtOAc/petroleum ether); ^1H NMR (300 MHz, CDCl_3) δ 1.09 (s, 9H), 1.42, 1.51 [both s, 3H ea, $\text{C}(\text{CH}_3)_2$], 2.04 (s, 3H, CH_3CO), 3.98 (m, 2H, H-4a,b), 4.54 (dd, $J = 3.3, 9.2$ Hz, 1H, H-2), 5.33 (m, 1H, H-3), 6.33 (d, $J = 3.3$ Hz, 1H, H-1), 7.40–7.70 (m 10H, Ph), 7.91 (s, 1H). ^{13}C NMR (75 MHz, CDCl_3) δ 21.3 (CH_3CO), 26.3 [$\text{C}(\text{CH}_3)_2, 1\text{C}$], 27.1 ($(\text{CH}_3)_3\text{CSi}$), 28.4 [$\text{C}(\text{CH}_3)_2, 1\text{C}$], 63.4, 71.3, 76.0 (3C, C-2, 3, 4), 93.5 (C-1), 112.7 [$\text{C}(\text{CH}_3)_2$], 127.8, 129.9, 133.3, 135.7 (Ph), 159.5 (HC = O), 170.0 ($\text{CH}_3\text{C} = \text{O}$). ESMS m/z 504.3 $[\text{M} + \text{NH}_4]^+$.

4-*O*-*tert*-Butyldiphenylsilyl-1,2-*O*-isopropylidene-D-erythro-S-phenyl monothiohemiacetal (13). $\text{BF}_3 \cdot \text{OEt}_2$ (1.60 mL, 12.7 mmol) was slowly added to a solution of **16** (4.10 g, 8.4 mmol) and thiophenol (2.16 mL, 21.1 mmol) in anhydrous CH_2Cl_2 (40 mL) at -78°C under an atmosphere of argon. The reaction was warmed to -40°C and stirred at this temperature for 1 h, or until TLC indicated complete disappearance of the starting material. Triethylamine (6.0 mL) was then added, and the reaction mixture was diluted with satd aq NaHCO_3 and extracted with ether. The organic phase was washed with brine, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The crude material was dissolved in methanol (50 mL) and treated with a solution of NaOMe in methanol at rt for 30 min. Most of the solvent was then removed under reduced pressure. FCC of the residue provided **13** (3.49 g, 81%) as a colorless oil; $R_f = 0.42$ (10% EtOAc/petroleum ether); IR (neat) 3485 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.07 (s, 9H,



(CH₃)₃CSi], 1.40, 1.53 [both s, 3H ea, C(CH₃)₂], 2.55 (d, J = 4.4 Hz, 1H, D₂O ex, OH), 3.81 (m, 3H, H-4a, 4b, H-3), 4.18 (t, J = 5.7 Hz, 1H, H-2), 5.58 (d, J = 5.5 Hz, 1H, H-1), 7.26–7.71 (m, 15H, Ph); ¹³C NMR (75 MHz, CDCl₃) δ 26.3 [C(CH₃)₂, 1C], 26.8 (CH₃)₃CSi), 27.1 (CH₃)₃CSi), 27.8 [C(CH₃)₂, 1C], 64.6, 72.4, 81.2 (3C, C-2, 3, 4), 86.0 (C-1), 111.7 [C(CH₃)₂], 127.2–135.6 (Ph). HRMS(EI) *m/z* calcd for C₂₃H₃₁O₄Si (M–SPh) 399.1992. Found 399.1996.

Methyl 2,3,6-tri-*O*-benzyl-4-*C*-(carboxymethyl)-4-deoxy- α -D-glucopyranoside (17). The ethyl ester of **17**^[27] (7.20 g, 10.0 mmol) was treated with a 1:1 mixture of aq 3N NaOH:ethanol (120 mL). After 4 h most of the ethanol was removed under reduced pressure. The resulting mixture was acidified with concentrated HCl to pH 4–5 and extracted with ethyl acetate. The combined organic layer was concentrated in vacuo. The residue was purified by FCC to give unreacted starting material and acid **17** (3.63g, 86% based on recovered starting material) as a colorless oil; *R*_f = 0.49 (40% EtOAc/petroleum ether); IR (film) 2500–3200 (br), 1707 cm^{–1}; ¹H NMR (300 MHz, CDCl₃) δ 2.03 (m, 2H, H-4, H-2'a), 2.57 (dd, J = 5.5, 16.5 Hz, 1H, H-2'b), 3.38 (s, 3H, OCH₃), 3.61, 3.83 (both m, 3H, 2H resp, H-2, 3, 5, 6a, 6b), 4.47 (d, J = 11.7 Hz, 1H, PhCH), 4.57–4.81 (m, 5H, H-1, 4 × PhCH), 5.02 (d, J = 11.0 Hz, 1H, PhCH), 7.20–7.35 (m, 15H, Ph). ¹³C NMR (75 MHz, CDCl₃) δ 32.9, 41.2 (C-4, 2'), 56.1 (OCH₃), 70.4, 70.9, 73.7, 74.2, 76.0, 78.6, 82.4 (7C, C-2, 3, 5, 6, PhCH₂), 99.2 (C-1), 128.2–139.3 (Ph), 178.7 (C = O). HRMS(FAB) *m/z* calcd for C₃₀H₃₄O₇Na [M + Na]⁺ 529.2201. Found 529.2202.

Methyl 2,3,6-tri-*O*-benzyl-[(4-*C*-(4-*O*-*tert*-butyldiphenylsilyl-1,2-*O*-isopropylidene-D-erythro-*S*-phenyl monothiohemiacetal)-2-ethanoate]-4-deoxy- α -D-glucopyranoside (18). DCC (2.71 g, 13.15 mmol) was added at 0°C to a solution of alcohol **13** (2.67 g, 5.26 mmol), acid **17**, (3.63 g, 7.17 mmol), and DMAP (128 mg, 1.05 mmol) in anhydrous benzene (50 mL). The reaction was warmed to rt and stirred for 2 h. The mixture was then diluted with ether (20 mL) and filtered. The filtrate was washed with 0.1 N aq HCl and brine, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residue was purified by FCC to give **18** (4.47 g, 85%) as a colorless oil; *R*_f = 0.42 (20% EtOAc/petroleum ether); IR (neat) 1734 cm^{–1}; ¹H NMR (500 MHz, CDCl₃) δ 1.00 [s, 9H, (CH₃)₃CSi], 1.31, 1.42 [both s, 3H ea, C(CH₃)₂], 2.26 (m, 2H, H-4, 2'a), 2.55 (m, 1H, H-2'b), 3.10 (s, 3H, OCH₃), 3.33 (m, 1H, H-1), 3.52 (m, 2H, H-6a, 6b), 3.79 (m, 4H, H-3, 5, 4a, 4b), 4.31 (t, J = 6.0 Hz, 1H, H-2''), 4.39 (d, J = 12.0 Hz, 1H, PhCH), 4.51 (m, 3H, 3 × PhCH), 4.61 (d, J = 11.0 Hz, 1H, PhCH), 4.62 (d, J = 4.0 Hz, 1H, H-1), 4.89 (m, 1H, H-3''), 4.91 (d, J = 11.0 Hz, 1H, PhCH), 5.37 (d, J = 6.5 Hz, 1H, H-1''), 7.10–7.70 (m, 30 H, Ph); ¹³C NMR (75 MHz, CDCl₃) δ 26.7 [C(CH₃)₂, 1C], 27.6 (CH₃)₃CSi), 28.1 [C(CH₃)₂, 1C], 33.5 (C-2'), 41.2 (C-4), 55.9 (OCH₃), 62.8 (C-4''), 70.6, 71.1, 73.5, 74.1, 74.4, 75.9, 78.9, 79.5, 82.5 (9C, C-2, 3, 5, 6, 2'', 3'', PhCH₂), 85.6 (C-1''), 99.1 (C-1), 112.2 [C(CH₃)₂], 127.8–139.3 (Ph), 171.6 (C = O). HRMS (FAB) *m/z* calcd for C₅₉H₆₈O₁₀SSiNa [M + Na]⁺ 1019.4203. Found 1019.4200.

Methyl 2,3,6-tri-*O*-benzyl-4-*C*-[(4-*O*-*tert*-butyldiphenylsilyl-1,2-*O*-isopropylidene-D-erythro-*S*-phenyl monothiohemiacetal)-2-propene]-4-deoxy- α -D-glucopyranoside (19). A solution of Tebbe reagent^{[31,32]d} in THF (9.3 mL, 0.5 M, 4.7 mmol),

^dThe Tebbe reagent used in this research was purchased from Sigma-Aldrich Inc.



was added dropwise, under an atmosphere of argon at -78°C , to ester **18** (1.82 g, 1.86 mmol) and pyridine (0.15 mL) in anhydrous 3:1 toluene/THF (20 mL). The reaction mixture was warmed to rt, stirred for 2 h, and then slowly poured into a solution of 1 N aq NaOH at 0°C . The resulting suspension was extracted with ether, and the organic phase was washed with brine, dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by FCC on basic alumina (Brockmann I, 150 mesh) to give enol ether **18** (1.09 g, 60%) as a colorless oil, $R_f = 0.61$ (15% EtOAc/petroleum ether); ^1H NMR (300 MHz, C_6D_6) δ 1.10 [s, 9H, $(\text{CH}_3)_3\text{CSi}$], 1.47, 1.62 [both s, 3H ea, $\text{C}(\text{CH}_3)_2$], 2.30 (dd, $J = 4.8, 14.5$ Hz, 1H, H-2'a), 2.40 (m, 1H, H-4), 2.69 (dd, $J = 3.6, 14.5$ Hz, 1H, H-2'b), 3.16 (s, 3H, OCH_3), 3.54 (dd, $J = 3.5, 9.7$ Hz, 1H, H-2), 3.75–4.15, 4.28 (both m, 8H, 1H, resp, H-3, 5, 6a, 6b, 3'', 4''a, 4''b, = CH_2), 4.30–4.65 (m, 4H, $2 \times \text{PhCH}_2$), 4.74 (m, 2H, H-1, PhCH), 4.88 (bt, $J = 5.0$ Hz, 1H, H-2''), 5.19 (d, $J = 11.2$ Hz, 1H, PhCH), 6.10 (d, $J = 6.2$ Hz, 1H, H-1''), 6.94–7.84 (m, 30H, Ph); ^{13}C NMR (75 MHz, C_6D_6) δ 26.8 [$\text{C}(\text{CH}_3)_2$, 1C], 27.6 $(\text{CH}_3)_3\text{CSi}$, 28.2 [$\text{C}(\text{CH}_3)_2$, 1C], 33.8 (C-2'), 42.1 (C-4), 55.2 (OCH_3), 62.3 (C-4''), 71.5, 71.9, 72.7, 74.1, 75.2, 77.4, 78.6, 81.4, 83.8, 85.7 (10C, C-2, 3, 5, 6, 2'', 3'', = CH_2 , PhCH_2) 86.1 (C-1''), 98.7 (C-1), 112.4 [$\text{C}(\text{CH}_3)_2$], 127.5–140.7 (Ph), 160.1 ($\text{OC} = \text{CH}_2$). HRMS (FAB) m/z calcd for $\text{C}_{60}\text{H}_{71}\text{O}_9\text{SSiH}$ [$\text{M} + \text{H}$] $^+$ 995.4587. Found 995.4588.

Methyl 4-C-(2,6-anhydro-7-O-tert-butylidiphenylsilyl-4,5-O-isopropylidene-1,3-dideoxy-D-ribo-hept-2-enitol-1-C-yl)-2,3,6-tri-O-benzyl-4-deoxy- α -D-glucopyranoside (20). A mixture of enol ether **19** (740 mg, 0.08 mmol), 2,6-di-tert-butyl-4-methylpyridine (1.54 g, 7.51 mmol), and freshly activated, powdered 4 Å molecular sieves (300 mg) in anhydrous CH_2Cl_2 (25 mL) was stirred for 15 min at rt under an argon atmosphere, then cooled to 0°C . Methyl triflate (1.0 mL, 7.55 mmol) was introduced, and the mixture was warmed to rt and stirred for an additional 18 h, or until all the starting material had disappeared, at which time triethylamine (2.5 mL) was added. The mixture was diluted with ether, washed with satd aq NaHCO_3 and brine, dried (Na_2SO_4), filtered and concentrated under reduced pressure. The residue was purified by FCC over basic alumina (Brockmann I, 150 mesh) to give **20** (460 mg, 70%) as a clear oil, $R_f = 0.60$ (20% EtOAc/petroleum ether); IR (film) 1647 cm^{-1} ; ^1H NMR (300 MHz, C_6D_6) δ 1.24 [s, 9H, $(\text{CH}_3)_3\text{CSi}$], 1.31, 1.37 [both s, 3H ea, $\text{C}(\text{CH}_3)_2$], 2.40 (dd, $J = 5.7, 14.4$ Hz, 1H, H-1'a), 2.56 (m, 2H, H1'b, 4), 3.24 (s, 3H, OCH_3), 3.70 (m, 2H, H-2, 7'a), 3.85–4.00–4.20 (both m, 2H, 5H resp., H-3, 5, 6a, 6b, 3', 6', 7'b), 4.27 (t, $J = 5.0$ Hz, 1H, H-5'), 4.60 (m, 4H, $2 \times \text{PhCH}_2$), 4.83 (m, 2H, H-1, PhCH), 4.93 (d, $J = 5.6$ Hz, 1H, H-4'), 5.27 (d, $J = 11.7$ Hz, 1H, PhCH), 7.11–7.46 and 7.90 (both m, 25H, Ph); ^{13}C NMR (75 MHz, C_6D_6) δ 26.8 [$\text{C}(\text{CH}_3)_2$, 1C], 27.7 $(\text{CH}_3)_3\text{CSi}$, 28.2 [$\text{C}(\text{CH}_3)_2$, 1C], 33.9 (C-1'), 42.2 (C-4), 55.3 (OCH_3), 62.4 (C-7'), 71.6, 72.0, 72.8, 74.2, 75.3, 77.5, 78.7, 81.6, 83.9, 85.7, 86.2 (11C, C2, 3, 5, 6, 3', 4', 5', 6', $3 \times \text{PhCH}_2$), 100.0 (C-1), 112.5 [$\text{C}(\text{CH}_3)_2$], 128.0–140.8 (Ph), 160.2 (C-2'). HRMS (FAB) m/z calcd for $\text{C}_{54}\text{H}_{65}\text{O}_9\text{Si}$ [$\text{M} + \text{H}$] $^+$ 885.4399. Found 885.4398.

Methyl 4-C-(2,6-anhydro-4,5-O-isopropylidene-7-O-tert-butylidiphenylsilyl-1-deoxy-D-glycero-D-manno-heptitol-1-C-yl)-2,3,6-tri-O-benzyl-4-deoxy- α -D-glucopyranoside (21) and Methyl 4-C-(2,6-anhydro-4,5-O-isopropylidene-7-O-tert-butylidiphenylsilyl-1-deoxy-D-glycero-D-allo-heptitol-1-C-yl)-2,3,6-tri-O-benzyl-4-deoxy- α -D-glucopyranoside (22). $\text{BH}_3 \cdot \text{Me}_2\text{S}$ (2.7 mL of 1 M solution in CH_2Cl_2 , 2.7 mmol) was added at 0°C to a solution of the glycol **20** (340 mg, 0.39 mmol) in anhydrous



THF (15 mL) under an atmosphere of argon. The mixture was warmed to rt and stirred for an additional 1 h. At that time the solution was cooled to 0°C and treated with a mixture of 3N NaOH (2.5 mL) and 30% aqueous H₂O₂ (2.5 mL) for 30 min. The mixture was diluted with ether, washed with satd aq NaHCO₃ and brine, dried (Na₂SO₄), filtered and concentrated under reduced pressure. FCC of the residue afforded a mixture of **21** and **22** (264 mg, ca. ratio 5:1, 84%) which had identical TLC mobilities (R_f = 0.25, 10% EtOAc/petroleum ether). For characterization purposes, a pure sample of **21** was obtained by resubjecting this material to a second chromatography and collection of a later eluting fraction. Compound **21** was isolated as a colorless oil; ¹H NMR (300 MHz, CDCl₃) δ 1.07 [s, 9H, (CH₃)₃CSi], 1.29, 1.46 [both s, 3H ea, C(CH₃)₂], 1.51 (m, 1H, H-1'a), 1.75 (m, 1H, H-1'b), 2.05 (m, 1H, H-4), 3.00 (bs, 1H, D₂O ex), 3.32 (m, 1H), 3.37 (s, 3H), 3.41 (m, 1H), 3.55 (m, 2H), 3.60–3.95 (m, 7H), 4.12 (m, 1H), 4.44 (s, 2H), 4.65–4.85 (m, 4H), 5.03 (d, J = 12.0 Hz, 1H), 7.26–7.78 (m, 25H); ¹³C NMR (75 MHz, CDCl₃) δ 27.7 (CH₃)₃CSi, C(CH₃)₂, 1C), 28.8 [C(CH₃)₂, 1C], 33.1 (C-1'), 40.5 (C-4), 55.9 (OCH₃), 65.4 (C-7'), 71.4, 72.4, 73.5, 74.0, 74.6, 75.5, 76.6, 78.8, 81.2, 82.6 (12C, C-2, 3, 5, 6, 2', 3', 4', 5' 6', 3 \times PhCH₂), 99.0 (C-1), 109.5 [C(CH₃)₂], 128.1–139.1 (Ph). HRMS (FAB) m/z calcd for C₅₄H₆₆O₁₀SiNa [M + Na]⁺ 925.4321. Found 925.4323. Compound **22** was characterized as triol **24** and triacetate **26** (vide infra).

Methyl 4-C-(2,6-anhydro-7-O-tert-butyldiphenylsilyl-1-deoxy-D-glycero-D-manno-heptitol-1-C-yl)-2,3,6-tri-O-benzyl-4-deoxy- α -D-glucopyranoside (23) and Methyl 4-C-(2,6-anhydro-7-O-tert-butyldiphenylsilyl-1-deoxy-D-glycero-D-allo-heptitol-1-C-yl)-2,3,6-tri-O-benzyl-4-deoxy- α -D-glucopyranoside (24). To a portion of the mixture of **21** and **22** from the previous step (103 mg, 0.11 mmol) in dry methanol was added a 1 M solution of HCl in ether (0.7 mL). The reaction was stirred for 1.5 h and then neutralized with a solution of NaOMe in methanol. Removal of the volatiles followed by FCC of the residue provided triols **23** (65 mg, 66%) and **24** (12 mg, 12%). Compound **23** was isolated as a clear gum; R_f = 0.50 (50% EtOAc/petroleum ether); ¹H NMR (300 MHz, C₆D₆) δ 1.18 [s, 9H, (CH₃)₃CSi], 1.82 (m, 1H, H-1'a), 1.98 (m, 1H, H-1'b), 2.32 (m, 1H, H-4), 2.75 (s, 1H, D₂O ex, OH), 2.92 (s, 1H, D₂O ex, OH), 3.22 (s, 3H, OCH₃), 3.51 (s, 1H, D₂O ex, OH), 3.52 (dd, J = 3.3, 9.9 Hz, partially hidden by s at δ 3.51, 1H), 3.60 (dd, J = 2.6, 8.4 Hz, 1H), 3.72 (m, 2H), 3.82 (m, 4H), 3.93 (dd, J = 6.3, 9.3 Hz, 1H), 4.01 (t, J = 9.6 Hz, 1H), 4.09 (bs, 1H), 4.26 (t, J = 6.2 Hz, 1H), 4.44 (ABq, $\Delta\delta$ = 0.06 ppm, J = 11.7 Hz, 1H), 4.49 (ABq, $\Delta\delta$ = 0.05 ppm, J = 12.4 Hz, 1H), 4.74 (d, J = 3.3 Hz, 1H, H-1), 4.87 (ABq, $\Delta\delta$ = 0.41 ppm, J = 10.6 Hz, 1H), 7.10–7.76 (m, 25H); ¹³C NMR (75 MHz, CDCl₃) δ 27.2 (CH₃)₃CSi], 33.4 (C-1'), 40.0 (C-4), 55.4 (OCH₃), 62.9 (C-7'), 69.0, 69.1, 70.8, 72.0, 72.9, 73.6, 73.7, 73.8, 74.4, 76.5, 81.3, 82.2, (12C, C-2, 3, 5, 6, 2', 3', 4' 5' 6', 3 \times PhCH₂), 98.4 (C-1), 127.6–138.4 (Ph). HRMS (FAB) m/z calcd for C₅₁H₆₂O₁₀NaSi [M + Na]⁺ 885.4010. Found 885.4013.

Compound **24** was isolated as a clear gum; R_f = 0.64 (50% EtOAc/petroleum ether); ¹H NMR (500 MHz, C₆D₆) δ 1.16 [s, 9H, (CH₃)₃CSi], 1.68 (m, 1H, H-1'a), 2.20 (m, 1H, H-1'b), 2.42 (m, 1H, H-4), 2.44 (bs, 1H, D₂O ex, OH), 2.58 (d, J = 7.5 Hz, 1H, D₂O ex, OH), 2.93 (d, J = 5.5 Hz, 1H, D₂O ex, OH), 3.07 (t, J = 7.3 Hz, 1H), 3.18 (s, 3H, OCH₃), 3.60 (m, 3H), 3.80 (m, 3H), 3.88–4.00 (m, 4H), 4.04 (dd, J = 4.5, 11.0 Hz, 1H), 4.42 (ABq, J = 12.0 Hz, $\Delta\delta$ = 0.05 ppm, 2H, PhCH₂), 4.48 (ABq, $\Delta\delta$ = 0.04 ppm, J = 12.0 Hz, 2H, PhCH₂), 4.76 (d, J = 3.0 Hz, 1H, H-1), 4.89 (ABq, $\Delta\delta$ = 0.44 ppm,



$J = 11.0$ Hz, 2H, PhCH_2), 7.05–7.82 (m, 25H); HRMS (FAB) m/z calcd for $\text{C}_{51}\text{H}_{62}\text{O}_{10}\text{NaSi}$ $[\text{M} + \text{Na}]^+$ 885.4010. Found 885.4030.

Methyl 4-*C*-(2,6-anhydro-1-deoxy-*D*-glycero-*D*-manno-heptitol-1-*C*-yl)-2,3,6-tri-*O*-benzyl-4-deoxy- α -*D*-glucopyranoside (25). A solution of HCl in ether (0.3 mL of ca. 1M) was added to **21** (419 mg, 0.49 mmol) in dry methanol (30 mL). The reaction was stirred at rt for approximately 4 h, then neutralized with a solution of NaOMe in methanol. Addition of Bu_4NF (2.8 mL of a 1 M solution in THF, 2.8 mmol) to the reaction mixture, followed by stirring for an additional 1 h, then removal of the volatiles under reduced pressure, and FCC of the residue, provided **25** (164 mg, 58%) as a colorless oil; $R_f = 0.37$, (20% acetone/EtOAc); ^1H NMR (300 MHz, CDCl_3) δ 1.62 (m, 1H, H-1'a), 1.77 (m, 1H, H-1'b), 2.07 (m, 1H, H-4), 2.86 (bs, 1H, D_2O ex, OH), 3.29 (m, 2H), 3.38 (s, 3H, OCH_3), 3.43–3.67 (m, 8H), 3.85 (m, 2H), 4.50–4.76 (m, 6H, H-1, 5 \times PhCH), 5.03 (d, $J = 10.6$ Hz, 1H, PhCH), 7.24–7.37 (m, 15H, Ph); ^{13}C NMR (75 MHz, CDCl_3) δ 32.3 (C-1'), 39.4 (C-4), 55.5 (OCH_3), 59.4 (C-7'), 68.6, 70.6, 71.5, 72.3, 72.5, 73.0, 73.4, 73.6, 76.3, 77.5, 81.6, 82.0 (12C, C-2, 3, 5, 6, 2', 3', 4', 5', 6', 3 \times PhCH_2), 98.3 (C-1), 127.7–138.2 (Ph). HRMS (FAB) m/z calcd for $\text{C}_{35}\text{H}_{44}\text{O}_{10}\text{Na}$ $[\text{M} + \text{Na}]^+$ 647.2832. Found 647.2834.

Methyl 4-*C*-(2,6-anhydro-3,4,5,7-tetra-*O*-acetyl-1-deoxy-*D*-glycero-*D*-manno-heptitol-1-*C*-yl)-2,3,6-tri-*O*-benzyl-4-deoxy- α -*D*-glucopyranoside (26). A solution of tetraol **25** (25.5 mg, 0.04 mmol), DMAP (1.00 mg, 4 μmol), and acetic anhydride (0.04 mL, 0.40 mmol) in ethyl acetate (3.0 mL), was stirred at rt for 20 min. Methanol was then added and the reaction mixture concentrated under reduced pressure. The residue was purified by FCC to give **26** (26.7 mg, 82%) as a colorless oil; $R_f = 0.66$ (40% EtOAc/petroleum ether); $[\alpha]_D + 23^\circ$ (c 1.3 CHCl_3); IR (film) 1743 cm^{-1} ; ^1H NMR (500 MHz, C_6D_6) δ 1.59 (s, 3H, CH_3CO), 1.64, (m, buried under s at δ 1.64, 1H, H-1'a) 1.64, 1.69, 1.78 (all s, 3H ea, 3 \times CH_3CO), 2.12 (t, $J = 10.5$ Hz, 1H, H-1'b), 2.23 (apparent q, $J = 7.5$ Hz, 1H, H-4), 3.25 (s, 3H, OCH_3), 3.56 (bd, $J = 9.0$ Hz, 1H, H-2), 3.62 (m, 1H, H-6a), 3.70 (bd, $J = 10.5$ Hz, 1H, H-6b), 3.81 (bd, $J = 10.5$ Hz, 1H, H-5), 3.89 (bd, $J = 11.0$ Hz, 1H, H-7'a), 4.06 (m, 2H, H-3, 6'), 4.28 (m, 2H, H-2', 7'b), 4.43 (m, 4H, 2 \times PhCH_2), 4.74 (bs, 1H, H-1), 4.94 (ABq, $\Delta\delta = 0.49$ ppm, $J = 10.5$ Hz, 2H, PhCH_2), 5.33 (t, $J = 8.5$ Hz, 1H, H-3'), 5.43 (dd, $J = 2.5, 8.5$ Hz, 1H, H-4'), 5.49 (d, $J = 2.5$ Hz, 1H, H-5'), 7.10–7.51 (m, 15H, Ph); ^{13}C NMR (75 MHz, C_6D_6) δ 20.7, 20.8, 20.9 (CH_3CO , 4C), 31.8 (C-1'), 41.1 (C-4), 55.5 (OCH_3), 62.3, 69.5, 70.6, 71.7, 72.0, 72.3, 72.9, 73.3 (2C), 74.2, 75.8, 81.4, 83.5 (13C, C-2, 3, 5, 6, 2', 3', 4', 5', 6', 7', 3 \times PhCH_2), 99.0 (C-1) 128.0–130.0 (Ph), 169.6, 169.7, 169.8, 170.1 (CH_3CO). HRMS (FAB) m/z calcd for $\text{C}_{43}\text{H}_{52}\text{O}_{14}\text{Na}$ $[\text{M} + \text{Na}]^+$ 815.8355. Found 815.3256.

Methyl 4-*C*-(2,6-anhydro-7-*O*-tert-butyldiphenylsilyl-3,4,5-tri-*O*-acetyl-1-deoxy-*D*-glycero-*D*-allo-heptitol-1-*C*-yl)-2,3,6-tri-*O*-benzyl-4-deoxy- α -*D*-glucopyranoside (27). Triol **24** (4.0 mg, 0.005 mmol) was subjected to the identical acetylation procedure that was described for the preparation of **26**. FCC of the crude reaction product provided **27** (4.2 mg, 91%) as a clear gum; $R_f = 0.50$ (30% EtOAc/petroleum ether); ^1H NMR (500 MHz, C_6D_6) δ 1.72 [s, 9H, $(\text{CH}_3)_3\text{CSi}$], 1.50, 1.63, 1.64 (all s, 3H ea, CH_3CO), 1.75 (m, 1H, H-1'a), 2.15 (bdd, $J = 4.5, 14.0$ Hz, 1H, H-1'b), 2.62 (m, 1H, H-4), 3.16 (s, 3H, OCH_3), 3.70 (dd, $J = 3.0, 8.8$ Hz, 1H, H-2), 3.76 (dd, $J = 3.5, 11.5$



Hz, 1H, H-7'a), 3.84 (m, 2H, H-6a, 6'), 3.88 (bd, $J = 11.5$ Hz, 1H, H-7'b), 3.98 (dd, $J = 3.5, 11.0$ Hz, 1H, H-6b), 4.12, (t, $J = 10.5$ Hz, 1H, H-3), 4.23 (m, 2H, H-5, H-2'), 4.44 (ABq, $\Delta\delta = 0.03$ ppm, $J = 12.0$ Hz, 2H, PhCH_2), 4.54 (ABq, $\Delta\delta = 0.07$ ppm, $J = 12.0$ Hz, 2H, PhCH_2), 4.81, (d, $J = 3.0$ Hz, 1H, H-1), 4.90 (dd, $J = 2.0, 9.5$ Hz, 1H, H-3'), 4.99 (ABq, $\Delta\delta = 0.55$ ppm, $J = 11.5$ Hz, 2H, PhCH_2), 5.43 (dd, $J = 2.0, 9.9$ Hz, 1H, H-5'), 6.20 (bt, $J = 2.0$ Hz, 1H, H-4'), 7.20, 7.80 (both m, 25H, Ph); ^{13}C NMR (75 MHz, CDCl_3) δ 20.7, 20.8, 20.9 (CH_3CO), 27.2, 27.6 ($(\text{CH}_3)_3\text{CSi}$), 30.0 (C-1'), 40.4 (C-4), 55.3 (OCH_3), 63.1, 64.5, 66.8, 68.8, 70.3, 70.7, 71.0, 71.8, 73.1, 73.8, 74.7, 75.0, 82.5 (13C, C-2, 3, 5, 6, 2', 3', 4', 5', 6', 7', $3 \times \text{PhCH}_2$), 98.6 (C-1), 127.4–139.3 (Ph), 168.9, 169.3, 169.9 (CH_3CO). ESMS m/z 1006.5 $[\text{M} + \text{NH}_4]^+$.

Methyl 4-C-(2,6-anhydro-1-deoxy-D-glycero-D-manno-heptitol-1-C-yl)-4-deoxy- α -D-glucopyranoside (2). A mixture of **25** (124 mg, 0.20 mmol), 20% Pd on carbon (400 mg), formic acid (0.35 mL) and methanol (7.0 mL) was stirred under an atmosphere of hydrogen (balloon), for 12 h. The reaction mixture was purged with argon and filtered through a bed of Celite. The filtrate was concentrated in vacuo, and the residue was purified by FCC to provide **2** (55.0 mg, 83%) as a white solid; $R_f = 0.68$ (50% MeOH/ CHCl_3); $[\alpha]_D + 93^\circ$ (c 0.54 MeOH); ^1H NMR (500 MHz, D_2O , 50°C) δ 1.83 (ddd, $J = 2.5, 10.0, 13.5$ Hz, 1H, H-1'a), 1.93 (m, 1H, H-4), 2.08 (dd, $J = 6.0, 13.5$ Hz, 1H, H-1'b), 3.53 (s, 3H, OCH_3), 3.61 (bd, $J = 9.5$ Hz, 1H, H-7'a), 3.64 (dd, $J = 4.0, 9.0$ Hz, 1H, H-2), 3.79–3.86 (m, 4H, H-5, 2', 3', 6'), 3.92* (m, 3H, H-4', 7'a, 7'b), 4.01* (bdd, $J = 8.0, 11.5$ Hz, H-6a), 4.08* (m, 2H, H-6b, 5'), 4.95 (d, $J = 3.5$ Hz, H-1); ^{13}C NMR (75 MHz, D_2O) δ 33.8 (C-1'), 42.8 (C-4), 58.6 (OCH_3), 62.1, 65.3 (C-6, 7'), 72.2, 74.7, 75.2, 75.5, 76.1, 76.2, 76.6, 81.8 (8C, C-2, 3, 5, 2', 3', 4', 5', 6'), 103.2 (C-1). HRMS (FAB) m/z calcd for $\text{C}_{14}\text{H}_{27}\text{O}_{10}\text{Na}$ $[\text{M} + \text{Na}]^+$ 355.1604. Found 355.1602.

ACKNOWLEDGMENT

We thank the National Institutes of Health (NIH), General Medical Sciences (GM 57865) for their support of this research. Research Centers in Minority Institutions (RCMI), award RR-03037 from the National Center for Research Resources of the NIH, which supports the infrastructure (and instrumentation) of the Chemistry Department at Hunter, is also acknowledged.

REFERENCES

1. Compain, P.; Martin, O.R. Carbohydrate mimetics-based glycosyltransferase inhibitors. *Bioorg. Med. Chem. Lett.* **2001**, 9, 3077–3092.
2. Chapleur, Y. *Carbohydrate Mimics, Concepts and Methods*; Wiley-VCH: Weinheim, 1998.
3. Palcic, M.M.; Heerze, L.D.; Srivastava, O.P.; Hindsgaul, O. A bisubstrate analog inhibitor for $\alpha(1\rightarrow2)$ -fucosyl transferase. *J. Biol. Chem.* **1989**, 264, 17174–17181.
4. Yuasa, H.; Palcic, M.M.; Hindsgaul, O. Synthesis of the carbocyclic analog of



- uridine 5'-(α -D-galactospyranosyl diphosphate) (UDP-Gal) as an inhibitor of β -(1 \rightarrow 4)-galactosyl transferase. *Can. J. Chem.* **1995**, *73*, 2190–2195.
5. Schmidt, R.R.; Frische, K. A new galactosyl transferase inhibitor. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 1747–1750.
 6. Hashimoto, H.; Endo, T.; Kajihara, Y. Synthesis of the first tricomponent bisubstrate analogue that exhibits potent inhibition against GlcNAc: β -1,4-galactosyltransferase. *J. Org. Chem.* **1997**, *62*, 1914–1915.
 7. Waldscheck, B.; Streiff, M.; Notz, W.; Kinzy, W.; Schmidt, R.R. Galactosyl transferase inhibition based on a new type of disubstrate analogue. *Angew. Chem., Int. Ed. Engl.* **2001**, *40*, 4007–4011.
 8. Heskamp, B.M.; Veeneman, G.H.; van der Marel, G.A.; van Boeckel, C.A.A.; van Boom, J.H. Synthesis and glycosylating properties of ketopyranosyl donors. *Tetrahedron* **1995**, *51*, 5657–5670.
 9. Pasquarello, C.; Picasso, S.; Demange, R.; Malissard, M.; Berger, E.G.; Vogel, P.J. The C-disaccharide α -C-(1 \rightarrow 3)-mannopyranoside of N-acetylgalactosamine is an inhibitor of glycohydrolases and of human α -(1 \rightarrow 3)-fucosyltransferase VI. Its epimer α -(1 \rightarrow 3)-mannopyranoside of N-acetyltalosamine is not. *J. Org. Chem.* **2000**, *65*, 4251–4260.
 10. Qiao, L.; Murray, B.W.; Shimazaki, M.; Schultz, J.; Wong, C.-H. Synergistic inhibition of human α -1,3-fucosyltransferase V. *J. Am. Chem. Soc.* **1996**, *118*, 7653–7662.
 11. Jeffries, I.; Bowen, B.R. Synthesis of inhibitors of α -1,3-fucosyltransferase. *Bioorg. Med. Chem. Lett.* **1997**, *7*, 1171–1174.
 12. Asensio, J.L.; Cañada, F.J.; García-Herrero, A.; Murillo, M.T.; Fernández-Mayoralas, J.B.A.; Kozak, J.; Zhu, Z.; Johnson, C.R.; Jiménez-Barbero, J. Conformational behavior of aza-C-glycosides: experimental demonstration of the relative role of the exo-anomeric effect and 1,3-type interactions in controlling the conformation of regular glycosides. *J. Am. Chem. Soc.* **1999**, *121*, 11318–11329.
 13. Montero, E.; García-Herrero, J.L.; Asensio, H.K.; Ogawa, S.; Santoyo-González, C.F.J.; Jiménez-Barbero, J. Conformational behaviour of non-hydrolyzable lactose analogues: the thioglycoside, carboglycoside, and carba-iminoglycoside cases. *Eur. J. Org. Chem.* **2000**, 1945–1952. For reviews on C-glycoside synthesis, see Refs. [14–21].
 14. Du, Y.; Linhardt, R.J.; Vlahov, J.R. Recent advances in stereoselective C-glycoside synthesis. *Tetrahedron* **1998**, *54*, 9913–9959.
 15. Togo, H.; He, W.; Waki, Y.; Yokoyama, M. C-Glycosidation technology with free radical reactions. *Synlett* **1998**, 700–717.
 16. Beau, J.-M.; Gallagher, T. Nucleophilic C-glycosyl donors for C-glycoside synthesis. *Top. Curr. Chem.* **1997**, *187*, 1–54.
 17. Skrydstrup, T.; Vauzeilles, B.; Beau, J.-M. Synthesis of C-oligosaccharides. In *Carbohydrates in Chemistry and Biology*; Ernst, B., Hart, G.W., Sinaÿ, P., Eds.; Wiley-VCH: Weinheim, 2000; Vol. 1, 495–530.
 18. Beau, J.-M.; Vauzeilles, B. C-glycosyl analogs of oligosaccharides and glycosyl amino acids. In *Glycoscience: Chemistry and Chemical Biology*; Fraser-Reid, B., Tatsuta, K., Thiem, J., Eds.; Springer: Heidelberg, 2001; Vol. 3, 2679–2724.
 19. Postema, M.H.D. *C-Glycoside Synthesis*; CRC Press: Boca Raton, 1995.



20. Liu, L.; McKee, M.; Postema, M.H.D. Synthesis of C-saccharides. *Curr. Org. Chem.* **2001**, *5*, 1133–1167.
21. Levy, D.; Tang, C. *The Synthesis of C-Glycosides*; Elsevier/Pergamon: Oxford, 1995.
22. Khan, N.; Cheng, X.; Mootoo, D.R. 1-Thio-1,2-*O*-isopropylidene acetals: novel precursors for the synthesis of complex C-glycosides. *J. Am. Chem. Soc.* **1999**, *121*, 4918–4919.
23. Cheng, X.; Khan, N.; Mootoo, D.R. Synthesis of the C-glycoside analogue of a novel sialyl Lewis X mimetic. *J. Org. Chem.* **2000**, *65*, 2544–2547.
24. Brill, G.M.; Jackson, M.; Whittern, D.N.; Buko, A.M.; Hill, P.; Chen, R.H.; McAlpine, J.B. Altromycins E, F, G, H and I; additional novel components of the altromycin complex. *J. Antibiot.* **1994**, *47*, 1160–1164.
25. Kaskar, B.; Heise, G.L.; Michalak, R.S.; Vishnuvajjala, B.R. Convergent large scale synthesis of protected D-ribonolactone from D-ribose. *Synthesis* **1990**, 1031–1032.
26. Armas, P.D.; Francisco, C.G.; Suarez, E. Reagents with hypervalent iodine: formation of convenient chiral synthetic intermediates by fragmentation of carbohydrate anomeric alkoxy radicals. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 772–773.
27. Wang, Y.; Babirad, S.A.; Kishi, Y. Preferred conformation of C-glycosides. 8. Synthesis of 1,4-linked carbon disaccharides. *J. Org. Chem.* **1992**, *57*, 468–481.
28. Liu, L.; Postema, M.H.D. A unified approach to differentially linked C-disaccharides by ring-closing metathesis. *J. Am. Chem. Soc.* **2001**, *123*, 8602–8603.
29. Immel, S.; Fujita, K.; Lichtenthaler, F.W. Solution geometries and lipophilicity patterns of α -cycloaltrin. *Chem. Eur. J.* **1999**, *5*, 3185–3192.
30. Dowd, M.K.; French, A.D.; Reilly, P.J. Modeling of aldopyranosyl ring puckering with MM3 (92). *Carbohydr. Res.* **1994**, *264*, 1–19.
31. Pine, S.H.; Pettit, R.J.; Geib, G.D.; Cruz, S.G.; Gallego, C.H.; Tijerina, T.; Pine, R.D. Carbonyl methylenation using a titanium-aluminum (Tebbe) complex. *J. Org. Chem.* **1985**, *50*, 1212–1218.
32. Pine, S.H.; Shen, G.S.; Hoang, H. Ketone methylenation using the Tebbe and Wittig reagents. A comparison. *Synthesis* **1991**, 165–167.

Received March 11, 2003

Accepted July 24, 2003

