This article was downloaded by: [Stony Brook University] On: 01 November 2014, At: 23:44 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lsyc20</u>

A CONVENIENT SYNTHESIS OF 3,3-DIMETHYLBUTYRALDEHYDE

Zhi Guo^a, Rachel Sawyer^a & Indra Prakash^b

^a The NutraSweet Company , Research and Development, 601 E. Kensington Road, Mt. Prospect, Illinois, 60056, U.S.A.

^b The NutraSweet Company, Research and Development, 601 E. Kensington Road, Mt. Prospect, Illinois, 60056, U.S.A.

Published online: 20 Aug 2006.

To cite this article: Zhi Guo , Rachel Sawyer & Indra Prakash (2001) A CONVENIENT SYNTHESIS OF 3,3-DIMETHYLBUTYRALDEHYDE, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 31:21, 3395-3399, DOI: <u>10.1081/SCC-100106050</u>

To link to this article: http://dx.doi.org/10.1081/SCC-100106050

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

SYNTHETIC COMMUNICATIONS, 31(21), 3395–3399 (2001)

A CONVENIENT SYNTHESIS OF 3,3-DIMETHYLBUTYRALDEHYDE

Zhi Guo, Rachel Sawyer, and Indra Prakash*

Research and Development, The NutraSweet Company, 601 E. Kensington Road, Mt. Prospect, Illinois 60056, USA

ABSTRACT

3,3-Dimethylbutyraldehyde is synthesized from the reaction of 1-chloro-3,3-dimethylbutane with DMSO in presence of a base and substoichiometric amounts of MX (M = Na, K; X = Br, I).

Neotame (*N*-[*N*-(3,3-dimethylbutyl)-*L*- α -aspartyl]-*L*-phenylalanine-*L*-methyl ester) is the next generation of non-nutritive sweetener from the NutraSweet Company. Neotame has a sweetness intensity about 8000 times that of sucrose and 30–40 times that of aspartame and is expected to have a wide range of applications in food categories.¹ A petition was submitted in December 1998 to the US FDA for the general use of neotame as a sweetener. Neotame is synthesized by reductive alkylation of aspartame with 3,3-dimethylbutyraldehyde (A) (Eq. 1).²

3395

Copyright © 2001 by Marcel Dekker, Inc.

www.dekker.com

^{*}Corresponding author.

ORDER		REPRINTS
-------	--	----------



In the process development of neotame, we have required large quantities of high quality **A**. Although **A** can be purchased from various vendors, it is advantageous from a supply standpoint to prepare **A** economically to reduce the overall cost of neotame. Therefore it was decided to explore economic synthetic routes to **A**.

It has been reported in the literature that **A** can be synthesized from the hydrolysis of 1,1-dichloro-3,3-dimethylbutane,³ oxidation of 3,3-dimethylbutanol,⁴ ozonolysis of 4,4-dimethyl-1-pentene,⁵ and reduction of 3,3-dimethylbutyl chloride.⁶ However, none of these methods were able to meet our cost goals.

The oxidation of a primary alkyl chloride or bromide with dimethyl sulfoxide (DMSO) to generate an aldehyde was described by Engel and co-workers.⁷ The reaction was facilitated by addition of 1.5 equivalent of sodium iodide. The reaction proceeds by formation of an alkyl iodide which is oxidized by DMSO.

This methodology seems applicable to the synthesis of **A** from 1-chloro-3,3-dimethylbutane, which can be readily prepared from the condensation of ethylene with *tert*-butyl chloride using $AlCl_3$ as a catalyst.⁸ However, the usage of 1.5 equivalent of iodide is still undesirable on an industrial scale due to its cost.

We describe herein a procedure for the synthesis of **A** from 1-chloro-3,3-dimethylbutane in the presence of a base and a substoichiometric amount of alkali bromide or iodide (Eq. 2).

1-Chloro-3,3-dimethylbutane itself does not react with DMSO at an appreciable rate without the addition of bromide or iodide. Alkali bromide



1-Chloro-3,3-dimethylbutane

M = Na, K; X = Br, I.

Marcel Dekker, Inc.

270 Madison Avenue, New York, New York 10016



3,3-DIMETHYLBUTYRALDEHYDE

or iodide facilitates the reaction by the formation of 1-bromo-3,3-dimethylbutane or 1-iodo-3,3-dimethylbutane, which is subsequently oxidized by DMSO to generate **A**. Since alkali bromide costs much less than alkali iodide, it would be economically advantageous to use bromide in Eq. 2. The reaction also works well with 1-bromo-3,3-dimethylbutane or 1-iodo-3,3-dimethylbutane.

In a typical experiment, a mixture of 1-chloro-3,3-dimethylbutane, DMSO, MX (M = Na, K; X = Br, I) and a base was loaded into a reactor and heated to 130–140°C with agitation for up to 14 h. The usage of alkali bromide and iodide is about 50% and 10–50% (w/w) respectively. An excess amount of a base, such as bicarbonate, carbonate, or zinc oxide, was used to neutralize HCl. Dimethyl sulfide was formed and distilled from the mixture during the reaction, which was terminated when > 95% of the 1-chloro-3,3-dimethylbutane reacted. The reaction mixture was subsequently distilled under house vacuum to yield the crude aldehyde product, which can be purified by further fractional distillation and used for most organic syntheses. The crude aldehyde may also be purified through formation and isolation of the solid A/sodium bisulfite adduct, Eq. 3. Regeneration from a base such as sodium bicarbonate in aqueous media generates A with a purity > 99%.



The final yield of **A** is 58% based on 1-chloro-3,3-dimethylbutane after purification through the aldehyde/bisulfite adduct and regeneration. Compound **A** produced from this process has been used successfully to produce high quality neotame.

In conclusion, we have developed a cost-effective synthetic route to 3,3-dimethylbutyraldehyde (A), a precursor to neotame. This process is robust and can be readily practiced on a commercial scale. We have also demonstrated that iodide may not be necessary, at least in our case, for the oxidation of primary alkyl chloride or bromide with DMSO to generate the corresponding aldehyde.

Marcel Dekker, Inc.

270 Madison Avenue, New York, New York 10016

ORDER		REPRINTS
-------	--	----------

GUO, SAWYER, AND PRAKASH

EXPERIMENTS

Preparation of 1-chloro-3,3-dimethylbutane. This compound was synthesized according to the literature.⁸ The catalyst used in the reaction, AlCl₃, may react with *tert*-butyl chloride at room temperature to generate an inactive material. It is desirable to add AlCl₃ at low temperature and to introduce ethylene as soon as possible after the catalyst dissolves.

Reaction of 1-chloro-3,3-dimethylbutane with DMSO/NaI. A 100 ml round bottom flask was loaded with 2.0 g 1-chloro-3,3-dimethylbutane, 2.0 sodium bicarbonate, 1.0 g NaI and 20 g DMSO- d_6 . The mixture was stirred and heated to 140°C for 2.5 h. ¹H NMR indicated that 62% 1-chloro-3,3-dimethylbutane was converted to 3,3-dimethylbutyraldehyde.

Reaction of 1-chloro-3,3-dimethylbutane with DMSO/KBr. A 100 ml round bottom flask was loaded with 3.1 g 1-chloro-3,3-dimethylbutane, 2.1 sodium bicarbonate, 1.5 g KBr and 18 g DMSO- d_6 . The mixture was stirred and heated to 140°C for 4.5 h. ¹HNMR indicated that the yield of 3,3-dimethylbutyraldehyde was 70%.

Reaction of 1-chloro-3,3-dimethylbutane with DMSO/NaBr. A 500 ml round bottom flask fitted with a distillation setup was loaded with 46.1 g 1-chloro-3,3-dimethylbutane, 23 g sodium bromide, 20 g zinc oxide and 276 g DMSO. The mixture was stirred and heated with an oil bath to 135–140°C under the nitrogen atmosphere for 12–14 h, during which some low boiling material (mostly dimethyl sulfide) was collected in a receiver.⁹ The reaction mixture was cooled to below 40°C and distilled under house vacuum to collect 31.6 g crude product (80% 3,3-dimethylbutyraldehyde) in a new receiver.

Purification of crude 3,3-dimethylbutyraldehyde. To a mixture of 30.6 g of the above described crude product (80% 3,3-dimethylbutyraldehyde) and 240 ml methyl tert-butyl ether (MTBE) was added drop wise 30 g sodium bisulfite in 60 ml water over a period of 20 min with vigorous agitation at $0-5^{\circ}$ C. A white solid precipitated upon addition of the bisulfite solution. The thus formed slurry was stirred for another 80 min with ice cooling followed by filtration. The solid was washed with 100 ml MTBE and $2 \times 100 \text{ ml} 60/40$ (vol/vol) MTBE/methanol and dried in a house vacuum oven $(40^{\circ}C)$ overnight. The weight of the dry solid is 48.0 g. 40.0 g of this dry solid, 22 g sodium bicarbonate and 200 ml water were loaded into a 250 ml round bottom flask. This mixture was heated to boiling and distilled to yield an azeotropic mixture of 3,3-dimethylbutyraldehyde and water (b.p. 82–85°C). Phase separation gave 18.0 g 3,3-dimethylbutyraldehyde.¹⁰ The overall yield is 58% based on 1-chloro-3,3-dimethylbutane. The purity of the aldehyde is >99% by GC. b.p. 106° C. ¹H NMR (DMSO- d_6): δ 1.01 (s, 9H), 2.24 (d, 2H), 9.75 (t, 1H).

3398



ORDER		REPRINTS
-------	--	----------

3,3-DIMETHYLBUTYRALDEHYDE

REFERENCES

- 1. Nofre, C.; Tinti, J.-M. US Patent 1996, 5480668.
- (a) Nofre, C.; Tinti, J.-M. US Patent 1996, 5510508; (b) Prakash, I. US Patent 1998, 5728862; (c) Prakash, I.; Bishay, I.; Schroeder, S. Synth. Commun. 1999, 29, 4461.
- 3. Schmerling, I. J. Amer. Chem. Soc. 1946, 68, 1650.
- 4. (a) Wiberg, K.B.; Squires, R.R. J. Amer. Chem. Soc. 1981, 103, 1650.
 (b) Cheung, C.K.; Wedinger, R.S.; le Noble, W.J. J. Org. Chem. 1989, 54, 570.
- 5. Whitmore, F.C.; Homeyer, A.H. J. Amer. Chem. Soc. 1933, 55, 4557.
- 6. Pospisek, J.; Blaha, K. Collect Czech. Chem. Comm. 1987, 52, 514.
- 7. Dave, P.; Byun, H.-S.; Engel, R. Synth. Comm. 1986, 16, 1343.
- 8. Branstrom, A. Acta Chem. Scand. 1959, 13, 611.
- 9. Alternatively 1-chloro-3,3-dimethylbutane was added drop-wise over a period of several hours into a preheated mixture of 1-chloro-3,3-dimethylbutane, sodium bromide, zinc oxide and DMSO at 130–140°C. This mixture continued to be heated at 130–140°C for a total of 12–14 h and was subsequently subjected to a similar work up procedure to produce 3,3-dimethylbutyraldehyde with similar yields.
- 10. 3,3-Dimethylbutyraldehyde may be oxidized by air into 3,3-dimethylbutyracid. Cautions should be taken to limit its contact with air during 3,3-dimethylbutyraldehyde's handling and storage.

Received in the USA January 21, 2001

Downloaded by [Stony Brook University] at 23:44 01 November 2014



3399

Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the <u>U.S. Copyright Office</u> for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on <u>Fair Use in the Classroom</u>.

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our <u>Website</u> <u>User Agreement</u> for more details.

Order now!

Reprints of this article can also be ordered at http://www.dekker.com/servlet/product/DOI/101081SCC100106050