

HETEROCYCLES, Vol. 64, 2004, pp. 61 - 64

Received, 13th September, 2004, Accepted, 2nd November, 2004, Published online, 9th November, 2004

**SYNTHESIS OF *N*-TRIFLUOROACETYL-2-HYDROXYNORAPORPHINE
BY RADICAL REACTION OF *p*-QUINOL ACETATE OF *N*-TRIFLUORO-
ACETYL-1-(2-IODOVERATRYL)METHYL-1, 2, 3, 4-TETRAHYDRO-7-
METHOXYISOQUINOLIN-6-OL[†]**


Akiko Moriya and Osamu Hoshino*,¹

Faculty of Pharmaceutical Sciences, Tokyo University of Science, 2641,

Yamazaki, Noda-shi, Chiba 278-8510, Japan

Abstract — Radical reaction of *p*-quinol acetate (**8b**) of *N*-trifluoroacetyl-1-(2-iodoveratryl)methyl-1, 2, 3, 4-tetrahydro-7-methoxyisoquinolin-6-ol (**7b**) using tributyltin hydride gave the corresponding noraporphine (**9**)(45 %), while that of the bromo congener (**8a**) failed. Similar reaction of **8b** using tris(trimethylsilyl)silane improved the yield (75 %) of **9**.

Previously,² we reported that lead tetraacetate (LTA) oxidation of *N*-trifluoroacetyl-1-veratrylmethyl-1, 2, 3, 4-tetrahydro-7-methoxyisoquinolin-6-ol (**1**) in CH₂Cl₂ gives rise to the corresponding *o*-quinol acetate (*o*-




Scheme 1

[†] Dedicated to Dr. Pierre Potier on the occasion of his 70th birthday.

QA)(2) in good yield, reaction of which with AcOH at 30-40°C produces *p*-QA (3)(Scheme 1).

Although treatment³ of *p*-QAs (5) of *N*-acetyltetrahydroisoquinolin-7-ols (4) with CF₃CO₂H in CH₂Cl₂




Scheme 2

afforded the corresponding 1-hydroxynoraporphines (6)(Scheme 2), attempts⁴ to synthesize 2-hydroxynoraporphine by reaction of 2 or 3 under a variety of acidic conditions were unfruitful, in which the reaction gave only undetermined complex mixtures along with reversed 1.


Therefore, in order to synthesize 2-hydroxynoraporphine from *p*-QA (8) and to examine its reactivity, radical reaction of 1-(2-bromo- and iodoveratryl)methyl *p*-QAs (8)⁵ was performed. This paper deals with successful formation of *N*-trifluoroacetyl-2-hydroxynoraporphine (9) from 1-(2-iodoveratryl)methyl *p*-QA (8b).

LTA oxidation of *N*-trifluoroacetyl-1-(2-bromo- and iodoveratryl)methyltetrahydro-7-methoxyisoquinolin-6-ols (7)(7a,^{6, 7} mp 157-158°C; 7b,^{6, 9} mp 190-191°C) in AcOH at 30-40°C for 5 h furnished the corresponding 1-(2-haloveratryl)methyl *p*-QAs (8)(8a,⁶ mp 162-163°C; 8b,⁶ mp 175-176°C) in good yield (Scheme 3).




Scheme 3

Unexpectedly, reaction of 8a in boiling benzene (1 mmol %) containing AIBN (0.55 eq.) and tributyltin hydride (1.5 eq.) for 4 h and purification on preparative TLC (SiO₂; developing solvent: benzene : EtOAc = 1 : 1) gave reversed 7a (54 %) and untact 8a (13 %), respectively (Scheme 4).



Scheme 4


However, similar reaction¹⁰ of **8b** followed by purification gave desired *N*-trifluoroacetyl-2-hydroxy-noraporphine (**9**) in 45 % yield, in which starting material (**8b**) and reversed **7b** were not determined (Scheme 5). Furthermore, an analogous reaction of **8b** in boiling benzene (1 mmol %) using (TMS)₃SiH (1.5 eq.) in place of Bu₃SnH afforded **9** in 70 % yield (Scheme 5).



Scheme 5

The difference in reaction would be attributable to the reactivity¹¹ of aryl bromide and iodide in radical reaction. Also this assumption is supported by formation of reversed **7a**, which would be produced by reductoin of a carbonyl group in **8a** with Bu₃SnH followed by elimination of AcOH.

A pathway on formation of **9** is considered to proceed *via* an intermediate (**10**)(Scheme 6), although a pathway *via* direct cyclization of **7b**, which might be formed from **8b** by reaction similar to that described for **8a**, could not be excluded completely. Because reaction of **7b** with AIBN (0.5 eq.) and Bu₃SnH (1.5 eq.) in boiling benzene (1 mmol %) for 7 h gave **9** (11 %) together with **1** (24 %).



Scheme 6

In conclusion, synthesis of noraporphine (**9**) by radical reaction of 1-(2-iodoveratryl)methyl *p*-QA (**8b**) was accomplished successfully. Thus, the reaction promises the preparation of noraporphine bearing no substituent(s) in D ring, which has been not yet synthesized by acid treatment of *p*-QA.

ACKNOWLEDGEMENTS

The authors are grateful to Mrs. F. Hasegawa of this faculty for MS spectral measurements.

REFERENCES AND NOTES

1. Present address: 683-56, Sashi-ougi, Nishi-ku, Saitama-shi, Saitama 331-0047, Japan.
2. O. Hoshino, M. Suzuki, and H. Ogasawara, *Tetrahedron*, 2001, **57**, 265.
3. O. Hoshino, H. Ogasawara, M. Suzuki, and B. Umezawa, *Heterocycles*, **1987**, 25, 151; O. Hoshino, H. Ogasawara, M. Suzuki, M. Arasawa, and B. Umezawa, *Heterocycles*, **1990**, 30, 385; O. Hoshino, H. Ogasawara, M. Arasawa, M. Suzuki, and K. Iizima, *Heterocycles*, **1993**, 35, 1005.
4. F. Mutoh, H. Ogasawara, and O. Hoshino, unpublished data.
5. Since *p*-QA was more stable than *o*-QA, the former was used in the reaction.
6. All new compounds described in this text gave satisfactory ^1H - and ^{13}C -NMR, and HRMS spectral data.
7. Starting material (**7a**) was prepared by *N*-trifluoroacetylation (trifluoroacetic anhydride, K_2CO_3 , CH_2Cl_2 ; K_2CO_3 , MeOH) of 1-(2-bromoveratryl)methyltetrahydroisoquinolin-6-ol⁸ from benzyloxyphenethylamine and 2-bromohomoveratric acid in a usual manner (Bischler-Napieralski reaction, NaBH_4 reduction, and debenzylation with conc. HCl).
8. M. S. Premila and B. R. Pai, *Indian J. Chem.*, **1975**, 13, 13.
9. Starting material (**7b**) was prepared from benzyloxyphenethylamine and 2-iodohomoveratric acid in a manner similar to that noted for **7a**.
10. A solution of **8b** (50 mg, 0.08 mmol) in benzene (8 mL) containing AIBN (6.7 mg, 0.04 mmol) and Bu_3SnH (35.8 mg, 0.12 mmol) was refluxed for 7 h. A residue remained on removal of the solvent *in vacuo* was dissolved in MeCN. The MeCN solution was washed with hexane. Usual work-up of the MeCN layer gave an oily product, which was purified on preparative TLC (SiO_2 ; developing solvent: benzene : EtOAc = 1 : 1) to produce **9** (15.5 mg, 45 %, mp 177-178°C); ^1H -NMR (270 MHz)(CDCl_3) δ : 3.61 (3H, s), 3.93 (6H, s), 5.90, 6.72, 6.80, and 7.99 (each 1H, s); ^{13}C -NMR (62.5 MHz)(CDCl_3) δ : 30.1, 32.8, 41.4, 52.6, 55.8, 56.1, 60.2, 110.8, 111.5, 113.2, 123.3, 123.4, 126.8, 128.9, 129.7, 142.8, 148.0, 148.6, 148.8, and 156.3; EI MS m/z (M^+): 423; HRMS m/z calcd for $\text{C}_{21}\text{H}_{20}\text{NO}_5\text{F}_3$ (M^+) 423.1293, found: 423.1302.
11. M. Poutsma, in 'Free Radicals', Vol. 2, ed. by J. K. Kochi, Wiley, N. Y., 1973, p.113; W. C. Danen, in 'Methods in Free Radical Chemistry', Vol. 5, ed. E. L. S. Huyser, Dekker, N. Y., 1974, p. 1.