

## Reactions of $\text{Me}_3\text{N}$ with $\text{GeH}_3\text{Br}$ : Evidence for Stabilized Germanium Dihydride Species

CHARLES A. HEIL and ARLAN D. NORMAN\*

Department of Chemistry and Biochemistry, University of Colorado, Boulder, Col. 80309, U.S.A.

(Received January 15, 1988)

### Abstract

Bromogermane ( $\text{GeH}_3\text{Br}$ ) and  $\text{Me}_3\text{N}$  (1:1 mole ratio) reaction at  $-78^\circ\text{C}$  yields  $\text{Me}_3\text{N}\cdot\text{GeH}_3\text{Br}$  (1). 1 with  $\text{H}_2\text{O}$  or  $\text{MeOH}$  yields  $(\text{GeH}_3)_2\text{O}$  or  $\text{MeOGeH}_3$ , respectively. 1 with a one molar excess of  $\text{Me}_3\text{N}$  forms  $(\text{Me}_3\text{N})_2\text{GeH}_3\text{Br}$  (2). Thermal decomposition of 2 produces mainly  $\text{Ge}_3\text{H}_8$  (25–30%) and  $\text{Me}_3\text{N}\cdot\text{NHBr}$ .  $\text{GeH}_3\text{Br}$  with excess  $\text{Me}_3\text{N}$  in  $\text{Et}_2\text{O}$  at  $0^\circ\text{C}$  yields  $\text{Me}_3\text{NHBr}$  and a species in solution formulated as  $\text{Me}_3\text{N}\cdot\text{GeH}_2$  (3). Decomposition of 3 in the presence of  $\text{Me}_3\text{GeH}$  or  $\text{Me}_2\text{GeH}_2$  results in small quantities of  $\text{Me}_3\text{Ge}_2\text{H}_3$  or  $\text{Me}_2\text{Ge}_2\text{H}_4$ , respectively. Above  $-63^\circ\text{C}$ , solid 1 undergoes intramolecular rearrangement to a product characterized as  $\text{Me}_3\text{N}\cdot\text{NHBr}\cdot 1/x(\text{GeH}_2)_x$  (4), a material which contains stabilized  $\text{GeH}_2$  units. The characterization of 1–4 based on their spectral properties and reactivity towards  $\text{B}_2\text{H}_6$  and/or  $\text{HBr}$ , and their potential in further germane syntheses, are discussed.

### Introduction

The existence of germynes ( $\text{GeX}_2$ ;  $\text{X}$  = halogen, aryl, and alkyl) stabilized by coordination to bases [1] or to metals and bases [2] is well established. Germanium dihydride ( $\text{GeH}_2$ )<sub>x</sub>, in the condensed phase has been reported as an intermediate or product in  $\text{Ge}_2\text{H}_6$  [3] or  $\text{NaGeH}_3\text{--PhBr}$  [4] in liquid  $\text{NH}_3$ ,  $\text{R}_3\text{N--GeH}_3\text{Cl}$  [5],  $\text{LiX}$  catalyzed decomposition of  $\text{Ge}_2\text{H}_6$  [6],  $\text{KGeH}_3\text{BH}_3$  [7] thermolysis, or  $\text{GeH}_3\text{Cl--NH}_3$  [8] reactions [7–12]; however, reports of its properties vary. No clear evidence for monomeric  $\text{GeH}_2$  other than in the gaseous phase [9–13] has appeared. Since  $\text{GeH}_2$ ,  $(\text{GeH}_2)_x$ , or its complexes could be important in the synthesis of new germane polymers or germanium subhydrides (e.g.  $\text{GeH}_{<1}$ ) for use as electronic materials, we have undertaken studies of its synthesis in the condensed phase. Now, in the  $\text{GeH}_3\text{Br--Me}_3\text{N}$  reaction system, we have found evidence for novel stabilized  $(\text{GeH}_2)_x$  species. Our studies are reported below.

\* Author to whom correspondence should be addressed.

### Experimental

#### Apparatus and Materials

All manipulations were carried out in standard vacuum lines or under a  $\text{N}_2$  atmosphere [14].  $^1\text{H}$  NMR spectra were obtained with a Varian A-60A spectrometer.  $\delta$  values were measured relative to  $\text{Me}_4\text{Si}$ ;  $+\delta$  values are downfield from the standard. Infrared and mass spectra were obtained using Beckman Model 4250 and Varian MAT CH-5 spectrometers, respectively.

Bromogermane [15],  $\text{Me}_2\text{GeH}_2$  [16, 17],  $\text{Me}_3\text{GeH}$  [16, 18],  $\text{B}_2\text{H}_6$  [19] and  $\text{DCI}$  [20] were prepared and purified as described previously.  $\text{Me}_3\text{GeBr}$  (Alfa Inorganics) was purified by fractional condensation.  $\text{Me}_2\text{O}$  and diglyme were distilled from  $\text{LiAlH}_4$ . Acetonitrile (Baker) and  $\text{Me}_3\text{N}$  (Matheson Gas) were distilled from  $\text{P}_4\text{O}_{10}$ . Deuterium oxide (>99.8% D, Mallinckrodt) and  $\text{MeOH}$  (Fischer, reagent) were used as obtained.

Reaction products from reactions below were identified by comparison of either their IR, NMR or mass spectra with published spectra.

#### Reactions of $\text{Me}_3\text{N}$ with $\text{GeH}_3\text{Br}$

##### $\text{Me}_3\text{N}\cdot\text{GeH}_3\text{Br}$ (1)

$\text{Me}_3\text{N}$  (0.117–0.135 mmol) was condensed onto  $\text{GeH}_3\text{Br}$  (0.213 mmol) at  $-196^\circ\text{C}$ . The mixture was warmed to  $-78^\circ\text{C}$ , mixed, and then warmed to  $-63^\circ\text{C}$ . A white solid formed immediately. Materials volatile at  $-63^\circ\text{C}$  were removed and measured. Typically, the  $\text{Me}_3\text{N}:\text{GeH}_3\text{Br}$  reactant ratio ranged from 1.09–0.97:1.00. In reactions involving  $\text{Me}_3\text{N}:\text{GeH}_3\text{Br}$  reactant ratios >1, small quantities (2–8% of total Ge) of  $\text{GeH}_4$  [21],  $\text{Ge}_2\text{H}_6$  [22, 23] and  $\text{Ge}_3\text{H}_8$  [23, 24] formed also. Under analogous conditions, but in  $\text{Me}_2\text{O}$  solvent, identical reactant combination was observed but with less  $\text{GeH}_4$ ,  $\text{Ge}_2\text{H}_6$ , or  $\text{Ge}_3\text{H}_8$  evolution.

##### $(\text{Me}_3\text{N})_2\text{GeH}_3\text{Br}$ (2)

$\text{Me}_3\text{N}$  (0.5 mmol) was condensed onto 1 (typically, 0.100–0.145 mmol) and allowed to stand at  $-78^\circ\text{C}$ . After 5–14 h, excess  $\text{Me}_3\text{N}$  was removed and

measured. The 1:Me<sub>3</sub>N consumed ranged from 1.0–1.05:1.00, consistent with formation of (Me<sub>3</sub>N)<sub>2</sub>-GeH<sub>3</sub>Br (2).

B<sub>2</sub>H<sub>6</sub> at –63 °C did not react with 2. A 0.05 mmol sample of 2 reacted with excess HBr at –63 °C to form GeH<sub>3</sub>Br (0.023 mmol) and traces of Ge<sub>2</sub>H<sub>6</sub> and Ge<sub>3</sub>H<sub>8</sub>. Thermolysis of 2 at 25 °C to 80 °C produces Me<sub>3</sub>NHBr, high yields of Ge<sub>3</sub>H<sub>8</sub> (25–30%) along with H<sub>2</sub> and traces of GeH<sub>4</sub> and Ge<sub>2</sub>H<sub>6</sub>.

#### Excess Me<sub>3</sub>N: Me<sub>3</sub>NGeH<sub>2</sub> (3)

Typically, Me<sub>3</sub>N (3–20 mmol) was allowed to react at temperatures of –78 to 0 °C with a GeH<sub>3</sub>Br (0.10–0.41 mmol)–diglyme (3.0–4.0 ml) solution. The solution turned bright yellow and a white precipitate formed. The solution (3) was filtered at 0 °C. The solid was extracted with H<sub>2</sub>O and found to contain typically 92–95% of the expected Me<sub>3</sub>NHBr. Solution 3, transferred at 0 °C to an NMR tube, showed only resonances due to Me<sub>3</sub>N.

Excess HBr (3–5 mmol) was condensed onto filtrate 3 and the solution was warmed to 25 °C. The yellow solution immediately turned colorless. Unreacted HBr, GeH<sub>3</sub>Br (typically 12–22% of initial Ge) and MeBr (0.2–0.3 mmol) were separated from the diglyme by fractional condensation. In control experiments, HBr reacted with diglyme to yield comparable amounts of MeBr under analogous reaction time and temperature conditions. No MeBr formed in a 1 h reaction of HBr with Me<sub>3</sub>NHBr.

Solutions of 3 were prepared as above, with the addition that a 10–30 fold excess of Me<sub>2</sub>GeH<sub>2</sub> or Me<sub>3</sub>GeH was dissolved in the initial diglyme solution. The stirred solutions were warmed to 0 °C (0.3–2 h) and 25 °C (0.5–17 h). The volatile products included GeH<sub>4</sub> (2–10%), Ge<sub>3</sub>H<sub>8</sub> (14–40%), traces of Ge<sub>4</sub>H<sub>10</sub> and Me<sub>2</sub>Ge<sub>2</sub>H<sub>4</sub> (with Me<sub>2</sub>GeH<sub>2</sub> coreactant) and Me<sub>3</sub>Ge<sub>2</sub>H<sub>3</sub> (with Me<sub>3</sub>GeH coreactant).

GeH<sub>3</sub>Br–excess Me<sub>3</sub>N reactions were carried out as above, except that Me<sub>2</sub>O replaced diglyme as solvent. The reaction solution was filtered at –78 °C and the yellow solution was evaporated to dryness *in vacuo*. The yellow solid obtained was unreactive towards H<sub>2</sub>O and yielded only traces of GeH<sub>3</sub>Cl and Ge<sub>3</sub>H<sub>8</sub> upon treatment with excess HCl at 25 °C.

Samples of neat GeH<sub>3</sub>Br (0.12 mmol) and Me<sub>3</sub>N (*ca.* 8 mmol) were allowed to warm to –100 °C in the probe of the <sup>1</sup>H NMR spectrometer. Between –80 and –40 °C, weak resonances at δ3.65 ppm (area 1) and δ1.27 ppm (area 3) relative to internal Me<sub>3</sub>N were observed. Above –40 °C these disappeared and only the Me<sub>3</sub>N and Me<sub>3</sub>NH<sup>+</sup> resonances remained.

#### Gaseous phase reactions

Equimolar quantities of GeH<sub>3</sub>Br and Me<sub>3</sub>N (at 50 to 250 torr) were placed in separate bulbs of a double-bulb gas mixing apparatus at 23 to 90 °C.

Opening a stopcock between bulbs allowed the reactants to mix. A volatile white solid formed immediately. Upon standing, the solid became non-volatile, turned yellow, and slowly evolved GeH<sub>4</sub>, Ge<sub>3</sub>H<sub>8</sub>, and traces of Ge<sub>2</sub>H<sub>6</sub> and Ge<sub>4</sub>H<sub>10</sub>.

#### Reactions of Me<sub>3</sub>N·GeH<sub>3</sub>Br (1)

##### With HBr and/or B<sub>2</sub>H<sub>6</sub>

Samples of Me<sub>3</sub>N·GeH<sub>3</sub>Br (1) (0.18–0.35 mmol), prepared at –78 °C, were allowed to react with excess HBr or B<sub>2</sub>H<sub>6</sub> as shown in Table I (experiments 1 and 2). After reaction, products were removed and analyzed. Reactions with HBr and B<sub>2</sub>H<sub>6</sub> yielded Me<sub>3</sub>NHBr and Me<sub>3</sub>NBH<sub>3</sub> [25], respectively, GeH<sub>3</sub>Br and traces of germanes.

TABLE I. Reaction of 1:1 Me<sub>3</sub>N–GeH<sub>3</sub>Br Complex with HBr and/or B<sub>2</sub>H<sub>6</sub>

Experiment	Complex history <sup>a</sup>		Acid	GeH <sub>3</sub> Br product yield; % of Ge in initial Me <sub>3</sub> NGeH <sub>3</sub> Br
	temperature (°C)	time (h)		
1	–78	1.0	HBr	97.8
2	–78	1.0	B <sub>2</sub> H <sub>6</sub>	92.5
3	–63	1.0	HBr	98.6
4	–63	1.0	B <sub>2</sub> H <sub>6</sub>	79.5
5	–23	0.5	B <sub>2</sub> H <sub>6</sub>	58
			HBr	23
6	23	4.0	B <sub>2</sub> H <sub>6</sub>	0
			HBr	<1

<sup>a</sup>Me<sub>3</sub>N·GeH<sub>3</sub>Br prepared at –78 °C, then warmed to the temperature for the time shown.

#### Thermolysis

Samples of 1 were allowed to warm to –63, –23 or 23 °C for times as shown in Table I (exps. 3–6). Samples were then treated at –78 °C with HBr or B<sub>2</sub>H<sub>6</sub> followed by HBr. As above, Me<sub>3</sub>NBH<sub>3</sub> along with GeH<sub>3</sub>Br and traces of germanes formed. The percent GeH<sub>3</sub>Br produced in each case is given in the Table. Intractable nonvolatile residues left after reaction in experiments 3–6 were not analyzed further.

1 (0.11 mmol, prepared at –78 °C) was warmed to –23 °C during 4 h in the presence of Me<sub>3</sub>GeH (3.5 mmol). Traces of GeH<sub>4</sub>, Ge<sub>2</sub>H<sub>6</sub>, and Ge<sub>3</sub>H<sub>8</sub> and unreacted Me<sub>3</sub>GeH were obtained.

#### With H<sub>2</sub>O or MeOH

Water (0.051 mmol) was condensed onto 1 (0.135 mmol, prepared at –78 °C) and the mixture was allowed to warm to –63 °C. After 3 h, H<sub>2</sub>O was consumed and (GeH<sub>3</sub>)<sub>2</sub>O [26] (0.051 mmol) and Me<sub>3</sub>NHBr were formed exclusively. From a 1–D<sub>2</sub>O reaction, (GeH<sub>3</sub>)<sub>2</sub>O and Me<sub>3</sub>NDBr were obtained quantitatively. The (GeH<sub>3</sub>)<sub>2</sub>O showed no Ge–D IR spectral absorption.

MeOH (0.07 mmol) and **1** (0.108 mmol) were allowed to react at  $-63^{\circ}\text{C}$ . After 6 h,  $\text{MeOGeH}_3$  [27] (0.064 mmol) was recovered.

#### Synthesis of $\text{Me}_3\text{NHBr} \cdot 1/x(\text{GeH}_2)_x$ (**4**)

Solid **1**, prepared in  $\text{Me}_2\text{O}$  at  $-78^{\circ}\text{C}$ , was warmed to  $25^{\circ}\text{C}$  *in vacuo* during 15 min in the dark. Traces of  $\text{GeH}_4$ ,  $\text{Ge}_2\text{H}_6$ ,  $\text{Ge}_3\text{H}_8$  and  $\text{GeH}_3\text{Br}$  ( $<5\%$  of total germanium) evolved. The resulting solid **4** was pure white and non-volatile.

$^1\text{H}$  NMR spectra of **4** could not be obtained owing to the insolubility of **4** in solvents in which it did not decompose, e.g.  $\text{CCl}_4$ ,  $\text{C}_6\text{H}_6$ , ethers and hexane.

Freshly prepared samples of **4** were powdered and pressed into KBr disks in a darkened room under  $\text{N}_2$  for IR spectral analysis. Preparation time varied from 15–30 min. Initial IR spectra were obtained in  $<0.5$  h, and subsequent spectra were obtained at 48 h intervals over a 10 day period. Samples were kept in the dark between analyses. Typical spectra are shown in Fig. 1.

**4** showed only slight discoloration after 7 days *in vacuo* in the dark at  $25^{\circ}\text{C}$ ; however, after 1 h in sunlight it turned deep yellow. After 6 days exposure to ambient laboratory fluorescent light, intense yellow discoloration along with evolution of traces of germanes occurred. At  $250$ – $300^{\circ}\text{C}$  rapid conversion of the white solid to yellow, red, brown and finally metallic grey (metallic Ge) occurred along with evolution of  $\text{H}_2$ ,  $\text{GeH}_4$  and higher germanes.  $\text{Me}_3\text{NHBr}$  (93–95%) sublimed from the pyrolysis zone.

**4** did not react with  $\text{B}_2\text{H}_6$  at  $25^{\circ}\text{C}$  in 2 h or with  $\text{HBr}$  at  $25^{\circ}\text{C}$  in 0.3 h. During 17 h at  $25^{\circ}\text{C}$ ,  $\text{DCl}$

(three-fold excess) reacted with **4** to produce  $\text{GeH}_3\text{Cl}$ , along with lesser amounts of  $\text{GeH}_2\text{DCl}$  (22–24% yield, overall).

**4** prepared under  $\text{N}_2$  was subjected to X-ray powder pattern analysis. After 3 h in  $\text{Cu K}\alpha$  irradiation, **4** turned deep yellow. The powder pattern was superimposable with that of known  $\text{Me}_3\text{NHBr}$ .

#### Dissolution of $\text{Me}_3\text{NHBr} \cdot 1/x(\text{GeH}_2)_x$ (**4**) in Acetonitrile

Freshly prepared **4** (0.10–0.15 mmol) was extracted with acetonitrile (5 ml) in the dark at  $25^{\circ}\text{C}$  during 3 h. Partial dissolution of the solid occurs. The  $^1\text{H}$  NMR spectrum of the acetonitrile solution showed resonances due to the  $\text{Me}_3\text{NH}^+$  ion and acetonitrile. A grey waxy solid (**5**) remained undissolved. Evaporation of the acetonitrile solution to dryness *in vacuo* yielded  $\text{Me}_3\text{NHBr}$  (92.5–94.0% of theoretical). **5** decomposes slowly at  $25^{\circ}\text{C}$ , although it yellows slowly upon exposure to light. Thermolysis above  $150^{\circ}\text{C}$  causes rapid evolution of  $\text{H}_2$ ,  $\text{GeH}_4$ ,  $\text{Ge}_2\text{H}_6$  and  $\text{Ge}_3\text{H}_8$ .

Elemental composition of **5** was determined by pyrolysis of **5** at  $400^{\circ}\text{C}$ , followed by Toepler analysis [14] of  $\text{H}_2$  and determination of Ge by peroxide titration of the resulting Ge metal [5, 6]. *Anal.* Calc. for  $\text{GeH}_2$ : Ge, 97.30; H, 2.70. Found: Ge, 95.8–96.4; H, 2.45–2.85%.

#### Reaction of $\text{Me}_3\text{GeBr}$ with $\text{Me}_3\text{N}$

Typically,  $\text{Me}_3\text{N}$  (0.120 mmol) and  $\text{Me}_3\text{GeBr}$  (0.090 mmol) were allowed to mix at  $25^{\circ}\text{C}$  and cooled to  $-78^{\circ}\text{C}$ . Excess  $\text{Me}_3\text{N}$  (0.031 mmol) was removed. Adduct  $\text{Me}_3\text{N} \cdot \text{Me}_3\text{GeBr}$  (**7**) remained. During a 2 h period at  $-63^{\circ}\text{C}$ , around 20% of the adduct in an equimolar mixture of  $\text{Me}_3\text{N}$  and  $\text{Me}_3\text{GeBr}$  (determined by IR spectra) was removed *in vacuo*. Reaction of **7** with excess  $\text{B}_2\text{H}_6$  at  $-78^{\circ}\text{C}$  resulted in consumption of a half molar equivalent of  $\text{B}_2\text{H}_6$ , and quantitative recovery, within experimental limits, of the  $\text{Me}_3\text{GeBr}$  and  $\text{Me}_3\text{NBH}_3$ .

## Results and Discussion

Bromogermane ( $\text{GeH}_3\text{Br}$ ) reacts with  $\text{Me}_3\text{N}$  to form the products summarized in Scheme 1. At  $-78^{\circ}\text{C}$ ,  $\text{Me}_3\text{N}$  with excess  $\text{GeH}_3\text{Br}$  reacts neat or in  $\text{Me}_2\text{O}$  solvent to form a 1:1  $\text{Me}_3\text{N} \cdot \text{GeH}_3\text{Br}$  complex (**1**) in a process analogous to that which occurs between  $\text{GeH}_3\text{Cl}$  and  $\text{Me}_3\text{N}$  reported in brief form earlier [5, 26]. Even in the absence of solvent,  $\text{GeH}_3\text{Br} \cdot \text{Me}_3\text{N}$  reactions are remarkably clean and are accompanied by formation of only traces of  $\text{GeH}_4$ ,  $\text{Ge}_2\text{H}_6$  and  $\text{Ge}_3\text{H}_8$ . Adduct **1** is a colorless solid, which appears unchanged after several hours at  $-78^{\circ}\text{C}$ ; however, upon warming it undergoes conversion to material(s) with different physical and chemical properties.







Fig. 1. Infrared spectrum of: (A), freshly prepared **4**; (B), **4** after 10 days at  $25^{\circ}\text{C}$ ; (C), pure  $\text{Me}_3\text{NHBr}$ .



Scheme 1.




Three formulations for the  $\text{Me}_3\text{N} \cdot \text{GeH}_3\text{Br}$  adduct can be considered: **I**, a five-coordinate structure in which  $\text{Me}_3\text{N}$  is coordinatively bonded to the Ge atom of  $\text{GeH}_3\text{Br}$ ; **II**, a salt in which a  $\text{GeH}_3^+$  moiety has been transferred to the  $\text{Me}_3\text{N}$  to form a quaternary salt; and **III**, a bromogermanide salt in which  $\text{Me}_3\text{N}$  has abstracted a  $\text{H}^+$  from  $\text{GeH}_3\text{Br}$  to form the  $\text{GeH}_2\text{Br}^-$  ion.



Precedents for species analogous to **I–III** exist. Five-coordinated silicon and germanium halides, where coordination involves amine, phosphine or ether bases are well known [9, 11, 28]. Type **II** complexes, e.g.  $[\text{Me}_3\text{NSiH}_3^+]\text{Br}^-$  and  $[\text{Me}_3\text{N}(\text{Me}_3\text{Ge})^+]\text{Br}^-$ , analogs of tetraalkylammonium ions, from  $\text{Me}_3\text{N-SiH}_3\text{Br}$  and  $\text{Me}_3\text{N-Me}_3\text{GeBr}$  reactions, respectively, have been tentatively reported by Srivastava and Onyszchuk [29]. A type **III** species would be analogous to the  $\text{GeH}_3^-$  anion formed upon deprotonation of  $\text{GeH}_4$  [30a] or the  $\text{SiCl}_3^-$  ion formed in reaction of  $\text{Me}_3\text{N}$  with  $\text{HSiCl}_3$  [30b]. Although  $\text{GeH}_4$  requires bases stronger than  $\text{Me}_3\text{N}$  for deprotonation [30, 31] it is possible that  $\text{GeH}_3\text{Br}$ , because of the

electron withdrawing effect of the Br atom, might be more acidic.

Characterization of **1** by spectral data (IR and  $^1\text{H}$  NMR) was unsuccessful because of the low thermal stability and solubility of **1** in solvents with which it did not react. However, from its chemical reactivity properties, definitive structural inferences can be made. **1** reacts quantitatively with  $\text{H}_2\text{O}$  (or  $\text{D}_2\text{O}$ ) or  $\text{HBr}$  and in high (ca. 90%) yield with  $\text{B}_2\text{H}_6$  or  $\text{MeOH}$  as shown in eqns. (2)–(5)



Substitution of  $\text{D}_2\text{O}$  for  $\text{H}_2\text{O}$  yields  $\text{Me}_3\text{NDBr}$  and  $(\text{GeH}_3)_2\text{O}$ ; within experimental error no incorporation of deuterium into the germyl ether occurs. Reactions in eqns. (4) and (5) are in fact good preparative syntheses for  $(\text{GeH}_3)_2\text{O}$  and  $\text{MeOGeH}_3$  [32], species which otherwise have been difficult to obtain in reasonable yields.

Of the structures possible for  $\text{Me}_3\text{N} \cdot \text{GeH}_3\text{Br}$  (**I–III**), **I** seems most likely. Since **1** reacts quantitatively with  $\text{D}_2\text{O}$  to form  $(\text{GeH}_3)_2\text{O}$  and no  $(\text{GeD}_3)_2\text{O}$ , the complex must contain an intact  $\text{GeH}_3$  unit; hence formulation **III** is ruled out. Since  $\text{GeH}_3\text{Br}$  is easily displaced from the complex by  $\text{B}_2\text{H}_6$ , we conclude

the  $\text{GeH}_3$  unit is not bonded to nitrogen in a quaternary cationic species as in II.

Upon warming slightly or standing at  $-63^\circ\text{C}$ , the character of 1 changes dramatically and its reactivity toward  $\text{H}_2\text{O}$  and  $\text{B}_2\text{H}_6$  decreases. After 1 h at  $-63^\circ\text{C}$  (exp. 3), the  $\text{Me}_3\text{N}\cdot\text{GeH}_3\text{Br}$  complex with  $\text{B}_2\text{H}_6$  produces only 79.5% of the theoretically expected  $\text{GeH}_3\text{Br}$ , although it still is in such a form that  $\text{GeH}_3\text{Br}$  can be displaced quantitatively (exp. 4) by  $\text{HBr}$ . Later, after 3 h at  $-23^\circ\text{C}$ , the  $\text{GeH}_3\text{Br}$  that can be displaced by  $\text{B}_2\text{H}_6$  has dropped further to 58%. Again, considerable reactivity towards  $\text{HBr}$  is retained. After reaction with  $\text{B}_2\text{H}_6$ ,  $\text{HBr}$  displaces another 23% of the  $\text{GeH}_3\text{Br}$  from the product. This behavior suggests involvement of a transient (intermediate) species whose formulation is different from 1 and that of the subsequently formed product 4. Such an intermediate could be  $[\text{Me}_3\text{NGeH}_3^+]\text{Br}^-$  (5; Scheme 1), a species which contains a  $\text{GeH}_3$  unit bonded to a quaternized nitrogen center as a result of cleavage of the  $\text{Ge}-\text{Br}$  bond in 1. Again, a species such as  $[\text{Me}_3\text{NH}^+][\text{GeH}_2\text{Br}]$  (type III, above) appears ruled out because in reactions with  $\text{D}_2\text{O}$  the digermyl ether obtained shows no significant  $\text{Ge}-\text{D}$  IR absorptions.

Further warming of the 1:1  $\text{Me}_2\text{N}\cdot\text{GeH}_3\text{Br}$  product to ambient room temperature ( $23^\circ\text{C}$ ) yields a pure white, light sensitive solid (4) which is essentially unreactive towards  $\text{B}_2\text{H}_6$ ,  $\text{HBr}$  (exp. 6, Table I) and  $\text{H}_2\text{O}$ . 4 is stable at ambient temperatures in the absence of light for periods of days, showing only barely decomposition and traces of germane evolution.

Solid 4 dissolves partially and slowly (3 h) in acetonitrile or water at room temperature. A colorless solution, which upon evaporation was found to contain 92–95% of the expected  $\text{Me}_3\text{NHBBr}$ , and a grey unstable gum (6) are formed. Compound 6 after drying *in vacuo*, analyzed closely for  $(\text{GeH}_2)_x$ . Upon heating, 6 discolors to yellow, orange and finally brown. During this decomposition  $\text{GeH}_4$ ,  $\text{Ge}_2\text{H}_6$  and  $\text{Ge}_3\text{H}_8$  evolve. Final heating at  $>300^\circ\text{C}$  yields a black residue which was presumed to be metallic germanium. It should be noted that dissolution of 4 is markedly different from that of  $\text{Me}_3\text{NHBBr}$ .

The IR spectra of 4, as a function of time, provide some insight into its composition. Spectral data are shown in Fig. 1. The spectrum of pure  $\text{Me}_3\text{NHBBr}$  (Fig. 1C) is shown for comparison purposes. The initial spectrum of a freshly prepared sample (Fig. 1A) which had been exposed minimally to light prior to spectral analysis showed absorptions clearly attributable to the  $\text{Me}_3\text{NH}^+$  ion. In addition, strong absorptions at 2045 (a), 755 (c), and 610 (d)  $\text{cm}^{-1}$ , and a weak absorption at 860 (b)  $\text{cm}^{-1}$  is present. Absorption a occurs in the characteristic  $\text{Ge}-\text{H}$  stretching mode region. Figure 1B shows the spectrum after 10 days at  $25^\circ\text{C}$ . During this period, the

sample was intentionally exposed to ambient laboratory light. A noticeable decrease in the intensity of absorptions a, b and c is evident along with some absorption line broadening. These are assigned to germanium hydride species. Although our data do not allow unambiguous characterization of 4, or the material to which it decomposes thermally or photochemically, they do show that 4 contains  $\text{Me}_3\text{NH}^+$  ions from the outset, and in a concentration which seems invariant with time.


It is known that 4 forms from 1, with no change in overall compound composition (excepting loss of traces of germanes due to decomposition), that 4 contains  $\text{Me}_3\text{NH}^+$  ions from the outset, and that 4 yields  $\text{Me}_3\text{NHBBr}$  upon dissolution in acetonitrile. Thus, a reasonable formulation for 4 is  $\text{Me}_3\text{NH}^+\text{Br}\cdot 1/x(\text{GeH}_2)_x$ , a composition containing  $\text{Me}_3\text{NH}^+$  and  $\text{Br}^-$  ions and  $(\text{GeH}_2)$  units. The germylene unit is written as  $(\text{GeH}_2)_x$ , since  $\text{GeH}_2$  monomers are likely not the main species present. Mono  $\text{GeH}_2$  species can be present only to a minor degree, since only 20–25% reaction with  $\text{HBr}$  to  $\text{GeH}_3\text{Br}$  occurs. It is assumed that  $(\text{GeH}_2)_x$  is less reactive than the mono  $\text{GeH}_2$  and therefore forms no  $\text{GeH}_3\text{Br}$  upon treatment with  $\text{HBr}$ . Also, 4 with  $\text{Me}_3\text{GeH}$  gave no  $\text{Me}_3\text{Ge}_2\text{H}_3$ , suggesting that the  $\text{GeH}_2$  species present is not very reactive towards insertion into  $\text{Ge}-\text{H}$  bonds.

The  $\text{GeH}_2$  species obtained in 4 seems different from that reported previously [3–5, 12]. The  $(\text{GeH}_2)_x$  from  $\text{Ge}_2\text{H}_6$  disproportionation or  $\text{GeH}_3^--\text{RBr}$  reactions reportedly decomposes quickly above  $-63$  or  $25^\circ\text{C}$ , respectively, to germanes and  $(\text{GeH}_{<2})$  polymer. The  $(\text{GeH}_2)_x$  in 4 does not discolor or evolve germane in significant quantities for days. This suggests that the  $(\text{GeH}_2)_x$  moieties are stabilized in the solid either by the  $\text{Me}_3\text{NH}^+$  ion or the  $\text{Br}^-$  ion, or both. It is known that bases stabilize  $\text{GeX}_2$  species in general [1, 2, 10], and there are cases where  $\text{GeX}_2$  units are stabilized by coordination both to a base and an acid, e.g. to  $\text{Fe}(\text{CO})_4$  and THF, in  $\text{THF}-\text{GeX}_2-\text{Fe}(\text{CO})_4$  [2]. In 4, weak coordination of  $(\text{GeH}_2)_x$  species by  $\text{Br}^-$  (acting as a base) and/or by  $\text{Me}_3\text{NH}^+$  (acting as an acid) could occur. In this way 4 represents a unique system in which  $(\text{GeH}_2)_x$  becomes stabilized and displays behavior unique from that of other reported species.

Compound 6, the product left after acetonitrile extraction of  $\text{Me}_3\text{NHBBr}$  from 4, analyzes for  $\text{GeH}_2$ . It also is more stable than previously reported  $(\text{GeH}_2)_x$ . It is unreactive towards  $\text{HBr}$  and shows no photodecomposition or significant thermal decomposition over moderately long periods. Either it contains entrapped  $\text{Me}_3\text{NHBBr}$  in concentrations below what we could detect by our analytical techniques, which stabilizes it, or it contains  $\text{GeH}_2$  units bonded differently from other known systems. Timms has suggested that solid  $(\text{GeH}_2)_x$  might


contain Ge—Ge bonds [12]. However, another arrangement involving Ge—H—Ge bonding analogous to that in  $\text{GeX}_2$  [10] is possible. **6** might be of the former type, and thus is expected to be unreactive towards HBr. Further studies of compound **6** and its properties are warranted.

Bromogermane with excess  $\text{Me}_3\text{N}$  yields products which are more complex. Compound **1** with an additional equivalent of  $\text{Me}_3\text{N}$  reacts to form a 2:1 complex as




The structure of the 2:1 complex is unclear. Since it is relatively unreactive towards  $\text{B}_2\text{H}_6$ , it likely is not a simple six-coordinate complex of  $\text{GeH}_3\text{Br}$ . A more complex formulation such as  $[\text{Me}_3\text{NH}^+][\text{Me}_3\text{NGeH}_2\text{Br}^-]$ , in which the second  $\text{Me}_3\text{N}$  deprotonates the acidic  $\text{Me}_3\text{NGeH}_3\text{Br}$  complex is possible. However, complete characterization awaits further study.

In the presence of  $\text{Me}_3\text{N}$  in large excess,  $\text{Me}_3\text{N}$  and  $\text{GeH}_3\text{Br}$  yield  $\text{Me}_3\text{NHBBr}$  and a yellow solution. Based on reaction stoichiometry data and the fact that  $\text{Me}_3\text{NHBBr}$  is recovered quantitatively, reaction appears to occur as



a reaction similar to what has been noted previously in the formation of the ylid  $\text{Me}_3\text{NCH}_2$  from  $\text{Me}_3\text{N}$  and  $\text{CH}_3\text{I}$  [33]. The species in solution, formulated as  $\text{Me}_3\text{NGeH}_2$  (**3**), decomposes rapidly but for a time retains reactivity towards Ge—H bonds. Reaction of **3** with  $\text{Me}_2\text{GeH}_2$  or  $\text{Me}_3\text{GeH}$  results in low yield formation of  $\text{Me}_2\text{Ge}_2\text{H}_4$  and  $\text{Me}_3\text{Ge}_2\text{H}_3$ , respectively, in addition to substantial quantities (13–37%) of  $\text{Ge}_3\text{H}_8$ . Similar behavior has been noted previously for the ylid  $\text{Me}_3\text{NCH}_2$  [33].

Reaction of  $\text{Me}_3\text{N}$  with  $\text{Me}_3\text{GeBr}$  results in formation of a 1:1 complex  $\text{Me}_3\text{N} \cdot \text{Me}_3\text{GeBr}$  (**7**) as



analogous to that which occurs between  $\text{Me}_3\text{N}$  and  $\text{GeH}_3\text{Br}$  and reported earlier for  $\text{Me}_3\text{GeBr}$  with amines [5, 28]. Reaction of **7** with  $\text{B}_2\text{H}_6$  yields  $\text{Me}_3\text{NBH}_3$  quantitatively; complex **7** is partially dissociated even at  $-63^\circ\text{C}$  to the extent that  $\text{Me}_3\text{GeBr}$  and  $\text{Me}_3\text{N}$  can be removed slowly *in vacuo*. However, beyond this, the similarity between **7** and **1** ends. Upon warming, **7** dissociates without formation of other products. In excess  $\text{Me}_3\text{N}$ , it shows no tendency to form products analogous to those in eqn. (7). Clearly, because the  $\text{GeH}_3\text{Br}$  has transferrable H atoms, it is unique in the products it can form.

## Acknowledgements

Support of this work by National Science Foundation grants GP-8090 and CHE-8312856 and the

Colorado Advanced Materials Institute is gratefully acknowledged.

## References

- (a) R. B. King, *Inorg. Chem.*, **2**, 199 (1963); (b) P. Jutzi, H.-J. Hoffman, D. J. Brauer and C. Kruger, *Angew. Chem., Int. Ed. Engl.*, **12**, 1002 (1973); (c) W.-W. duMont and H. Schuman, *J. Organomet. Chem.*, **85**, C45 (1975); (d) P. Jutzi, H.-J. Hoffman and K.-H. Wyes, *J. Organomet. Chem.*, **83**, 341 (1974); (e) W.-W. duMont, B. Neudert, G. Rudolph and H. Schuman, *Angew. Chem., Int. Ed. Engl.*, **15**, 308 (1976); (f) C. D. Silverman and M. Zeldin, *Inorg. Chem.*, **19**, 272 (1980).
- (a) T. J. Marks and A. R. Newman, *J. Am. Chem. Soc.*, **95**, 769 (1973); (b) P. Jutzi and W. Steiner, *Angew. Chem., Int. Ed. Engl.*, **15**, 684 (1976); (c) J. D. Cotton, P. J. Davidson and M. F. Lappert, *J. Chem. Soc., Dalton Trans.*, 2275 (1976).
- R. M. Dreyfuss and W. L. Jolly, *Inorg. Chem.*, **7**, 2645 (1968).
- S. N. Glarum and C. A. Kraus, *J. Am. Chem. Soc.*, **72**, 5398 (1950).
- T. N. Srivastava and M. Onyszczuk, *137th Nat. Meeting Amer. Chem. Soc.*, Cleveland, 1960, Abstracts.
- W. R. Bornhorst and M. A. Ring, *Inorg. Chem.*, **7**, 1009 (1968).
- D. S. Rustad and W. L. Jolly, *Inorg. Chem.*, **7**, 213 (1968).
- M. L. Dennis and R. W. Work, *J. Am. Chem. Soc.*, **55**, 4486 (1933).
- F. Glockling, 'The Chemistry of Germanium', Academic Press, New York, 1969.
- J. Satgé, M. Massol and P. Riviere, *J. Organomet. Chem.*, **56**, 1 (1973).
- E. Wiberg and E. Amberger, 'Hydrides of the Elements of Main Groups I - IV', Elsevier, Amsterdam, 1971.
- P. L. Timms, *Prep. Inorg. Reactions*, **4**, 59 (1968).
- R. L. Jenkins, R. A. Kedrowski, L. E. Elliot, D. C. Toppans, D. J. Schleyer and M. A. Ring, *J. Organomet. Chem.*, **86**, 347 (1975).
- D. F. Shriver and M. A. Drezdson, 'The Manipulation of Air-Sensitive Compounds', 2nd. edn., McGraw-Hill, New York, 1986.
- J. W. Anderson, G. K. Barker, J. E. Drake and R. T. Hemmings, *Can. J. Chem.*, **50**, 1607 (1972).
- A. E. Finholt, A. C. Bond, K. E. Wilzbach and H. I. Schlesinger, *J. Am. Chem. Soc.*, **69**, 2692 (1947).
- D. F. Van deVondel and G. P. Van derKelen, *Bull. Soc. Chim. Belg.*, **74**, 467 (1965).
- R. J. Mathias, J. Satgé and F. Mathis, *Spectrochim. Acta*, **18**, 1463 (1962).
- A. D. Norman and W. L. Jolly, *Inorg. Synth.*, **11**, 15 (1968).
- H. C. Brown and C. Groot, *J. Am. Chem. Soc.*, **64**, 2223 (1942).
- (a) J. W. Straley, C. H. Tindal and H. H. Nielson, *Phys. Rev.*, **62**, 161 (1942); (b) V. A. Crawford, K. H. Rhea and M. K. Wilson, *J. Chem. Phys.*, **37**, 2377 (1962).
- D. A. Davis and R. M. Hexter, *J. Chem. Phys.*, **24**, 1029 (1962).
- W. L. Jolly and J. E. Drake, *Inorg. Synth.*, **7**, 34 (1963).
- J. Drake and W. L. Jolly, *J. Chem. Soc.*, 2807 (1962).
- B. Rice, R. J. Galiano and W. J. Lehmann, *J. Phys. Chem.*, **61**, 1222 (1957).
- T. D. Goldfarb and S. Sujishi, *J. Am. Chem. Soc.*, **86**, 1679 (1964).
- G. A. Gibbon, J. T. Wang and C. H. VanDyke, *Inorg. Chem.*, **6**, 1990 (1967).

- 28 E. A. V. Ebsworth, 'Volatile Silicon Compounds', Macmillan, New York, 1963.
- 29 T. N. Srivastava and M. Onyszchuk, *Angew. Chem.*, 75, 578 (1963).
- 30 (a) T. Birchall and I. Drommond, *J. Chem. Soc. A*, 1859 (1970); (b) M. A. Ring, R. L. Jenkins, R. Zangreb and H. C. Brown, *J. Am. Chem. Soc.*, 93, 265 (1971).
- 31 D. S. Rustad and W. L. Jolly, *Inorg. Chem.*, 6, 1986 (1967).
- 32 C. A. Heil and A. D. Norman, *Inorg. Chim. Acta*, 64, 1197 (1982).
- 33 (a) W. K. Musker, *J. Chem. Educ.*, 45, 200 (1968), (b) W. K. Musker and R. R. Stevens, *Inorg. Chem.*, 8, 255 (1969).