

Medicinal Flowers

Part 29¹⁾

Acylated Oleanane-Type Triterpene Bisdesmosides: Perennisaponins G, H, I, J, K, L, and M with Pancreatic Lipase Inhibitory Activity from the Flowers of *Bellis perennis*

by Toshio Morikawa^{a)}, Xuezheng Li^{b)}, Eriko Nishida^{a)}, Seikou Nakamura^{b)}, Kiyofumi Ninomiya^{a)}, Hisashi Matsuda^{b)}, Yoshimi Oda^{b)}, Osamu Muraoka^{a)}, and Masayuki Yoshikawa^{a,b)}

^{a)} Pharmaceutical Research and Technology Institute, Kinki University, 3-4-1 Kowakae, Higashi-osaka, Osaka 577-8502, Japan

^{b)} Kyoto Pharmaceutical University, Misasagi, Yamashina-ku, Kyoto 607-8412, Japan
(phone: +81-75-5954633; fax: +81-75-5954768; e-mail: myoshika@mb.kyoto-phu.ac.jp)

The MeOH extract from the flowers of *Bellis perennis* was found to show pancreatic-lipase inhibitory activity (IC_{50} 455 μ g/ml). From the extract, seven new triterpene saponins named perennisaponins G (**1**; IC_{50} 163 μ M), H (**2**; 137 μ M), I (**3**; 147 μ M), J (**4**; 148 μ M), K (**5**; 223 μ M), L (**6**; 81.4 μ M), and M (**7**; 195 μ M) were isolated as pancreatic lipase inhibitors. The stereostructures of **1–7** were elucidated on the basis of chemical and spectroscopic evidence.

Introduction. – The Asteraceae plant *Bellis perennis* is distributed widely in Europe and North Africa, and the whole flowering plant has been used for bruises, bleeding, muscular pain, purulent skin diseases, and rheumatism in European folk medicine [2][3]. This herbal medicine is also well-known as an ornamental plant, and its flowers and young leaves are edible as a salad. During the course of our studies on bioactive constituents from medicinal flowers [1–20], we found that the MeOH extract from the flowers of *B. perennis* was found to inhibit the increase of plasma triglyceride (TG) levels in olive-oil-loaded mice [2]. From the MeOH extract, 13 acylated saponin constituents, perennisosides I–VII [2] and perennisaponins A–F [3], were isolated, together with eight saponins and two glycosides [2][3].

Our continuing search now led to the isolation and characterization of seven new acylated oleanane-type triterpene oligoglycosides named perennisaponins G–M (**1–7**), which were obtained from the flowers of *B. perennis*. Furthermore, we examined the inhibitory effects against pancreatic lipase of perennisaponins **1–7**.

Results and Discussion. – The flowers of *B. perennis* cultivated in Albania were treated with MeOH to give an extract (25.8% from the dried flowers), as described previously [2]. The MeOH extract, which was found to inhibit pancreatic lipase activity (IC_{50} 455 mg/ml), was partitioned between AcOEt and H₂O (1:1) to furnish an

¹⁾ For Part 28, see [1].

AcOEt-soluble fraction (6.7%) and an aqueous phase. The aqueous phase was subjected to column chromatography (*Diaion HP-20*, $\text{H}_2\text{O} \rightarrow \text{MeOH}$) to give H_2O - and MeOH-eluted fractions (12.5 and 6.4%), respectively [2]. The MeOH-eluted fraction was subjected to normal- and reversed-phase column chromatographies and finally HPLC to give **1** (0.124%), **2** (0.139%), **3** (0.0038%), **4** (0.060%), **5** (0.027%), **6** (0.0028%), and **7** (0.065%) (*Fig. 1*).




Fig. 1. Perennisaponins G–M (1–7) isolated from Bellis perennis

Compound **1** was obtained as an amorphous powder with a negative optical rotation. The IR spectrum of **1** showed absorption bands at 1736 and 1638 cm^{-1} ascribable to ester C=O and olefin moieties, and broad bands at 3450 and 1049 cm^{-1} , suggestive of an oligoglycoside structure. In the positive- and negative-ion-mode FAB-MS of **1**, quasi-molecular-ion peaks were observed at m/z 1327 ($[M + \text{Na}]^+$) and 1303 ($[M - \text{H}]^-$), respectively, and a positive-mode HR-FAB-MS analysis revealed the molecular formula of **1** to be $\text{C}_{63}\text{H}_{100}\text{O}_{28}$. On alkaline hydrolysis of **1** with 10% aqueous KOH in $\text{H}_2\text{O}/1,4\text{-dioxane}$ 1 : 1, polygalactic acid 3-(α -L-rhamnopyranoside) (**1a**) [21] was obtained together with AcOH, which was identified by HPLC analysis of the

corresponding 4-nitrobenzyl derivative [3–7][18]. Treatment of **1** with 0.5% MeONa/MeOH provided a deacyl derivative, bellissaponin BS1 (= (2 β ,3 β ,4 α ,16 α)-3-[(6-deoxy- α -L-mannopyranosyl)oxy]-2,16,23-trihydroxyolean-12-en-28-oic acid *O*-6-deoxy- α -L-mannopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-6-deoxy- α -L-mannopyranosyl-(1 $\rightarrow$ 2)-6-deoxy- β -D-galactopyranosyl ester; **1b**) [3]. The ^1H - and ^{13}C -NMR data of **1** (Tables 1 and 2), which were assigned by various NMR experiments (including DEPT, DQF, HMQC, HMBC, and TOCSY) showed signals assignable to six Me s at $\delta(\text{H})$ 0.93 (Me(29)), 1.03 (Me(30)), 1.17 (Me(26)), 1.23 (Me(24)), 1.58 (Me(25)), and 1.75 (Me(27)), an oxygenated CH_2 group at $\delta(\text{H})$ 3.72 (2 br. s, $\text{CH}_2(23)$), three oxygenated CH groups at $\delta(\text{H})$ 4.39 (br. s, H-C(3)), 4.72–4.76 (*m*, H-C(2)), and 5.16 (br. s, H-C(16)), an olefinic H-atom at $\delta(\text{H})$ 5.64 (*t*-like, $J \approx 3$ Hz, H-C(12)), a fucopyranosyl moiety at $\delta(\text{H})$ 1.24 (*d*, $J = 6.4$ Hz, Me(6'')) and 6.02 (*d*, $J = 8.3$ Hz, H-C(1'')), three rhamnopyranosyl moieties at $\delta(\text{H})$ 1.60 (*d*, $J = 6.0$ Hz, Me(6')), 1.63 (*d*, $J = 6.0$ Hz, Me(6''')), 1.75 (*d*, $J = 6.2$ Hz, Me(6''')), 5.76 (br. s, H-C(1')), 6.07 (br. s, H-C(1'')), and 6.21 (br. s, H-C(1''')), and a xylopyranosyl moiety at $\delta(\text{H})$ 5.14 (*d*, $J = 7.6$ Hz, H-C(1''')), together with two AcO groups at $\delta(\text{H})$ 1.98 and 1.99 (2 s, 2 Me). Comparison of the ^{13}C -NMR data of **1** with those of the deacetyl derivative **1b** revealed acetylation shifts around the 4''-position of the β -D-fucopyranosyl moiety (**1**: $\delta(\text{C})$ 73.9 (C(3'')), 74.4 (C(4'')), and 70.4 (C(5'')); **1b**: $\delta(\text{C})$ 76.5 (C(3'')), 73.1 (C(4'')), and 72.5 (C(5''))) and around the 2'''-position of the inner α -L-rhamnopyranosyl moiety of the 28-*O*-sugars (**1**: $\delta(\text{C})$ 98.8 (C(1''')), 73.3 (C(2''')), and 70.0 (C(3''')); **1b**: $\delta(\text{C})$ 101.4 (C(1''')), 71.9 (C(2''')), and 72.6 (C(3'''))). The positions of the two AcO groups in **1** were clarified on the basis of an HMBC experiment, which showed $^1\text{H} \rightarrow ^{13}\text{C}$ long-range correlations between the following pairs as shown in Fig. 2: $\delta(\text{H})$ 1.98 (Me of Ac) and 5.51 (br. s, H-C(4'') of Fuc)/ $\delta(\text{C})$ 171.1 (C=O of Ac), and $\delta(\text{H})$ 1.99 (Me of Ac) and 5.99–6.01 (*m*, H-C(2''') of the inner Rha)/ $\delta(\text{C})$ 170.4 (C=O of Ac). On the basis




Fig. 2. Selected HMBCs (H $\rightarrow$ C) of **1**

of the above evidence, the structure of perennisaponin G (**1**) was determined as 3-*O*- α -L-rhamnopyranosylpolygalacic acid 28-[*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-2-*O*-acetyl- α -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-4-*O*-acetyl- β -D-fucopyranosyl] ester.

Table 1. Selected ^1H -NMR Data (600 MHz, 40°, (D₅)pyridine) of **1–3**. δ in ppm, J in Hz.

	1	2	3
H–C(2)	4.72–4.76 (<i>m</i>)	4.70–4.74 (<i>m</i>)	4.72–4.76 (<i>m</i>)
H–C(3)	4.39 (br. <i>s</i>)	4.38 (<i>d</i> , $J = 2.8$)	4.40 (<i>d</i> , $J = 3.1$)
H–C(12)	5.64 (<i>t</i> -like, $J \approx 3$)	5.63 (<i>t</i> -like, $J \approx 3$)	5.65 (<i>t</i> -like, $J \approx 3$)
H–C(16)	5.16 (br. <i>s</i>)	5.16 (br. <i>s</i>)	5.16 (br. <i>s</i>)
H–C(18)	3.39 (<i>dd</i> -like)	3.41 (<i>dd</i> , $J = 3.4, 14.4$)	3.40 (<i>dd</i> -like)
CH ₂ (23)	3.72 (br. <i>s</i>)	3.70 (br. <i>s</i>)	3.71 (br. <i>s</i>)
Me(24)	1.23 (<i>s</i>)	1.26 (<i>s</i>)	1.24 (<i>s</i>)
Me(25)	1.58 (<i>s</i>)	1.55 (<i>s</i>)	1.59 (<i>s</i>)
Me(26)	1.17 (<i>s</i>)	1.14 (<i>s</i>)	1.19 (<i>s</i>)
Me(27)	1.75 (<i>s</i>)	1.75 (<i>s</i>)	1.76 (<i>s</i>)
Me(29)	0.93 (<i>s</i>)	0.92 (<i>s</i>)	0.93 (<i>s</i>)
Me(30)	1.03 (<i>s</i>)	1.03 (<i>s</i>)	1.03 (<i>s</i>)
<i>3-O-Rha</i> :			
H–C(1')	5.76 (br. <i>s</i>)	5.76 (br. <i>s</i>)	5.78 (br. <i>s</i>)
Me(6')	1.60 (<i>d</i> , $J = 6.0$)	1.60 (<i>d</i> , $J = 6.2$)	1.62 (<i>d</i> , $J = 6.1$)
<i>28-O-Sugars</i> :			
<i>Fuc</i> :			
H–C(1'')	6.02 (<i>d</i> , $J = 8.3$)	6.06 (<i>d</i> , $J = 6.9$)	6.03 (<i>d</i> , $J = 8.3$)
H–C(4'')	5.51 (br. <i>s</i>)	5.47 (br. <i>s</i>)	5.59 (br. <i>d</i> , $J \approx 4$)
Me(6'')	1.24 (<i>d</i> , $J = 6.4$)	1.24 (<i>d</i> , $J = 6.2$)	1.32 (<i>d</i> , $J = 6.4$)
AcO–C(4'')	1.98 (<i>s</i>)	1.99 (<i>s</i>)	
<i>3HBO–C(4'')</i> :			
CH ₂ (2)			2.68 (<i>dd</i> , $J = 5.2, 14.4$), 2.78 (<i>dd</i> , $J = 7.6, 14.4$) 4.46–4.48 (<i>m</i>) 1.39 (<i>d</i> , $J = 6.4$)
H–C(3)			
Me(4)			
<i>Inner Rha</i> :			
H–C(1''')	6.07 (br. <i>s</i>)	6.07 (br. <i>s</i>)	6.09 (<i>d</i> , $J = 1.5$)
H–C(2''')	5.99–6.01 (<i>m</i>)	5.00–5.02 (<i>m</i>)	5.99 (<i>dd</i> , $J = 1.5, 3.4$)
H–C(3''')	4.79 (<i>dd</i> , $J = 3.4, 9.8$)	5.87 (<i>dd</i> , $J = 2.7, 9.6$)	4.78 (<i>dd</i> , $J = 3.6, 9.8$)
Me(6''')	1.75 (<i>d</i> , $J = 6.2$)	1.75 (<i>d</i> , $J = 6.2$)	1.75 (<i>d</i> , $J = 6.1$)
AcO–C(2''')	1.99 (<i>s</i>)		1.96 (<i>s</i>)
AcO–C(3''')		2.04 (<i>s</i>)	
<i>Xyl</i> :			
H–C(1''')	5.14 (<i>d</i> , $J = 7.6$)	4.98 (<i>d</i> , $J = 7.6$)	5.15 (<i>d</i> , $J = 7.7$)
<i>Terminal Rha</i> :			
H–C(1''')	6.21 (br. <i>s</i>)	6.21 (br. <i>s</i>)	6.17 (br. <i>s</i>)
Me(6''')	1.63 (<i>d</i> , $J = 6.0$)	1.66 (<i>d</i> , $J = 6.2$)	1.64 (<i>d</i> , $J = 6.1$)

Compound **2** was also obtained as an amorphous powder with a negative optical rotation. The IR spectrum of **2** showed absorption bands at 3450, 1734, 1638, and 1049 cm^{−1}, ascribable to OH, ester C=O, olefin, and ether functions. The molecular formula of **2**, C₆₃H₁₀₀O₂₈, was determined from the positive- and negative-ion-mode

Table 2. ^{13}C -NMR Data of **1–3**. At 150 MHz, 40°, in (D_5)pyridine; δ in ppm.

	1	2	3		1	2	3
C(1)	44.8	44.8	44.8	28-O-Sugars:			
C(2)	71.1	71.1	71.1	Fuc:			
C(3)	81.4	81.4	81.5	C(1'')	94.3	94.2	94.3
C(4)	43.0	43.0	43.0	C(2'')	75.1	76.8	75.2
C(5)	47.6	47.7	47.6	C(3'')	73.9	74.1	73.5
C(6)	18.3	18.3	18.3	C(4'')	74.4	74.4	74.4
C(7)	33.2	33.2	33.3	C(5'')	70.4	70.3	70.5
C(8)	40.2	40.2	40.3	C(6'')	16.4	16.4	16.5
C(9)	47.5	47.5	47.5	AcO–C(4'')	171.1, 20.7	171.1, 20.7	
C(10)	37.2	37.1	37.2	3HBO–C(4'')			
C(11)	24.1	24.1	24.1	C(1)			172.3
C(12)	122.8	122.8	122.8	C(2)			45.0
C(13)	144.2	144.2	144.2	C(3)			64.8
C(14)	42.3	42.2	42.3	C(4)			24.0
C(15)	36.2	36.1	36.2	Inner Rha:			
C(16)	74.1	74.1	74.1	C(1''')	98.8	102.7	98.7
C(17)	49.7	49.7	49.5	C(2''')	73.3	69.4	73.2
C(18)	41.6	41.6	41.6	C(3''')	70.0	75.5	70.1
C(19)	47.3	47.3	47.3	C(4''')	83.8	77.9	83.8
C(20)	30.7	30.8	30.8	C(5''')	68.5	68.9	68.5
C(21)	35.9	35.9	36.0	C(6''')	18.6	19.2	18.5
C(22)	32.0	31.9	32.0	AcO–C(2''')	170.4, 20.9		170.4, 20.9
C(23)	65.3	65.2	65.3	AcO–C(3''')		170.9, 21.4	
C(24)	14.9	15.0	15.0	Xyl:			
C(25)	17.7	17.7	17.8	C(1''')	106.8	105.8	106.8
C(26)	17.7	17.6	17.7	C(2''')	76.2	75.0	76.2
C(27)	27.1	27.1	27.1	C(3''')	83.3	83.3	83.4
C(28)	176.1	176.2	176.2	C(4''')	69.2	69.8	69.3
C(29)	33.1	33.1	33.1	C(5''')	67.3	67.1	67.3
C(30)	24.5	24.6	24.6	Terminal Rha:			
3-O-Rha:				C(1''')	102.6	102.6	102.6
C(1')	104.2	104.1	104.2	C(2''')	72.5	72.5	72.5
C(2')	72.4	72.4	72.5	C(3''')	72.6	72.7	72.6
C(3')	72.7	72.7	72.8	C(4''')	73.9	73.9	74.0
C(4')	73.2	73.5	74.0	C(5''')	69.9	69.8	70.0
C(5')	70.3	70.3	70.3	C(6''')	18.5	18.6	18.6
C(6')	18.6	18.6	18.6				

FAB-MS (m/z 1327 ($[M + \text{Na}]^+$) and 1303 ($[M - \text{H}]^-$), resp.) and by positive-ion-mode HR-FAB-MS, which was the same as that of **1**. Alkaline hydrolysis of **2** with 10% aqueous KOH in $\text{H}_2\text{O}/1,4$ -dioxane 1:1 liberated **1a** together with AcOH. Treatment of **2** with 0.5% MeONa/MeOH yielded **1b**. The ^1H - and ^{13}C -NMR spectra of **2** (Tables 1 and 2) indicated the presence of the following functions: a polygalacic acid part (six Me s at $\delta(\text{H})$ 0.92 (Me(29)), 1.03 (Me(30)), 1.14 (Me(26)), 1.26 (Me(24)), 1.55 (Me(25)), and 1.75 (Me(27))), an oxygenated CH_2 group at $\delta(\text{H})$ 3.70 (br. s, $\text{CH}_2(23)$), three oxygenated CH groups at $\delta(\text{H})$ 4.38 (*d*, $J = 2.8$ Hz, $\text{H}-\text{C}(3)$), 4.70–4.74 (*m*, $\text{H}-\text{C}(2)$), and 5.16 (br. s, $\text{H}-\text{C}(16)$), and an olefinic H-atom at $\delta(\text{H})$ 5.63 (*t*-like, $J \approx 3$ Hz,

H–C(12))), a fucopyranosyl moiety at $\delta(\text{H})$ 1.24 (*d*, $J = 6.2$ Hz, Me(6'')) and 6.06 (*d*, $J = 6.9$ Hz, H–C(1'')), three rhamnopyranosyl moieties at $\delta(\text{H})$ 1.60 (*d*, $J = 6.2$ Hz, Me(6')), 1.66 (*d*, $J = 6.2$ Hz, Me(6''')), 1.75 (*d*, $J = 6.2$ Hz, Me(6'')), 5.76 (br. *s*, H–C(1')), 6.07 (br. *s*, H–C(1'')), and 6.21 (br. *s*, H–C(1''')), and a xylopyranosyl moiety at $\delta(\text{H})$ 4.98 (*d*, $J = 7.6$ Hz, H–C(1''')), together with two AcO groups at $\delta(\text{H})$ 1.99 and 2.04 (2*s*, 2 Me). The ^1H - and ^{13}C -NMR data of **2** were superimposable on those of **1**, except for the signals due to the part around the inner Rha unit of the 28-*O*-sugars. The positions of the two AcO groups in **2** were determined by an HMBC experiment, which showed the following $^1\text{H} \rightarrow ^{13}\text{C}$ long-range correlations: $\delta(\text{H})$ 1.99 (*s*, Me of Ac) and 5.47 (br. *s*, H–C(4'')) of Fuc/ $\delta(\text{C})$ 171.1 (C=O of Ac), and $\delta(\text{H})$ 2.04 (*s*, Me of Ac) and 5.87 (*dd*, $J = 2.7, 9.6$ Hz, H–C(3''')) of the inner Rha/ $\delta(\text{C})$ 170.9 (C=O of Ac). Consequently, the structure of **2** was elucidated as 3-*O*- α -L-rhamnopyranosylpolygalactic acid 28-[*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-3-*O*-acetyl- α -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-4-*O*-acetyl- β -D-fucopyranosyl] ester.

Compound **3** was also obtained as an optically active amorphous powder. The positive- and negative-ion-mode FAB-MS of **3** showed quasi-molecular-ion peaks at m/z 1371 ($[M + \text{Na}]^+$) and 1347 ($[M - \text{H}]^-$), respectively. The positive-ion-mode HR-FAB-MS of **3** revealed the molecular formula to be $\text{C}_{65}\text{H}_{104}\text{O}_{29}$. Alkaline hydrolysis of **3** with 10% aqueous KOH in $\text{H}_2\text{O}/1,4$ -dioxane 1:1 liberated **1a** and two organic acids, AcOH and 3-hydroxybutanoic acid (3HBOH), which were identified by HPLC analysis of the corresponding 4-nitrobenzyl derivatives. In addition, treatment of **3** with 0.5% MeONa/MeOH gave **1b**, together with methyl (+)-(3*S*)-3-hydroxybutanoate [22], which was identified by HPLC and an optical-rotation detector [23–27]. The H- and C-atom signals in the ^1H - and ^{13}C -NMR spectra of **3** (Tables 1 and 2, resp.) were superimposable on those of **1**, except for the signals due to the acyl groups: an AcO group at $\delta(\text{H})$ 1.96 (*s*, Me of Ac) and a (3*S*)-3-hydroxybutanoyl (3HB) group at $\delta(\text{H})$ 1.39 (*d*, $J = 6.4$ Hz, Me(4)), 2.68 (*dd*, $J = 5.2, 14.4$ Hz, 1 H–C(2)), 2.78 (*dd*, $J = 7.6, 14.4$ Hz, 1 H–C(2)), and 4.46–4.48 (*m*, H–C(3)). In the HMBC experiment with **3**, the following $^1\text{H} \rightarrow ^{13}\text{C}$ long-range correlations were observed: $\text{CH}_2(2)$ of 3HB, H–C(3) of 3HB, and $\delta(\text{H})$ 5.59 (br. *d*, $J \approx 4$ Hz, H–C(4'')) of Fuc/ $\delta(\text{C})$ 172.3 (C=O of 3HB); and $\delta(\text{H})$ 1.96 (*s*, Me of Ac) and 5.99 (*dd*, $J = 1.5, 3.4$ Hz, H–C(2''')) of the inner Rha/ $\delta(\text{C})$ 170.4 (C=O of Ac). Thus, the connectivities of the acyloxy groups in **3** were elucidated. On the basis of above-mentioned evidence, the structure of **3** was determined as 3-*O*- α -L-rhamnopyranosylpolygalactic acid 28-[*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-2-*O*-acetyl- α -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-4-*O*-(3*S*)-3-hydroxybutanoyl- β -D-fucopyranosyl] ester.

Compounds **4** and **5** were obtained as amorphous powders with negative optical rotations. The molecular formula of both **4** and **5**, $\text{C}_{67}\text{H}_{106}\text{O}_{30}$, was determined from the positive- and negative-ion-mode FAB-MS at m/z 1413 ($[M + \text{Na}]^+$) and 1389 ($[M - \text{H}]^-$) and by positive-ion-mode HR-FAB-MS measurement. Alkaline hydrolysis of **4** and **5** with 10% aqueous KOH in $\text{H}_2\text{O}/1,4$ -dioxane 1:1 provided **1a** and two organic acids, AcOH and 3HBOH, respectively. Treatment of each **4** and **5** with 0.5% MeONa/MeOH gave **1b**, together with methyl (+)-(3*S*)-3-hydroxybutanoate, respectively. The H- and C-atom signals in the ^1H - and ^{13}C -NMR spectra of **4** (Tables 3 and 4) indicated the presence of an aglycon part, containing six Me *s* at $\delta(\text{H})$ 0.95 (Me(29)), 1.06 (Me(30)), 1.18 (Me(26)), 1.22 (Me(24)), 1.58 (Me(25)), and 1.75 (Me(27)), a CH_2 and

three CH groups bearing an O-atom function at $\delta(\text{H})$ 3.69 (br. s, $\text{CH}_2(23)$), 4.35 (*d*, $J = 3.5$ Hz, $\text{H}-\text{C}(3)$), 4.68–4.72 (*m*, $\text{H}-\text{C}(2)$), and 5.12 (br. s, $\text{H}-\text{C}(16)$), an olefinic H-atom at $\delta(\text{H})$ 5.65 (*t*-like, $J \approx 3$ Hz, $\text{H}-\text{C}(12)$), and a fucopyranosyl moiety at $\delta(\text{H})$ 1.25 (*d*, $J = 6.3$ Hz, $\text{Me}(6'')$) and 5.99 (*d*, $J = 8.0$ Hz, $\text{H}-\text{C}(1'')$), three rhamnopyranosyl moieties at $\delta(\text{H})$ 1.60 (*d*, $J = 6.1$ Hz, $\text{Me}(6')$), 1.67 (*d*, $J = 6.3$ Hz, $\text{Me}(6''')$), 1.75 (*d*, $J = 6.4$ Hz, $\text{Me}(6''')$), 5.71 (br. s, $\text{H}-\text{C}(1')$), 6.04 (*d*, $J = 1.8$ Hz, $\text{H}-\text{C}(1''')$), and 6.08 (br. s, $\text{H}-\text{C}(1''''')$), and a xylopyranosyl moiety at $\delta(\text{H})$ 5.10 (*d*, $J = 7.7$ Hz, $\text{H}-\text{C}(1''''')$), together with two AcO groups at $\delta(\text{H})$ 1.98 (2s, 2 Me) and a 3HBO group at $\delta(\text{H})$ 1.30 (*d*, $J = 6.3$ Hz, $\text{Me}(4)$), 2.68 (*dd*, $J = 5.5, 15.8$ Hz, 1 $\text{H}-\text{C}(2)$), 2.78 (*dd*, $J = 7.5, 15.8$ Hz, 1 $\text{H}-\text{C}(2)$), and 5.49–5.51 (*m*, $\text{H}-\text{C}(3)$). The H- and C-atom signals in the ^1H - and ^{13}C -NMR spectra of **4** resembled those of **3**, except for the signals due to the additional AcO group. Comparison of the ^{13}C -NMR spectra of **4** with those of **3** revealed an acetylation shift around the 3-position of the 3HBO–C(4'') group (**4**: $\delta(\text{C})$ 40.8 (C(2)), 67.5 (C(3)), and 19.9 (C(4)); **3**: $\delta(\text{C})$ 45.0 (C(2)), 64.8 (C(3)), and 24.0 (C(4))). The connectivity of the above-mentioned AcO group in **4** was clarified by an HMBC experiment, which showed $^1\text{H} \rightarrow ^{13}\text{C}$ long-range correlations between $\delta(\text{H})$ 1.98 (s, Me of Ac) and $\text{H}-\text{C}(3)$ of 3HBO–C(4'') and $\delta(\text{C})$ 170.0 (C=O of Ac). In turn, the H- and C-atom signals in the ^1H - and ^{13}C -NMR spectra of **5** (Tables 3 and 4) resembled those of **4**, except for the signals due to the inner Rha of the 28-*O*-sugars. The position of an AcO group at the inner Rha of **5** was established by an HMBC experiment, which showed $^1\text{H} \rightarrow ^{13}\text{C}$ long-range correlations between $\delta(\text{H})$ 2.05 (s, Me of Ac) and 5.84 (*dd*, $J = 2.5, 8.9$ Hz, $\text{H}-\text{C}(3''')$ of the inner Rha) and $\delta(\text{C})$ 170.9 (C=O of Ac). On the basis of the above-mentioned evidence, the structures of **4** and **5** were elucidated to be 3-*O*- α -L-rhamnopyranosylpolygalactic acid 28-{*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-2-*O*-acetyl- α -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-4-*O*-[(3*S*)-3-(acetyloxy)butanonyl]- β -D-fucopyranosyl} ester and 3-*O*- α -L-rhamnopyranosylpolygalactic acid 28-{*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-3-*O*-acetyl- α -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-4-*O*-[(3*S*)-3-(acetyloxy)butanonyl]- β -D-fucopyranosyl} ester, respectively.

The molecular formulas of compound **6**, $\text{C}_{69}\text{H}_{110}\text{O}_{31}$, and compound **7**, $\text{C}_{73}\text{H}_{116}\text{O}_{33}$, were determined from their respective positive- and negative-ion-mode FAB-MS and by positive-ion-mode HR-FAB-MS data. Alkaline hydrolysis of **6** and **7** with 10% aqueous KOH in $\text{H}_2\text{O}/1,4$ -dioxane 1 : 1 provided **1a** and two organic acids, AcOH and 3HBOH, respectively. Treatment of **6** and **7** with 0.5% MeONa/MeOH gave **1b**, together with (+)-(3*S*)-3-hydroxybutanoic acid. The H- and C-atom signals in the ^1H - and ^{13}C -NMR spectra of **6** (Tables 3 and 4) showed signals assignable to a bellissaponin BS1 (**1b**) moiety, an AcO group at $\delta(\text{H})$ 1.99 (s, Me) and two 3HBO groups at $\delta(\text{H})$ 1.35 (*d*, $J = 6.2$ Hz, $\text{Me}(4)$), 1.38 (*d*, $J = 6.9$ Hz, $\text{Me}(4')$), 2.61–2.65 and 2.71–2.75 (2 *m*, $\text{CH}_2(2')$), 2.68–2.72 and 2.82–2.86 (2 *m*, $\text{CH}_2(2)$), 4.53–4.57 (*m*, $\text{H}-\text{C}(3')$), and 5.59–5.61 (*m*, $\text{H}-\text{C}(3)$). The connectivities of the acyl moieties in **6** were determined through an HMBC experiment, which exhibited $^1\text{H} \rightarrow ^{13}\text{C}$ long-range correlations between the following pairs: $\delta(\text{H})$ 1.99 (s, Me of Ac) and 6.01–6.03 (*m*, $\text{H}-\text{C}(2''')$ of the inner Rha)/ $\delta(\text{C})$ 170.5 (C=O of Ac), $\text{CH}_2(2)$ of 3HBO–C(4''), $\text{H}-\text{C}(3)$ of 3HBO–C(4''), and $\delta(\text{H})$ 5.53 (br. *d*, $J \approx 3$ Hz, $\text{H}-\text{C}(4'')$ of Fuc)/ $\delta(\text{C})$ 170.8 (C=O of 3HBO–C(4'')), and $\text{CH}_2(2')$ of 3HBO–C(3), $\text{H}-\text{C}(3')$ of 3HBO–C(3), and $\text{H}-\text{C}(3)$ of 3HBO–C(4'')/ $\delta(\text{C})$ 171.3 (C=O of 3HBO–C(3)). By comparison of

Table 3. Selected ^1H -NMR Data (600 MHz, 40°, (D₅)pyridine) of **4–7**. δ in ppm, J in Hz.

	4	5	6	7
H–C(2)	4.68–4.72 (<i>m</i>)	4.71 (br. <i>d</i> , $J \approx 3$)	4.74–4.78 (<i>m</i>)	4.70 (br. <i>s</i>)
H–C(3)	4.35 (<i>d</i> , $J = 3.5$)	4.39 (<i>d</i> , $J = 2.9$)	4.41 (br. <i>s</i>)	4.36 (<i>d</i> , $J = 3.2$)
H–C(12)	5.65 (<i>t</i> -like, $J \approx 3$)	5.65 (<i>t</i> -like, $J \approx 3$)	5.67 (<i>t</i> -like, $J \approx 3$)	5.63 (<i>t</i> -like, $J \approx 3$)
H–C(16)	5.12 (br. <i>s</i>)	5.14 (br. <i>s</i>)	5.16 (br. <i>s</i>)	5.12 (br. <i>s</i>)
H–C(18)	3.39 (<i>dd</i> , $J = 4.3, 14.1$)	3.40 (<i>dd</i> , $J = 3.4, 11.9$)	3.41 (<i>dd</i> , $J = 4.2, 13.7$)	3.40 (<i>dd</i> -like)
CH ₂ (23)	3.69 (br. <i>s</i>)	3.70 (br. <i>s</i>)	3.72 (br. <i>s</i>)	3.70 (br. <i>s</i>)
Me(24)	1.22 (<i>s</i>)	1.24 (<i>s</i>)	1.25 (<i>s</i>)	1.25 (<i>s</i>)
Me(25)	1.58 (<i>s</i>)	1.57 (<i>s</i>)	1.60 (<i>s</i>)	1.59 (<i>s</i>)
Me(26)	1.18 (<i>s</i>)	1.17 (<i>s</i>)	1.19 (<i>s</i>)	1.18 (<i>s</i>)
Me(27)	1.75 (<i>s</i>)	1.75 (<i>s</i>)	1.76 (<i>s</i>)	1.75 (<i>s</i>)
Me(29)	0.95 (<i>s</i>)	0.94 (<i>s</i>)	0.94 (<i>s</i>)	0.94 (<i>s</i>)
Me(30)	1.06 (<i>s</i>)	1.07 (<i>s</i>)	1.04 (<i>s</i>)	1.06 (<i>s</i>)
<i>3-O-Rha</i> :				
H–C(1')	5.71 (br. <i>s</i>)	5.72 (br. <i>s</i>)	5.79 (br. <i>s</i>)	5.72 (br. <i>s</i>)
Me(6')	1.60 (<i>d</i> , $J = 6.1$)	1.60 (<i>d</i> , $J = 6.0$)	1.62 (<i>d</i> , $J = 6.2$)	1.60 (<i>d</i> , $J = 6.0$)
<i>28-O-Sugars</i> :				
<i>Fuc</i> :				
H–C(1'')	5.99 (<i>d</i> , $J = 8.0$)	6.03 (<i>d</i> , $J = 7.6$)	6.02 (<i>d</i> , $J = 8.2$)	6.00 (<i>d</i> , $J = 8.0$)
H–C(4'')	5.50 (br. <i>d</i> , $J \approx 4$)	5.48 (br. <i>d</i> , $J \approx 3$)	5.53 (br. <i>d</i> , $J \approx 3$)	5.50 (br. <i>s</i>)
Me(6'')	1.25 (<i>d</i> , $J = 6.3$)	1.25 (<i>d</i> , $J = 6.0$)	1.27 (<i>d</i> , $J = 6.2$)	1.25 (<i>d</i> , $J = 6.3$)
<i>3HBO–C(4'')</i> :				
CH ₂ (2)	2.68 (<i>dd</i> , $J = 5.5, 15.8$), 2.78 (<i>dd</i> , $J = 7.5, 15.8$)	2.68 (<i>dd</i> , $J = 4.6, 13.1$), 2.78 (<i>dd</i> , $J = 6.2, 13.1$)	2.68–2.72 (<i>m</i>), 2.82–2.86 (<i>m</i>)	2.65–2.70 (<i>m</i>) ^a , 2.74–2.81 (<i>m</i>) ^b
H–C(3)	5.49–5.51 (<i>m</i>)	5.50–5.52 (<i>m</i>)	5.59–5.61 (<i>m</i>)	5.52–5.56 (<i>m</i>) ^c
Me(4)	1.30 (<i>d</i> , $J = 6.3$)	1.30 (<i>d</i> , $J = 6.3$)	1.35 (<i>d</i> , $J = 6.2$)	1.34 (<i>d</i> , $J = 6.3$) ^d
AcO–C(3)	1.98 (<i>s</i>)	2.00 (<i>s</i>)		
<i>3HBO–C(3)</i> :				
CH ₂ (2')			2.61–2.65 (<i>m</i>), 2.71–2.75 (<i>m</i>)	2.65–2.70 (<i>m</i>) ^a , 2.74–2.81 (<i>m</i>) ^b
H–C(3')			4.53–4.57 (<i>m</i>)	5.52–5.56 (<i>m</i>) ^c
Me(4')			1.38 (<i>d</i> , $J = 6.9$)	1.31 (<i>d</i> , $J = 6.3$) ^d
<i>3HBO–C(3'')</i> :				
CH ₂ (2'')				2.60 (<i>dd</i> , $J = 5.2, 14.9$), 2.74–2.81 (<i>m</i>) ^b
H–C(3'')				4.52–4.54 (<i>m</i>)
Me(4'')				1.38 (<i>d</i> , $J = 6.1$)
<i>Inner Rha</i> :				
H–C(1''')	6.04 (<i>d</i> , $J = 1.8$)	6.03 (br. <i>s</i>)	6.10 (br. <i>s</i>)	6.05 (br. <i>s</i>)
H–C(2''')	5.94 (<i>dd</i> , $J = 1.8, 3.5$)	4.95–4.97 (<i>m</i>)	6.01–6.03 (<i>m</i>)	5.94 (<i>dd</i> , $J = 1.5, 3.2$)
H–C(3''')	4.73 (<i>dd</i> , $J = 3.5, 9.4$)	5.84 (<i>dd</i> , $J = 2.5, 8.9$)	4.80–4.84 (<i>m</i>)	4.74 (<i>dd</i> , $J = 3.2, 9.5$)
Me(6''')	1.75 (<i>d</i> , $J = 6.4$)	1.74 (<i>d</i> , $J = 6.4$)	1.78 (<i>d</i> , $J = 6.8$)	1.75 (<i>d</i> , $J = 6.4$)
AcO–C(2''')	1.98 (<i>s</i>)		1.99 (<i>s</i>)	1.99 (<i>s</i>)
AcO–C(3''')		2.05 (<i>s</i>)		
<i>Xyl</i> :				
H–C(1''')	5.10 (<i>d</i> , $J = 7.7$)	4.95 (<i>d</i> , $J = 7.6$)	5.16 (<i>d</i> , $J = 6.8$)	5.10 (<i>d</i> , $J = 7.7$)
<i>Terminal Rha</i> :				
H–C(1''')	6.08 (br. <i>s</i>)	6.13 (br. <i>s</i>)	6.18 (br. <i>s</i>)	6.08 (br. <i>s</i>)
Me(6''')	1.67 (<i>d</i> , $J = 6.3$)	1.65 (<i>d</i> , $J = 6.2$)	1.65 (<i>d</i> , $J = 6.2$)	1.62 (<i>d</i> , $J = 6.3$)

^a)^b)^c) Overlapped. ^d) May be interchangeable within the same column.

Table 4. ^{13}C -NMR Data of **4–7**. At 150 MHz, 40°, in (D₅)pyridine; δ in ppm.

	4	5	6	7		4	5	6	7
C(1)	44.8	44.8	44.9	44.8	28-O-Sugars:				
C(2)	71.0	71.0	71.1	71.0	Fuc:				
C(3)	81.6	81.6	81.4	81.6	C(1'')	94.4	94.2	94.3	94.4
C(4)	43.0	43.0	43.0	43.0	C(2'')	75.0	76.7	74.9	75.0
C(5)	47.7	47.7	47.6	47.7	C(3'')	73.3	73.4	73.2	73.3
C(6)	18.4	18.3	18.3	18.4	C(4'')	74.8	74.7	74.7	74.8
C(7)	33.3	33.2	33.3	33.3	C(5'')	70.3	70.3	70.3	70.4
C(8)	40.3	40.3	40.3	40.3	C(6'')	16.4	16.4	16.4	16.4
C(9)	47.6	47.6	47.6	47.6	3HBO–C(4'')				
C(10)	37.2	37.2	37.2	37.2	C(1)	170.7	170.7	170.8	170.7
C(11)	24.1	24.1	24.1	24.1	C(2)	40.8	40.8	40.7	40.7 ^{a)}
C(12)	122.8	122.8	122.8	122.8	C(3)	67.5	67.5	67.4	67.9 ^{b)}
C(13)	144.3	144.2	144.3	144.3	C(4)	19.9	19.8	19.8	19.8 ^{c)}
C(14)	42.4	42.2	42.4	42.4	AcO–C(3)	170.0, 21.1	170.1, 21.1		
C(15)	36.2	36.1	36.2	36.2	3HBO–C(3):				
C(16)	74.1	74.2	74.1	74.2	C(1')			171.3	169.7
C(17)	49.6	49.5	49.5	49.6	C(2')			45.2	41.2 ^{a)}
C(18)	41.7	41.6	41.6	41.7	C(3')			64.4	67.6 ^{b)}
C(19)	47.4	47.4	47.3	47.4	C(4')			23.9	19.9 ^{c)}
C(20)	30.8	30.7	30.8	30.8	3HBO–C(3')				
C(21)	36.0	36.0	36.0	36.0	C(1'')				171.4
C(22)	31.9	31.7	32.0	31.9	C(2'')				45.1
C(23)	65.4	65.3	65.3	65.4	C(3'')				64.4
C(24)	15.0	15.0	15.0	15.0	C(4'')				23.9
C(25)	17.8	17.7	17.8	17.7	Inner Rha:				
C(26)	17.7	17.7	17.7	17.7	C(1''')	98.8	102.5	98.7	98.7
C(27)	27.1	27.1	27.1	27.1	C(2''')	73.3	69.5	73.2	73.3
C(28)	176.2	176.3	176.2	176.2	C(3''')	70.1	75.5	70.1	70.0
C(29)	33.1	33.1	33.1	33.1	C(4''')	84.0	77.9	83.7	83.9
C(30)	24.7	24.7	24.5	24.7	C(5''')	68.6	68.9	68.6	68.6
3-O-Rha:					C(6''')	18.5	19.1	18.5	18.5
C(1')	104.1	104.0	104.2	104.1	AcO–C(2''')	170.4, 20.9		170.5, 20.9	170.4, 20.9
C(2')	72.5	72.4	72.5	72.5	AcO–C(3''')		170.9, 21.3		
C(3')	72.8	72.8	72.8	72.8	Xyl:				
C(4')	74.0	74.1	74.0	74.0	C(1''')	106.8	105.7	106.7	106.8
C(5')	70.3	70.2	70.3	70.4	C(2''')	76.1	75.0	76.2	76.1
C(6')	18.6	18.5	18.6	18.6	C(3''')	83.7	83.6	83.3	83.7
					C(4''')	69.3	69.8	69.3	69.3
					C(5''')	67.3	67.1	67.3	67.3
					Terminal Rha:				
					C(1''')	102.6	102.7	102.6	102.6
					C(2''')	72.4	72.4	72.4	72.4
					C(3''')	72.6	72.7	72.6	72.6
					C(4''')	74.0	74.0	74.0	74.0
					C(5''')	70.0	69.8	70.0	70.0
					C(6''')	18.5	18.5	18.6	18.5

^{a)}^{b)}^{c)} Interchangeable within the same column.

the ^1H - and ^{13}C -NMR spectra of **7** (Tables 3 and 4) with those of **6**, **7** had one more 3HBO group than **6** at $\delta(\text{H})$ (1.31 (*d*, $J = 6.3$ Hz), 1.34 (*d*, $J = 6.3$ Hz), 1.38 (*d*, $J = 6.1$ Hz), Me(4'), Me(4), and Me(4'')), 2.65–2.70 and 2.74–2.81 (2 *m*, 2 H each, CH₂(2) and CH₂(2'')), 2.60 (*dd*, $J = 5.2$, 14.9 Hz, 1 H–C(2'')), 2.74–2.81 (*m*, 1 H–C(2'')), 4.52–4.54 (*m*, H–C(3'')), and 5.52–5.56 (*m*, H–C(3) and H–C(3')). In the HMBC experiment of **7**, the following $^1\text{H} \rightarrow ^{13}\text{C}$ long-range correlations were observed: CH₂(2'') of 3HBO–C(3'), H–C(3') of 3HBO–C(3'), and H–C(3') of 3HBO–C(3)/ $\delta(\text{C})$ 171.4 (C=O of 3HBO–C(3')). Consequently, the structures of **6** and **7** were elucidated to be 3-*O*- α -L-rhamnopyranosylpolygalacic acid 28-*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-2-*O*-acetyl- α -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-4-*O*-[(3*S*)-3-[(3*S*)-3-(hydroxybutanoyl)oxy]butanoyl]- β -D-fucopyranosyl ester and 3-*O*- α -L-rhamnopyranosylpolygalacic acid 28-*O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 4)-*O*-2-*O*-acetyl- α -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-4-*O*-[(3*S*)-3-[(3*S*)-3-[(3*S*)-3-hydroxybutanoyl]oxy]butanoyl]oxy]butanoyl]- β -D-fucopyranosyl ester, respectively.

Pancreatic lipase is well known to play an important role in lipid digestion. Recently, inhibitory effects of saponins on pancreatic lipase were reported to be involved in anti-obese effects of saponins (*e.g.*, theasaponins, chikusetsusaponins) [28–31]. In the course of our characterization studies on anti-obese constituents from *B. perennis*, inhibitory effects of the constituents on pancreatic lipase activity were examined. Among the saponin constituents, perennisaponins G (**1**; IC_{50} 163 μM), H (**2**; 137 μM), I (**3**; 147 μM), J (**4**; 148 μM), K (**5**; 223 μM), L (**6**; 81.4 μM), and M (**7**; 195 μM) were found to inhibit pancreatic lipase activity (Table 5). These saponins **1**–**7** are more

Table 5. Inhibitory Effects of the MeOH Extract and of **1**–**7** from the Flowers of *B. perennis* against Pancreatic Lipase

	Concentration [$\mu\text{g}/\text{ml}$] ^{a)}					IC_{50} [$\mu\text{g}/\text{ml}$]
	0	100	200	400	800	
MeOH Extract	0.0 $\pm$ 0.6	8.3 $\pm$ 1.8	21.4 $\pm$ 2.1 ^{c)}	43.8 $\pm$ 1.9 ^{c)}	71.6 $\pm$ 4.5 ^{c)}	455
	Concentration [μM] ^{a)}					IC_{50} [μM]
	0	50	100	200	400	
Perennisaponin G (1)	0.0 $\pm$ 2.8	– 18.8 $\pm$ 10.4	21.3 $\pm$ 5.8	58.9 $\pm$ 3.7 ^{c)}	85.5 $\pm$ 1.7 ^{c)}	163
Perennisaponin H (2)	0.0 $\pm$ 1.5	– 33.5 $\pm$ 12.2	14.7 $\pm$ 9.3	71.9 $\pm$ 3.1 ^{c)}	87.4 $\pm$ 1.3 ^{c)}	137
Perennisaponin I (3)	0.0 $\pm$ 1.1	– 17.0 $\pm$ 18.9	27.5 $\pm$ 8.9	60.2 $\pm$ 2.4 ^{c)}	85.2 $\pm$ 4.1 ^{c)}	147
Perennisaponin J (4)	0.0 $\pm$ 3.0	16.6 $\pm$ 13.6	38.3 $\pm$ 10.0 ^{b)}	55.9 $\pm$ 5.1 ^{c)}	55.2 $\pm$ 8.6 ^{c)}	148
Perennisaponin K (5)	0.0 $\pm$ 3.8	35.0 $\pm$ 5.0 ^{c)}	47.1 $\pm$ 5.0 ^{c)}	43.1 $\pm$ 10.1 ^{c)}	55.6 $\pm$ 6.5 ^{c)}	223
Perennisaponin L (6)	0.0 $\pm$ 4.2	26.9 $\pm$ 7.2 ^{b)}	55.3 $\pm$ 2.8 ^{c)}	52.7 $\pm$ 5.1 ^{c)}	68.5 $\pm$ 6.9 ^{c)}	81.4
Perennisaponin M (7)	0.0 $\pm$ 5.7	22.6 $\pm$ 12.0	28.5 $\pm$ 11.3	53.9 $\pm$ 3.8 ^{c)}	67.2 $\pm$ 5.6 ^{c)}	195
	Concentration [μM] ^{a)}					IC_{50} [μM]
	0	100	200	400	800	
Theasaponin E ₁	0.0 $\pm$ 2.2	24.4 $\pm$ 4.3 ^{b)}	39.2 $\pm$ 7.9 ^{c)}	57.4 $\pm$ 9.1 ^{c)}	88.6 $\pm$ 2.6 ^{c)}	270

^{a)} Values represent the means $\pm$ s.e.m ($N = 4$). Significantly different from the control group: ^{b)} $p < 0.05$. ^{c)} $p < 0.01$.

efficient than theasaponin E₁ (IC_{50} 270 μ M) [30], which was isolated from *Camellia sinensis* [32], although their inhibitory activities were considerably weaker than that of the lipase inhibitor orlistat (IC_{50} 56 nM). On the basis of above *in vitro* results and our previous *in vivo* evidence [2], the saponin constituents from the flowers of *B. perennis* may be useful for the prevention of obesity.

O. M., T. M., and K. N. were supported by ‘High-tech Research Center’ Project for Private Universities: matching fund subsidy from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan, 2007–2011, and by a Grant-in Aid for Scientific Research from MEXT. M. Y., H. M., and S. N. were supported by the 21st COE Program, the Academic Frontier Project, and a Grant-in Aid for Scientific Research from MEXT. H. M. was also supported by the Hoh-ansha Foundation, Japan.

Experimental Part

General. Column chromatography (CC): normal-phase CC with silica gel BW-200 (SiO₂; 150–300 mesh; Fuji Silysia Chemical, Ltd., Japan), reversed-phase CC with Chromatorex ODS DM1020T (100–200 mesh; Fuji Silysia Chemical, Ltd., Japan). HPLC: Shimadzu-RID-6A refractive-index, SPD-10A UV/VIS, and Shodex-OR-2 optical-rotation detectors; Shimadzu-LC-6AD pump; Shimadzu-CTO-10A column oven; Shimadzu-C-R6A chromatopac; Cosmosil-5C₁₈-MS-II and -HILIC (Nacalai Tesque, Inc.), Wakopak-Navi-C-30-5 (Wako Pure Chemical Industries Ltd.), and YMC-Pack-ODS-A and YMC-Pack-ODS-AQ (YMC Co., Ltd.) columns; t_R in min. Optical rotations: Horiba-SEPA-300 digital polarimeter ($l = 5$ cm). IR Spectra: Shimadzu-FTIR-8100 spectrometer; in cm^{-1} . ¹H- and ¹³C-NMR Spectra: Jeol-JNM-ECA600 spectrometer; at 600 and 150 MHz, resp.; δ in ppm rel. to Me₄Si, J in Hz. FAB- and HR-FAB-MS: Jeol-JMS-SX-102A mass spectrometer; in m/z .

Plant Material. The flowers of *B. perennis* cultivated in Albania were purchased from Tochimoto Tenkaido Co., Ltd., in November 2006, as described previously [2][3].

Extraction and Isolation. Compounds **1**–**7** were isolated from previously reported fractions [2][3], Fr. 6.11 (4.850 g) and Fr. 6.12 (50.269 g), originally obtained from the MeOH-eluted fraction (6.4%, 140.0 g) of the MeOH extract from flowers of *B. perennis*. An aliquot of Fr. 6.11 (1820.0 mg) was purified by HPLC (Wakopak-Navi-C30-5, 250 $\times$ 20 mm i.d., MeCN/MeOH/1% aq. AcOH 32:16:52; 9.0 ml/min) to afford the following eight fractions: Fr. 6.11.1 (465.8 mg), Fr. 6.11.2 (272.1 mg), Fr. 6.11.3 (41.7 mg), Fr. 6.11.4 (44.5 mg), Fr. 6.11.5 (263.0 mg), Fr. 6.11.6 (210.0 mg), Fr. 6.11.7 (137.9 mg), and Fr. 6.11.8 (88.6 mg). Fr. 6.11.5 (263.0 mg) was purified by HPLC (Cosmosil 5C₁₈-MS-II, 250 $\times$ 20 mm i.d., MeCN/MeOH/1% aq. AcOH 32:16:52; 9.0 ml/min): perennisaponin I (**3**; 33.1 mg, 0.0038%; t_R 48.9). Fr. 6.11.7 (137.9 mg) was purified by HPLC (Cosmosil 5C₁₈-MS-II, 250 $\times$ 20 mm i.d., MeCN/MeOH/1% aq. AcOH 32:16:52; 9.0 ml/min): perennisaponin G (**1**, 42.0 mg, 0.0048%; t_R 64.1). Fr. 6.11.8 (88.6 mg) was purified by HPLC (Cosmosil 5C₁₈-MS-II, 250 $\times$ 20 mm i.d., MeCN/MeOH/1% aq. AcOH 32:16:52; 9.0 ml/min): perennisaponin L (**6**, 24.3 mg, 0.0028%; t_R 70.0). An aliquot of Fr. 6.12 (2015.0 mg) was further separated by HPLC (Wakopak-Navi-C30-5, 250 $\times$ 20 mm i.d., MeCN/1% aq. AcOH 40:60; 9.0 ml/min) to give the following six fractions: Fr. 6.12.1 (893.3 mg), Fr. 6.12.2 (324.4 mg), Fr. 6.12.3 (131.3 mg), Fr. 6.12.4 (350.4 mg), Fr. 6.12.5 (48.8 mg), and Fr. 6.12.6 (135.0 mg). Fr. 6.12.1 (893.3 mg) was separated by HPLC (Cosmosil 5C₁₈-MS-II, 250 $\times$ 20 mm i.d., MeCN/MeOH/1% aq. AcOH 32:16:52; 9.0 ml/min) to furnish the following fourteen fractions: Fr. 6.12.1.1 (13.0 mg), Fr. 6.12.1.2 (9.4 mg), Fr. 6.12.1.3 (34.5 mg), Fr. 6.12.1.4 (10.5 mg), Fr. 6.12.1.5 (23.0 mg), Fr. 6.12.1.6 (80.8 mg), Fr. 6.12.1.7 (49.7 mg), Fr. 6.12.1.8 (86.8 mg), Fr. 6.12.1.9 (34.9 mg), Fr. 6.12.1.10 (125.8 mg, 0.139%; perennisaponin H (**2**); t_R 58.2), Fr. 6.12.1.11 (57.9 mg), Fr. 6.12.1.12 (109.2 mg, 0.119%; perennisaponin G (**1**); t_R 64.0), Fr. 6.12.1.13 (56.0 mg), and Fr. 6.12.1.14 (22.2 mg). An aliquot of Fr. 6.12.2 (180.0 mg) was purified by HPLC (Cosmosil HILIC, 250 $\times$ 20 mm i.d., MeCN/H₂O 90:10; 9.9 ml/min): perennisaponins J (**4**, 12.7 mg, 0.026%; t_R 49.9) and M (**7**, 31.7 mg, 0.065%; t_R 52.7) together with perennisaponins C (9.3 mg, 0.019%; t_R 65.2), D (8.1 mg, 0.017%; t_R 57.0), E (10.4 mg, 0.021%; t_R 59.2), and F (28.5 mg, 0.059%; t_R 55.7).

Fr. 6.12.3 (131.3 mg) was purified by HPLC (*Cosmosil HILIC*, 250 × 20 mm i.d., MeCN/H₂O 90:10; 9.9 ml/min): perennisaponins J (**4**, 32.6 mg, 0.034%; *t_R* 54.2) and K (**5**, 25.6 mg, 0.027%; *t_R* 57.1).

Perennisaponin G (= (2β,3β,4α,16α)-3-[(6-Deoxy-α-L-mannopyranosyl)oxy]-2,16,23-trihydroxy-olean-12-en-28-oic Acid O-6-Deoxy-α-L-mannopyranosyl-(1 → 3)-O-β-D-xylopyranosyl-(1 → 4)-O-2-O-acetyl-6-deoxy-α-L-mannopyranosyl-(1 → 2)-4-O-acetyl-6-deoxy-β-D-galactopyranosyl Ester; **1**): Amorphous powder. $[\alpha]_D^{25} = -19.5$ (*c* = 4.48, MeOH). IR (KBr): 3450, 1736, 1638, 1049. ¹H- and ¹³C-NMR: *Tables 1* and 2. FAB-MS (pos.): 1327 ($[M + Na]^+$). FAB-MS (neg.): 1303 ($[M - H]^-$), 1157 ($[M - C_6H_{11}O_4]^-$). HR-FAB-MS (pos.): 1327.6296 ($[M + Na]^+$, C₆₃H₁₀₀NaO₂₈; calc. 1327.6299).

Perennisaponin H (= (2β,3β,4α,16α)-3-[(6-Deoxy-α-L-mannopyranosyl)oxy]-2,16,23-trihydroxy-olean-12-en-28-oic Acid O-6-Deoxy-α-L-mannopyranosyl-(1 → 3)-O-β-D-xylopyranosyl-(1 → 4)-O-2-O-acetyl-6-deoxy-α-L-mannopyranosyl-(1 → 2)-4-O-acetyl-6-deoxy-β-D-galactopyranosyl Ester; **2**): Amorphous powder. $[\alpha]_D^{25} = -30.9$ (*c* = 4.43, MeOH). IR (KBr): 3450, 1734, 1638, 1049. ¹H- and ¹³C-NMR: *Tables 1* and 2. FAB-MS (pos.): 1327 ($[M + Na]^+$). FAB-MS (neg.): 1303 ($[M - H]^-$), 1157 ($[M - C_6H_{11}O_4]^-$), 649 ($[M - C_{27}H_{43}O_{18}]^-$). HR-FAB-MS (pos.): 1327.6305 ($[M + Na]^+$, C₆₃H₁₀₀NaO₂₈; calc. 1327.6299).

Perennisaponin I (= (2β,3β,4α,16α)-3-[(6-Deoxy-α-L-mannopyranosyl)oxy]-2,16,23-trihydroxy-olean-12-en-28-oic Acid O-6-Deoxy-α-L-mannopyranosyl-(1 → 3)-O-β-D-xylopyranosyl-(1 → 4)-O-2-O-acetyl-6-deoxy-α-L-mannopyranosyl-(1 → 2)-6-deoxy-4-O-[(3S)-3-hydroxy-1-oxobutyl]-β-D-galactopyranosyl Ester; **3**): Amorphous powder. $[\alpha]_D^{25} = -22.0$ (*c* = 2.28, MeOH). IR (KBr): 3450, 1738, 1638, 1049. ¹H- and ¹³C-NMR: *Tables 1* and 2. FAB-MS (pos.): 1371 ($[M + Na]^+$). FAB-MS (neg.): 1347 ($[M - H]^-$), 1115 ($[M - C_{10}H_{17}O_6]^-$), 649 ($[M - C_{29}H_{47}O_{19}]^-$). HR-FAB-MS (pos.): 1371.6558 ($[M + Na]^+$, C₆₅H₁₀₄NaO₂₉; calc. 1371.6561).

Perennisaponin J (= (2β,3β,4α,16α)-3-[(6-Deoxy-α-L-mannopyranosyl)oxy]-2,16,23-trihydroxy-olean-12-en-28-oic Acid O-6-Deoxy-α-L-mannopyranosyl-(1 → 3)-O-β-D-xylopyranosyl-(1 → 4)-O-2-O-acetyl-6-deoxy-α-L-mannopyranosyl-(1 → 2)-4-O-[(3S)-3-(acetyloxy)-1-oxobutyl]-6-deoxy-β-D-galactopyranosyl Ester; **4**): Amorphous powder. $[\alpha]_D^{25} = -48.5$ (*c* = 1.20, MeOH). IR (KBr): 3445, 1734, 1638, 1049. ¹H- and ¹³C-NMR: *Tables 3* and 4. FAB-MS (pos.): 1413 ($[M + Na]^+$). FAB-MS (neg.): 1389 ($[M - H]^-$), 649 ($[M - C_{31}H_{49}O_{20}]^-$). HR-FAB-MS (pos.): 1413.6664 ($[M + Na]^+$, C₆₇H₁₀₆NaO₃₀; calc. 1413.6667).

Perennisaponin K (= (2β,3β,4α,16α)-3-[(6-Deoxy-α-L-mannopyranosyl)oxy]-2,16,23-trihydroxy-olean-12-en-28-oic Acid O-6-Deoxy-α-L-mannopyranosyl-(1 → 3)-O-β-D-xylopyranosyl-(1 → 4)-O-3-O-acetyl-6-deoxy-α-L-mannopyranosyl-(1 → 2)-4-O-[(3S)-3-(acetyloxy)-1-oxobutyl]-6-deoxy-β-D-galactopyranosyl Ester; **5**): Amorphous powder. $[\alpha]_D^{25} = -47.1$ (*c* = 1.00, MeOH). IR (KBr): 3445, 1734, 1638, 1049. ¹H- and ¹³C-NMR: *Tables 3* and 4. FAB-MS (pos.): 1413 ($[M + Na]^+$). FAB-MS (neg.): 1389 ($[M - H]^-$), 649 ($[M - C_{31}H_{49}O_{20}]^-$). HR-FAB-MS (pos.): 1413.6674 ($[M + Na]^+$, C₆₇H₁₀₆NaO₃₀; calc. 1413.6667).

Perennisaponin L (= (2β,3β,4α,16α)-3-[(6-Deoxy-α-L-mannopyranosyl)oxy]-2,16,23-trihydroxy-olean-12-en-28-oic Acid O-6-Deoxy-α-L-mannopyranosyl-(1 → 3)-O-β-D-xylopyranosyl-(1 → 4)-O-2-O-acetyl-6-deoxy-α-L-mannopyranosyl-(1 → 2)-6-deoxy-4-O-[(3S)-3-[(3S)-3-hydroxy-1-oxobutyl]oxy]-1-oxobutyl]-β-D-galactopyranosyl Ester; **6**): Amorphous powder. $[\alpha]_D^{25} = -22.4$ (*c* = 1.64, MeOH). IR (KBr): 3450, 1736, 1655, 1049. ¹H- and ¹³C-NMR: *Tables 3* and 4. FAB-MS (pos.): 1457 ($[M + Na]^+$). FAB-MS (neg.): 1433 ($[M - H]^-$), 1115 ($[M - C_{14}H_{23}O_8]^-$), 649 ($[M - C_{33}H_{53}O_{21}]^-$), 503 ($[M - C_{39}H_{62}O_{25}]^-$). HR-FAB-MS (pos.): 1457.6937 ($[M + Na]^+$, C₆₉H₁₁₀NaO₃₁; calc. 1457.6929).

Perennisaponin M (= (2β,3β,4α,16α)-3-[(6-Deoxy-α-L-mannopyranosyl)oxy]-2,16,23-trihydroxy-olean-12-en-28-oic Acid O-6-Deoxy-α-L-mannopyranosyl-(1 → 3)-O-β-D-xylopyranosyl-(1 → 4)-O-2-O-acetyl-6-deoxy-α-L-mannopyranosyl-(1 → 2)-6-deoxy-4-O-[(3S)-3-[(3S)-3-[(3S)-3-hydroxy-1-oxobutyl]oxy]-1-oxobutyl]oxy]-1-oxobutyl]-β-D-galactopyranosyl Ester; **7**): Amorphous powder. $[\alpha]_D^{25} = -24.2$ (*c* = 1.42, MeOH). IR (KBr): 3445, 1736, 1655, 1049. ¹H- and ¹³C-NMR: *Tables 3* and 4. FAB-MS (pos.): 1543 ($[M + Na]^+$). FAB-MS (neg.): 1519 ($[M - H]^-$), 649 ($[M - C_{37}H_{59}O_{23}]^-$). HR-FAB-MS (pos.): 1543.7290 ($[M + Na]^+$, C₇₃H₁₁₆NaO₃₃; calc. 1543.7297).

Alkaline Hydrolysis of 1–7. A soln. of each perennisaponins **1–7** (6 mg) in H₂O/1,4-dioxane 1:1 (1 ml) was treated with 10% aq. KOH soln. (1 ml), and the mixture was stirred at 40° for 12 h. The mixture was neutralized over *Dowex-HCR-W2* resin (H⁺ form), which was then removed by filtration.

The filtrate was concentrated, and the resulting product subjected to reversed-phase CC (2 g of *ODS*; $\text{H}_2\text{O} \rightarrow \text{MeOH}$) to afford H_2O - and MeOH -eluted fractions, resp. The H_2O -eluted fraction was dissolved in $\text{ClCH}_2\text{CH}_2\text{Cl}$ (2 ml). This soln. was treated with *N,N'*-diisopropyl *O*-(4-nitrobenzyl)-isourea (= (4-nitrophenyl)methyl *N,N'*-bis(1-methylethyl)carbamimidate; 10 mg) and stirred at 80° for 1 h. The mixture was then subjected to HPLC (*YMC-Pack ODS-A*, 250 × 4.6 mm i.d., $\text{MeOH}/\text{H}_2\text{O}$ 65 : 35; 0.8 ml/min, UV detection at 254 nm): 4-nitrobenzyl esters of 3-hydroxybutanoic acid (**a**; t_R 7.0) from **3–7** and of AcOH (**b**; t_R 8.3) from **1–7**. The MeOH -eluted fraction was subjected to normal-phase CC (2 g of SiO_2 , $\text{CHCl}_3/\text{MeOH}/\text{H}_2\text{O}$ 10 : 3 : 1, lower layer) to give (2 β ,3 β ,4 α ,16 α)-3-[(6-deoxy- α -L-mannopyranosyl)oxy]-2,16,23-trihydroxyolean-12-en-28-oic acid (**1a**; 2 mg each from **1–7**).

Deacylation of 1–7. A soln. of each perennissaponin **1–7** (2 mg) in 0.5% MeONa/MeOH (1 ml) was stirred at r.t. for 3 h. An aliquot of the mixture was subjected to HPLC (*YMC-Pack ODS-AQ*, 250 × 4.6 mm i.d.; $\text{MeOH}/\text{H}_2\text{O}$ 20 : 80; 0.7 ml/min, optical-rotation detector): methyl (+)-(3*S*)-3-hydroxybutanoate (t_R 9.2, pos. optical rotation) from **3–7**. The rest of each mixture was neutralized over *Dowex-HCR-W2* resin (H^+ form), which was then removed by filtration. The filtrate was concentrated and the resulting product purified by HPLC (*Cosmosil 5C₁₈-MS-II*, 250 × 20 mm i.d., $\text{MeCN}/\text{MeOH}/\text{H}_2\text{O}$ 32 : 16 : 52; 9.0 ml/min): bellissaponin BS1 (**1b**; 1.5 mg each from **1–7**).

Effect on Pancreatic Lipase Activity. A suspension of triolein (80 mg), phosphatidylcholine (10 mg), and sodium taurocholate (5 mg) in 9 ml of 0.1M *Tris*·HCl buffer (pH 7.0) containing 0.1M NaCl was homogeneously emulsified with a homogenizer (strait *Teflon* pestle with strait glass tube, volume 20 ml). The substrate suspension (0.1 ml) in a test tube was preincubated with 5 μl of test sample in DMSO and 95 μl of *Tris*·HCl buffer for 3 min at 37°. An aliquot of porcine pancreatic lipase (250 $\mu\text{g}/\text{ml}$, type II, *Sigma–Aldrich, Inc.*; 50 μl) or *Tris*·HCl buffer (50 μl) as a blank test was then added to start the reaction. After 30 min of incubation, the test tube was immediately immersed in boiling water for 2 min to stop the reaction and then cooled with water. Free fatty acid concentration was determined by a commercial kit (*Wako NEFA C* test, *Wako Pure Chemical Industries, Ltd.*). Theasaponin E_1 was isolated from *Camellia sinensis* [32] and used as a reference compound [28]. IC_{50} was determined graphically ($N=4$).

Statistics. Values were expressed as means $\pm$ s.e.m. For statistical analysis, one-way analysis of variance followed by *Dunnnett's* test was used. Probability (*p*) values less than 0.05 were considered significant.

REFERENCES

- [1] L.-B. Wang, T. Morikawa, S. Nakamura, K. Ninomiya, H. Matsuda, O. Muraoka, L.-J. Wu, M. Yoshikawa, *Heterocycles* **2009**, 78, 1235.
- [2] T. Morikawa, X. Li, E. Nishida, Y. Ito, H. Matsuda, S. Nakamura, O. Muraoka, M. Yoshikawa, *J. Nat. Prod.* **2008**, 71, 828.
- [3] M. Yoshikawa, X. Li, E. Nishida, S. Nakamura, H. Matsuda, O. Muraoka, T. Morikawa, *Chem. Pharm. Bull.* **2008**, 56, 559.
- [4] H. Matsuda, T. Morikawa, T. Ohgushi, T. Ishiwada, N. Nishida, M. Yoshikawa, *Chem. Pharm. Bull.* **2005**, 53, 387.
- [5] M. Yoshikawa, T. Morikawa, K. Yamamoto, Y. Kato, A. Nagatomo, H. Matsuda, *J. Nat. Prod.* **2005**, 68, 1360.
- [6] M. Yoshikawa, S. Sugimoto, S. Nakamura, H. Matsuda, *Chem. Pharm. Bull.* **2007**, 55, 571.
- [7] M. Yoshikawa, F. Xu, T. Morikawa, Y. Pongpiriyadacha, S. Nakamura, Y. Asao, H. Matsuda, *Chem. Pharm. Bull.* **2007**, 55, 308.
- [8] S. Nakamura, S. Sugimoto, H. Matsuda, M. Yoshikawa, *Heterocycles* **2007**, 71, 577.
- [9] M. Yoshikawa, S. Nakamura, Y. Kato, K. Matsuhira, H. Matsuda, *Chem. Pharm. Bull.* **2007**, 55, 598.
- [10] M. Yoshikawa, T. Morikawa, Y. Asao, E. Fujiwara, S. Nakamura, H. Matsuda, *Chem. Pharm. Bull.* **2007**, 55, 606.
- [11] M. Yoshikawa, S. Sugimoto, S. Nakamura, H. Sakumae, H. Matsuda, *Chem. Pharm. Bull.* **2007**, 55, 1034.

- [12] S. Nakamura, S. Sugimoto, H. Matsuda, M. Yoshikawa, *Chem. Pharm. Bull.* **2007**, *55*, 1342.
- [13] M. Yoshikawa, T. Wang, S. Sugimoto, S. Nakamura, A. Nagatomo, H. Matsuda, S. Harima, *Yakugaku Zasshi* **2008**, *128*, 141.
- [14] M. Yoshikawa, S. Sugimoto, Y. Kato, S. Nakamura, T. Wang, C. Yamashita, H. Matsuda, *Chem. Biodiversity* **2009**, *6*, 903.
- [15] M. Yoshikawa, S. Sugimoto, S. Nakamura, H. Matsuda, *Chem. Pharm. Bull.* **2008**, *56*, 1297.
- [16] Y. Xie, T. Morikawa, K. Ninomiya, K. Imura, O. Muraoka, D. Yuan, M. Yoshikawa, *Chem. Pharm. Bull.* **2008**, *56*, 1628.
- [17] S. Nakamura, Y. Okazaki, K. Ninomiya, T. Morikawa, H. Matsuda, M. Yoshikawa, *Chem. Pharm. Bull.* **2008**, *56*, 1704.
- [18] S. Sugimoto, M. Yoshikawa, S. Nakamura, H. Matsuda, *Heterocycles* **2009**, *78*, 1023.
- [19] S. Sugimoto, G. Chi, Y. Kato, S. Nakamura, H. Matsuda, M. Yoshikawa, *Chem. Pharm. Bull.* **2009**, *57*, 269.
- [20] T. Morikawa, L.-B. Wang, S. Nakamura, K. Ninomiya, E. Yokoyama, H. Matsuda, O. Muraoka, L.-J. Wu, M. Yoshikawa, *Chem. Pharm. Bull.* **2009**, *57*, 361.
- [21] T. Schöpke, C. Al-Tawaha, V. Wray, M. Nimiz, K. Hiller, *Phytochemistry* **1997**, *45*, 125.
- [22] M. J. Burk, T. G. P. Harper, C. S. Kalberg, *J. Am. Chem. Soc.* **1995**, *117*, 4423.
- [23] T. Morikawa, H. Matsuda, T. Ohgushi, N. Nishida, T. Ishiwada, M. Yoshikawa, *Heterocycles* **2004**, *63*, 2211.
- [24] M. Yoshikawa, T. Wang, T. Morikawa, H. Xie, H. Matsuda, *Chem. Pharm. Bull.* **2007**, *55*, 1308.
- [25] K. Ninomiya, T. Morikawa, H. Xie, H. Matsuda, M. Yoshikawa, *Heterocycles* **2008**, *75*, 1983.
- [26] T. Morikawa, H. Xie, T. Wang, H. Matsuda, M. Yoshikawa, *Chem. Pharm. Bull.* **2008**, *56*, 1438.
- [27] T. Morikawa, H. Xie, T. Wang, H. Matsuda, M. Yoshikawa, *Chem. Biodiversity* **2009**, *6*, 411.
- [28] L.-K. Han, Y. Kimura, M. Kawashima, T. Takaku, T. Taniyama, T. Hayashi, Y.-N. Zheng, H. Okuda, *Int. J. Obes.* **2001**, *25*, 1459.
- [29] L.-K. Han, Y.-N. Zheng, M. Yoshikawa, H. Okuda, Y. Kimura, *BMC Complem. Altern. Med.* **2005**, *5*, 9.
- [30] T. Morikawa, Y. Xie, Y. Asao, M. Okamoto, C. Yamashita, O. Muraoka, H. Matsuda, Y. Pongpiriyadacha, D. Yuan, M. Yoshikawa, *Phytochemistry* **2009**, *70*, 1166.
- [31] Y. Asao, T. Morikawa, Y. Xie, M. Okamoto, M. Hamao, M. Matsuda, O. Muraoka, D. Yuan, M. Yoshikawa, *Chem. Pharm. Bull.* **2009**, *57*, 198.
- [32] I. Kitagawa, K. Hori, T. Motozawa, T. Murakami, M. Yoshikawa, *Chem. Pharm. Bull.* **1998**, *46*, 1901.

Received July 9, 2009