This article was downloaded by: [McMaster University] On: 27 October 2014, At: 11:35 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/lsyc20

Mild and Solvent-Free Alkynylation of Ketones on the KF/Alumina

Ali Sharifi $^{\rm a}$, Mojtaba Mirzaei $^{\rm a}$ & M. Reza Naimi-Jamal $^{\rm a}$

^a Chemistry & Chemical Engineering Research Center of Iran, Tehran, Iran Published online: 16 Aug 2006.

To cite this article: Ali Sharifi , Mojtaba Mirzaei & M. Reza Naimi-Jamal (2005) Mild and Solvent-Free Alkynylation of Ketones on the KF/Alumina, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 35:8, 1039-1044, DOI: <u>10.1081/SCC-200054195</u>

To link to this article: http://dx.doi.org/10.1081/SCC-200054195

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

Synthetic Communications[®], 35: 1039–1044, 2005 Copyright © Taylor & Francis, Inc. ISSN 0039-7911 print/1532-2432 online DOI: 10.1081/SCC-200054195



Mild and Solvent-Free Alkynylation of Ketones on the KF/Alumina

Ali Sharifi, Mojtaba Mirzaei, and M. Reza Naimi-Jamal

Chemistry & Chemical Engineering Research Center of Iran, Tehran, Iran

Abstract: The solvent-free alkynylation of various ketones on the surface of KF/alumina under mild conditions is described.

Keywords: Alkynes, alkynylation, ketones, KF/alumina, solvent-free

INTRODUCTION

Carbon–carbon bond formations are among the most important reactions in organic synthesis.^[1] One significant example is alkynylation of carbonyl compounds for the preparation of propargylic alcohols. The traditional approach to these reactions involves the addition of alkynylide anions to carbonyl compounds. This process requires the conversion of alkynes to corresponding alkynylide anions using 1 equiv. of a strong air-sensitive base such as Grignard reagent,^[2] alkyl lithium,^[3] or sodium amide^[4] in an anhydrous solvent. Adding the carbonyl compound gradually to this mixture prevents the competitive aldol reaction. More recent methods, making use of catalytic amounts of bases such as *t*-BuOK,^[5] cesium hydroxide,^[6] zinc alkynylides,^[7] and quaternary ammonium hydroxide^[8] are more favorable. However, these reactions suffer from the lack of generality and require the use of a solvent. Replacement of the organic solvents with ionic liquids,^[9]

Received in Poland October 25, 2004

Address correspondence to Ali Sharifi, Chemistry & Chemical Engineering Research Center of Iran, P.O. Box 14335-186, Tehran, Iran. Tel.: +98-21-8036144; Fax: +98-21-8037185; E-mail: sharifi@ccerci.ac.ir; M. Reza Naimi-Jamal, Institut für Reine und Angew. Chemie, Universität Oldenburg, Oldenburg, Postfach 2503, D-26111 Oldenburg, Germany (current address). E-mail: naimi@uni.oldenburg.de supercritical CO_2 ,^[10] water,^[11] or an adsorbent^[12] seems to be more economical and ecological, and is garnering special interest. Recently, Miyamoto et al. have reported that the addition of ethynylbenzene to ketones using potassium *t*-butoxide or potassium hydroxide proceeds efficiently in the absence of a solvent.^[13] However, the generality of this method has yet to be investigated for aliphatic acetylenes.

Earlier studies have revealed that potassium fluoride (KF), either alone or in the form of an adsorbed reagent, can be employed as a mild, effective, and nonnucleophilic base in organic reactions. In 1979, Yamawaki and Ando introduced KF/alumina as a useful, versatile reagent in alkylation reactions.^[14] So far, the usefulness of this reagent has been widely investigated in many C–O, C–N, and C–C bond formation reactions.^[15] With our continuing interest in developing practical, solvent-free reactions in mind,^[16] we decided to investigate this reagent. We now report the reaction of terminal alkynes with ketones at room temperature using KF/alumina.

RESULTS AND DISCUSSION

This reaction of terminal alkynes with ketones proceeds on the surface of KF/alumina without solvent in up to 95% yield (Scheme 1). The results are summarized in Table 1. The solvent-free reaction appears to be more efficient than the solution phase reaction. For example, phenyl acetylene **1a** reacts with cyclohexanone **2h** on the surface of KF/alumina at room temperature to give 95% yield of the corresponding alcohol (Table 1, entry 8), whereas for the same substrates in DMSO using KOtBu as a base, 83% yield has been reported.^[13] The reaction of less acidic acetylenes fails when employing these conditions. For example, addition of 1-hexyne and



3a: R¹=Ph, R²=Me, R³=Me; **3b:** R¹=Ph, R²=Me, R³=Et; **3c:** R¹=Ph, R²=Et, R³=Et **3d:** R¹=Ph, R²=Me, R³=*i*-Pr; **3e:** R¹=Ph, R²=*i*-Pr, R³=*i*-Pr; **3f:** R¹=Ph, R²=Me, R³=*n*-penthyl **3g:** R¹=Ph, R²=*n*-Bu, R³=*n*-Bu; **3h:** R¹=Ph, R², R³= -CH₂-(CH₂)₃-CH₂- **3i:** R¹=Ph, R², R³= -CH₂-(CH₂)₄-CH₂-; **3j:** R¹=PhOCH₂, R²=Et, R³=Et **3k:** R¹=PhOCH₂, R²=Me, R³=*i*-Pr; **3l:** R¹=PhOCH₂, R²=*i*-Pr, R³=*i*-Pr **3m:** R¹=*n*-Pentyl, R²=Me, R³=*n*-penthyl; **3n:** R¹=*n*-Hexyl, R²=*n*-Bu, R³=*n*-Bu **3o:** R¹=2,4-DimethylphenylOCH₂, R², R³= -CH₂-(CH₂)₃-CH₂-; **3p:** R¹=1-NaphthylOCH₂, R², R³= -CH₂-(CH₂)₃-CH₂-

Scheme 1.

Entry	Alkyne	Ketone	Product ^a	Time/h (temp)	Yield $(\%)^b$
1	1a	2a	3a ^[8]	20 (rt)	90
2	1 a	2b	3b ^[13]	20 (rt)	95
3	1 a	2c	3 c ^[8]	20 (rt)	90
4	1a	2d	3d ^[8]	15 (rt)	42
				15 (60°C)	90
5	1 a	2e	3e ^[13]	17 (rt)	32
				15 (60°C)	95
6	1a	2f	3f ^[3a]	17 (rt)	46
7	1a	2g	3g ^[20]	17 (rt)	30
				15 (60°C)	93
8	1a	2h	3h ^[13]	17 (rt)	95
9	1a	2i	3i ^[13]	20 (rt)	40
				40 (rt)	70
10	1b	2h	3j ^[19]	20 (60°C)	90
11	1b	2 b	3k ^[19]	20 (60°C)	95
12	1b	2a	3l ^[19]	5 (rt)	20
				17 (rt)	70
13	1c	2h	3m	20 (rt)	с
14	1d	2h	3n	20 (rt)	С
15	1e	2h	30	20 (60°C)	92
16	1f	2h	3 p	20 (60°C)	90

Table 1. Solvent-free alkynylation of ketones on the surface of KF/alumina

^bIsolated yields.

^cNo reaction occurred.

1-heptyne to the cyclohexanone does not occur after 20 h at room temperature (Table 1, entries 13, 14).

The method is also suitable for functionalized acetylenes such as propargyl ethers **1b** and **1e–f**. They react with ketones to give the corresponding tertiary alcohols with moderate to high yields. These reactions are more efficient at higher temperatures (Table 1, entries 10, 15, 16). Raising the reaction temperature considerably affects the reaction rate and yield. For example, the addition of phenyl acetylene **1a** to the ketone **2g** results in 30% and 93% yields of the product **3g**, at room temperature and 60°C, respectively (Table 1, entry 7).

Using unsupported KF instead of KF/alumina dramatically decreases the reaction yields. The reason for this change might be explained by the lower basicity of unsupported KF compared with KF/alumina.^[17] According to Ando,^[17b] the basicity of KF/alumina not only results from the presence of active F⁻ (as in KF itself), but also from generated OH⁻ and Al–OH.

^{*a*}Mp and/or spectral data (IR, ¹H-NMR and ¹³C-NMR) of known compounds (**3a–1**) were compared with given references.

In summary, we report here that KF/alumina is a very mild and efficient catalyst to promote the addition of primary alkynes to ketones, in an environmentally friendly reaction, without need of any reaction solvent and at room temperature.

EXPERIMENTAL

1042

Mps were determined on a hot stage or oil-bath apparatus without correction. ¹H-NMR (80 MHz or 300 MHz) and ¹³C-NMR (75 MHz) spectra were recorded on a Bruker 80 MHz or Bruker WP 300 MHz in CDCl₃ using TMS as internal standard. HRMS were obtained on a Finnigan MAT system MAT 212.

Preparation of KF/Alumina^[18]

Neutral alumina (60–80 mesh, 30 g) in water (150 ml) is added to a stirred solution of potassium fluoride (20 g) in water (150 ml). After 30 min the water is evaporated in a rotary evaporator at \sim 60°C. When most of the water has been removed, the remaining mixture is heated to 140–150°C and maintained at that temperature under vacuum (5 mmHg) for 6 h to give 50 g of KF/alumina reagent.

General Procedure for the Preparation of Tertiary Alcohols from Alkynes and Ketones

Alkyne (1 mmol), ketone (1.2 mmol), and KF/alumina (1 g) were mixed in a 25-mL flask at room or mentioned temperature. The progress of reaction was monitored by GC. After the given time, the reaction mixture was washed with petrol and filtered, and the solvent was evaporated on the rotavapour. The residue was purified with the bulb-to-bulb distillation technique or with a short silica-gel column (eluent: light petroleum ether/ethyl acetate, 20:1).

30. Colorless oil, IR (KBr, ν , cm⁻¹): 3200–3500, 3020, 2932, 2856, 1256, 1215, 1130, 801. ¹H-NMR (CDCl₃): $\delta = 1.41-1.84$ (m, 10H), 2.15 (s, 3H), 2.19 (s, 3H), 2.30 (s, 1H), 4.83 (s, 2H), 6.60–6.80 (m, 3H_{arom}). ¹³C-NMR (CDCl₃): $\delta = 15.85$, 20.14, 22.88 (2C), 24.90, 39.47 (2C), 56.47, 58.50, 78.50, 91.34, 112.38, 126.63, 128.59, 130.63, 131.35, 154.93. MS (EI): m/z = 55 (46), 77 (38), 91 (40), 107 (42), 122 (100%), 136 (20), 160 (20), 258 (33) [M⁺]. HRMS (EI): m/z calcd. for C₁₇H₂₂O₂: 258.1620; found: 258.1620.

3p. Colorless oil, IR (KBr, ν , cm⁻¹): 3200–3500, 3055, 2925, 2858, 1580, 1504, 1266, 1228, 1173, 1097, 1018, 791. ¹H-NMR (CDCl₃):

Alkynylation of Ketones on KF/Alumina

$$\begin{split} &\delta = 1.43 - 1.75 \text{ (m, 10H)}, 2.10 \text{ (s, 1H)}, 4.86 \text{ (s, 2H)}, 6.80 \text{ (m, 1H}_{arom}), 7.30 - 7.50 \text{ (m, 4H}_{arom}), 7.70 - 7.80 \text{ (m, 1H}_{arom}), 8.20 - 8.40 \text{ (m, 1H}_{arom}). ^{13}\text{C-NMR} \\ &\text{(CDCl}_3): \ \delta = 22.95 \text{ (2C)}, 24.90, 39.52 \text{ (2C)}, 56.20, 68.32, 78.91, 91.36, \\ &105.75, 120.85, 121.91, 124.80, 125.09, 125.39, 126.22, 127.25, 134.12, \\ &153.48. \text{ MS (EI): } m/z = 41 \text{ (38)}, 55 \text{ (35)}, 115 \text{ (28)}, 152 \text{ (88)}, 171 \text{ (62)}, 181 \\ &\text{(100)}, 209 \text{ (53)}, 233 \text{ (42)}, 262 \text{ (22)}, 280 \text{ (93) } [\text{M}^+]. \text{ HRMS (EI): } m/z \text{ calcd.} \\ &\text{for $C_{19}H_{20}O_2$: 280.1463; found: 280.1464.} \end{split}$$

REFERENCES

- Trost, B. M. Comprehensive Organic Synthesis; Trost, B. M., Fleming, I., Eds.; Pergamon Press: London, 1991; Vol. 3.
- (a) Viehe, H. G.; Reinstein, M. Dichloromethane as solvent for Grignard reagents. *Chemische Berichte* 1962, 95, 2557–2562; (b) Fantazier, R. M.; Poutsma, M. L. Reduction of propargylic chlorides with tri-n-butyltin hydride: The ambident behavior of propargylic radicals. *J. Am. Chem. Soc.* 1968, 90, 5490–5498.
- (a) Albert, S.; Robin, S.; Rousseau, G. Preparation of oxetanes by 4-endo trig electrophilic cyclizations of cinnamic alcohols. *Tetrahedron Lett.* 2001, *42* (13), 2477–2480; (b) Marson, C. M.; Walker, A. J.; Pickering, J.; Harper, S.; Wrigglesworth, R.; Edge, S. J. Efficient routes to cyclic 2,3-epoxy alcohols from cycloalkenyl ketones, via cycloalkenyl alcohols. *Tetrahedron* 1993, *49* (45), 10317–10338.
- Saunders, J. H. Organic Synthesis; Wiley: New York, 1955; Collect. Vol. III; p. 416.
- Babler, J. H.; Liptak, V. P.; Phan, N. Alkoxide-catalyzed addition of terminal alkynes to ketones. J. Org. Chem. 1996, 61, 416–417.
- Tzalis, D.; Knochel, P. Cesium hydroxide: A superior base for the catalytic alkynylation of aldehydes and ketones and catalytic alkenylation of nitriles. *Angew. Chem., Int. Ed. Engl.* 1999, 38, 1463–1465.
- (a) Anand, N. K.; Carreira, E. M. A simple, mild, catalytic, enantioselective addition of terminal acetylenes to aldehydes. J. Am. Chem. Soc. 2001, 123, 9687–9688;
 (b) Moore, D.; Pu, L. BINOL-catalyzed highly enantioselective terminal alkyne additions to aromatic aldehydes. Org. Lett. 2002, 4, 1855;
 (c) Li, X.; Lu, G.; Kwok, W. H.; Chan, A. S.C. Highly enantioselective alkynylzinc addition to aromatic aldehydes catalyzed by self-assembled titanium catalysts. J. Am. Chem. Soc. 2002, 124, 12636–12637.
- Ishikawa, T.; Mizuta, T.; Hagiwara, K.; Aikawa, T.; Kudo, T.; Saito, S. Catalytic alkynylation of ketones and aldehydes using quaternary ammonium hydroxide base. J. Org. Chem. 2003, 68, 3702–3705.
- (a) Welton, T. Room-temperature ionic liquids, solvents for synthesis and catalysis. *Chem. Rev.* **1999**, *99*, 2071–2083; (b) Wasserscheid, P.; Keim, W. Ionic liquids—new solutions for transition metal catalysis. *Angew. Chem., Int. Ed. Engl.* **2000**, *39*, 3772–3789.
- Oakes, R. S.; Clifford, A. A.; Rayner, C. M. The use of supercritical fluids in synthetic organic chemistry. J. Chem. Soc., Perkin Trans. 1 2001, 9, 917–941.
- Li, C. J.; Chan, T. H. Organic Reactions in Aqueous Media; John Wiley and Sons: New York, 1997.

- (a) Kropp, P. J.; Breton, G. W.; Fields, J. D.; Tung, J. C.; Loomis, B. R. Surfacemediated reactions, 8: Oxidation of sulfides and sulfoxides with tert-butyl hydroperoxide and OXONE. *J. Am. Chem. Soc.* 2000, *122* (18), 4280–4285; (b) Das, B.; Madhusudhan, P.; Venkataiah, B. Studies on novel synthetic methodologies, Part 4: An efficient microwave-assisted one-pot conversion of aldehydes into nitriles using silica-gel-supported NaHSO4 catalyst. *Synlett* 1999, *10*, 1569–1570.
- Miyamoto, H.; Yasaka, S.; Tanaka, K. Solvent-free addition of ethynylbenzene to ketones. *Bull. Chem. Soc. J.* 2001, 74, 185–186.
- Yamawaki, J.; Ando, T. Potassium fluoride on inorganic solid supports: A search for further efficient reagents promoting hydrogen-bond-assisted alkylations. *Chem. Lett.* 1979, 7, 755–758.
- Blass, B. E. KF/Al₂O₃ mediated organic synthesis. *Tetrahedron* 2002, 58, 9301–9320.
- (a) Sharifi, A.; Mojtahedi, M. M.; Saidi, M. R. Microwave irradiation techniques for the Cannizzaro reaction. *Tetrahedron Lett.* **1999**, *40*, 1179–1180;
 (b) Sharifi, A.; Mohsenzadeh, F.; Naimi-Jamal, M. R. Solvent-free preparation of monoacylaminals assisted by microwave irradiation. *J. Chem. Res., Synop.* **2000**, *8*, 394–396; (c) Sharifi, A.; Mirzaei, M.; Naimi-Jamal, M. R. Solvent-free aminoalkylation of phenols and indoles assisted by microwave irradiation. *Monatsh. Chem.* **2001**, *132*, 875–880; (d) Sharifi, A.; Mirzaei, M.; Naimi-Jamal, M. R. Copper-catalyzed oxidative homo-coupling of terminal acetylenes on alumina assisted by microwave irradiation. *J. Chem. Res., Synop.* **2002**, 628–630.
- (a) Weinstock, L. M.; Stevenson, J. M.; Tomellini, S. A.; Pan, S. H.; Utne, T.; Jobson, R. B.; Rienhold, D. F. Characterization of the actual catalytic agent in potassium fluoride on ativated alumina systems. *Tetrahedron Lett.* **1986**, 27 (33), 3845–3848; (b) Ando, T.; Clark, J. H.; Cork, D. G.; Hanafusa, T.; Ichihara, J.; Kimura, T. Fluoride-alumina reagents: The active basic species. *Tetrahedron Lett.* **1987**, 28, 1421–1424; (c) Kabashima, H.; Tsuji, H.; Nakata, S.; Tanaka, Y.; Hattori, H. Activity for base-catalyzed reactions and characterization of alumina-supported KF catalysts. *Appl. Catal., A* **2000**, *194–195*, 227–240.
- Paquette, L. A. Encyclopedia of Reagents for Organic Synthesis; John Wiley & Sons: Chichester, 1995; Vol. 6, p. 4223.
- Moskowitz, H.; Blanc-Guenee, J.; Miocque, M. Polyfunctional acetylenic derivatives: Tertiary carbinols with alkoxy and aminoalkoxy groups. C. R. Seances Acad. Sci., Ser. 3: Sci. 1967, 264, 1406.
- Kunishima, M.; Nakata, D.; Tanaka, S.; Hioki, K.; Tani, S. Generation and reactions of alkynylsamariums. *Tetrahedron* 2000, 56, 9927–9936.