This article was downloaded by: [Rensselaer Polytechnic Institute] On: 29 September 2013, At: 18:51 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

### Compliancy of 1,4-Naphthoquinone for the Synthesis of New Heterocyclic Quinone Compounds Containing Imino Linkage

N. A. A. El Kanzi<sup>a</sup>, A. K. Khalafallah<sup>a</sup>, Abu El Hamd<sup>a</sup>, H. Mohamed<sup>a</sup> & Soleiman A. Soleiman<sup>a</sup>

<sup>a</sup> Chemistry Department, Faculty of Science, South Valley University, Aswan, Egypt Published online: 07 Jun 2007.

To cite this article: N. A. A. El Kanzi , A. K. Khalafallah , Abu El Hamd , H. Mohamed & Soleiman A. Soleiman (2007) Compliancy of 1,4-Naphthoquinone for the Synthesis of New Heterocyclic Quinone Compounds Containing Imino Linkage, Phosphorus, Sulfur, and Silicon and the Related Elements, 182:7, 1483-1496, DOI: 10.1080/10426500701242921

To link to this article: http://dx.doi.org/10.1080/10426500701242921

#### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with

primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



#### Compliancy of 1,4-Naphthoquinone for the Synthesis of New Heterocyclic Quinone Compounds Containing Imino Linkage

N. A. A. El Kanzi A. K. Khalafallah Abu El Hamd H. Mohamed Soleiman A. Soleiman Chemistry Department, Faculty of Science, South Valley University, Aswan, Egypt

Some new  $\beta$ -lactam, thiazolidinone derivatives were synthesized by reaction of new Schiff bases with compounds **4** and oxazoles, pyrazoles, pyrimidines, and pyrimidinethion were also synthesized by reaction of an arylidino system with hydroxyl amine hydrochloride, hydrazines, urea, and thiourea, which we expected to have biological activities such as bactericidal and fungicidal or other applications of certain interest.

**Keywords** Arylidino derivatives; isolated  $\beta$ -lactams; oxazoles; pyrazoles; pyrimidines; pyrimidinethion; Schiff bases; thiazolidinone derivatives

#### INTRODUCTION

Three membered rings containing nitrogen, oxygen, and sulphur atoms as the heteroatom are called aziridine, oxirane, and thiirane, respectively. A duly examination of the literature for fused three membered heterocycles with other hetero rings indicated that such systems too reave one class of antibiotics.<sup>1</sup> Also  $\beta$ -lactam and related derivatives exhibit antibacterial activities.<sup>2–7</sup> Thiazole derivatives and  $\beta$ -lactam rings were known and used as potent antibiotics,<sup>8</sup> antibiotic activity,<sup>9</sup> and inhibitory activity.<sup>10</sup> Diverse biological activities, such as bactericidal, pesticidal, fungicidal, insecticidal, anticonvulsant, tuberculostatic, anti-inflammatory, and antithydroidal have been found to be associated with thiazolidinone derivatives.<sup>11</sup> Thus, our initial strategy in this

Received May 29, 2006; accepted December 11, 2006.

Address correspondence to A. K. Khalafallah, Chemistry Department, Faculty of Science, South Valley University, Answan, Egypt. E-mail: magdi5aziz@yahoo.com

research project was divested to the synthesis of different structural formulas corresponding to the isolated  $\beta$ -lactams, thiazolidinone derivatives, pyrazoles, oxazoles, pyrimidines, and pyrimidinethion.

#### **RESULTS AND DISCUSSION**

The known importance of these heterocyclic compounds with respect to various types of biological effects promysted us to synthesize new heterocyclic compounds depending upon our previous research in this field.<sup>12–14</sup> Thus, 1, 4-naphthoquinone **1** was reacted with acetamide and mercaptoacetic acid in ethanol at reflux in the presence of piperidine to give 2-amido-2-methyl-2-H-naphtho[2, 3-b][1, 4]thiazine-5, 10-dione, 3-carboxylic acid 4. The formation of 4 was assumed to proceed via the intermediate 2, which cyclizes to 4, (c.f., Scheme 1). The IR spectrum of 4 revealed the presence of absorption bands at 1665, 3421, 3069  $\rm cm^{-1}$ assigned to CO, NH, OH groups, respectively. The <sup>1</sup>H-NMR spectrum (DMSO-d6) were showed a singlet at  $\delta$  1.02 assigned for methyl group, a singlet at  $\delta$  3.43 for a methyl group of the acetyl group,  $\delta$  6.01 for (CH-S), multaiplets at  $\delta$  7.10–8.10 for aromatic protons, and a broad singlet at  $\delta$  9.01 for (NH) protons, respectively, with lack of a peak due to an NH<sub>2</sub> group. The mass spectrum of compound 4 showed m/z at 346 ( $M^+$ , 346). The compound 4 contains methyl groups which render it susceptible to react with different nitroso compounds such as  $\beta$ nitroso $\alpha$ -naphthol,  $\alpha$ -nitroso $\beta$ -naphthol, P-nitrophenol, and P-nitro-N.Ndimethyl-aniline in the presence of piperidine as catalyst to give the corresponding Schiff base derivatives 7a-d (Scheme 1). The reaction of 7a-d with an equimolar ratio of chloroacetylchloride in a mixture of DMF and ethanol in the presence of piperidine as a catalyst afforded  $\beta$ -lactam derivatives **8a–d** (Scheme 1). Cycloaddition reaction of thioglycolic acid to the previously prepared Schiff base compounds 7a-d in a mixture of ethanol and DMF in the presence of piperidine as catalyst afforded thiazolidinone derivatives 9a-d (Scheme 1). The active methyl group in the new compound 4 condensed with different aromatic aldehydes in a mixture of ethanol and DMF under piperidine as a catalyst to yield the corresponding arylidino **10a–c**. The arylidino derivatives **10a–** c, when interacted with hydrazine, hydrate in the presence of acetic acid in ethanol as solvents under piperidine as catalyst, respectively, gave the required N-acetylpyrazolo **11a-d**, and/or N-phenylpyrazolo derivatives **12a–d**, respectively (Scheme 2). The arylidino derivatives **10a– c** when interacted with hydroxylamine hydrochloride in a mixture of ethanol and DMF as solvent under the effect of sodium hydroxide as catalyst, gave the required isooxazolino derivatives **13a–c**. The arylidino





**SCHEME 1** 



#### **SCHEME 2**

derivatives **10a–c**, when interacted with urea and/or thiourea in a mixture of ethanol and DMF as solvent under the effect of sodium hydroxide as catalyst, gave the required pyrimidino derivatives **14a–c** or, under the effect of hydrochloric acid as catalyst, gave the required pyrimidinethion derivatives **15a–c**.

The activity of the carbonyl group in compound **4** renders it available to react with different aromatic amine such as aniline, P-chloroaniline, and P-nitroaniline in the presence of a mixture of ethanol and DMF





**SCHEME 3** 

as a solvent and piperidine catalyst to give new Schiff base derivatives **16a–c** (Scheme 3). The reaction of **16a–c** with equimolar ratios of chloroacetylchloride in mixture of ethanol and DMF in the presence of piperidine catalyst afforded lactam derivatives **17a–c** (Scheme 3). Cycloaddition reaction of thioglycolic acid to the previously prepared Schiff bases **16a–c** proceeded successfully. Thus thioglycolic acid added to Schiff bases **16a–c** in a mixture of ethanol and DMF in the presence of piperidine as catalyst and afforded thiazolidinone derivatives **18a–c** (Scheme 3).

#### **EXPERIMENTAL**

All melting points were uncorrected. IR spectra were recorded on a pye Unicam SP 1100 spectrophotometer using KBr disks. <sup>1</sup>HNMR spectra were recorded on a Varian EM390 MHz spectrophotometer using DMSO-d6 as a solvent and TMS as an internal standard. Chemical shifts are expressed as ppm units. Mass spectra were recorded on a HP Ms6988 spectrometer. Analytical data were determined with a CE 440 Elemental Analyzer-Automatic Indicator at Cairo University.

## Synthesis of 2-Amido-2-methyl-2-H-naphtho[2, 3-b][1, 4]thiazine-5, 10-dione, 3-carboxylic acid 4

A solution of acetamide (0.59 g, 0.01 mol) and mercaptoaceticacid (0.92 g, 0.01 mol) was treated with 1,4-naphthoquinone (1.58 g, 0.01 mol) in ethanol in the presence of piperidine as catalyst and refluxed for 15 h. The reaction mixture was filtered from unreacted materials. The filtered was poured into ice water acidified by concentrated hydrochloric acid. A reddish brown precipitate separated from the filtrate and was washed several times with water. It was crystallized from ethanol and dimethyl formamide to give compound **8a–d**, mp 290°C (c.f., Table I).

## Synthesis of 2-diazaamido-2-methyl-2-H-naphthol[2, 3-b][1, 4]thiazine-5, 10-dione, 3-carboxylic acid 7a–d

Compound 4 (3.46 g, 0.01 mol) and nitroso compounds (1.73 g, 1.73 g, 1.50 g, 0.01 mol) in an equimolar ratio were dissolved in ethanol and dimethylformamide, and 0.5 ml of piperidine as catalyst was added. The mixture was refluxed about 9 h. The reaction mixture was allowed to cool at room temperature, then filtered, washed several times with water, dried, collected, and crystallized from ethanol to give **7a-d** (c.f., Table I).

Compound number	Solvent of crystallization	Melting point	Yield %	Molecular formula	Molecular weight	Mass spectra	
4	EtOH/DMF	290	85%	$C_{16}H_{14}O_5N_2S$	346.06	348(M+2)	
7a	EtOH/DMF	115	75%	$\mathrm{C_{26}H_{19}O_6N_3S}$	501.1	502(M+1)	
7b	EtOH/DMF	120	75%	$\mathrm{C}_{26}\mathrm{H}_{19}\mathrm{O}_{6}\mathrm{N}_{3}\mathrm{S}$	501.1	502(M+1)	
7c	EtOH/DMF	125	80%	$\mathrm{C}_{22}\mathrm{H}_{17}\mathrm{O}_{6}\mathrm{N}_{3}\mathrm{S}$	451.09	452(M+1)	
7d	EtOH/DMF	122	75%	$\mathrm{C}_{24}\mathrm{H}_{22}\mathrm{O}_5\mathrm{N}_4\mathrm{S}$	478.13	478	
8a	EtOH/DMF	260	65%	$C_{28}H_{20}O_7N_3SCl$	577.07	579(M+2)	
8b	EtOH/DMF	260	65%	$C_{28}H_{20}O_7N_3SCl$	577.07	579(M+2)	
8c	EtOH/DMF	250	67%	$\mathrm{C}_{24}\mathrm{H}_{17}\mathrm{O}_{7}\mathrm{N}_{3}\mathrm{SCl}$	530.10	530	
8d	EtOH/DMF	270	70%	$C_{26}H_{23}O_6N_4SCl$	554.10	554	
9a	EtOH/DMF	360	70%	$C_{28}H_{21}O_7N_3S_2$	575.08	575	
9b	EtOH/DMF	360	73%	$C_{28}H_{21}O_7N_3S_2$	575.08	575	
9c	EtOH/DMF	320	75%	$C_{24}H_{19}O_7N_3S_2$	525.06	525	
9d	EtOH/DMF	340	72%	$\mathrm{C}_{26}\mathrm{H}_{24}\mathrm{O}_{6}\mathrm{N}_{4}\mathrm{S}_{2}$	552.11	552	
10a	EtOH/DMF	250	80%	$\mathrm{C}_{23}\mathrm{H}_{18}\mathrm{O}_5\mathrm{N}_2\mathrm{S}$	434.09	432(M-2)	
10b	EtOH/DMF	220	82%	$\mathrm{C}_{23}\mathrm{H}_{18}\mathrm{O}_{6}\mathrm{N}_{2}\mathrm{S}$	450.09	449(M-1)	
10c	EtOH/DMF	190	78%	$C_{23}H_{17}O_7N_3S$	479.08	479	
11a	EtOH/DMF	260	85%	$\mathrm{C}_{25}\mathrm{H}_{22}\mathrm{O}_5\mathrm{N}_4\mathrm{S}$	490.97	449(M+3)	
11b	EtOH/DMF	230	85%	$\mathrm{C}_{25}\mathrm{H}_{22}\mathrm{O}_6\mathrm{N}_4\mathrm{S}$	506.96	508(M+1)	
11c	EtOH/DMF	200	82%	$\mathrm{C}_{25}\mathrm{H}_{21}\mathrm{O}_{7}\mathrm{N}_{5}\mathrm{S}$	535.45	535	
12a	EtOH/DMF	230	70%	$\mathrm{C}_{29}\mathrm{H}_{24}\mathrm{O}_4\mathrm{N}_4\mathrm{S}$	524.15	524	
12b	EtOH/DMF	235	68%	$\mathrm{C}_{29}\mathrm{H}_{24}\mathrm{O}_{5}\mathrm{N}_{4}\mathrm{S}$	540.15	540	
12c	EtOH/DMF	225	70%	$C_{29}H_{23}O_6N_5S$	569.14	569	
13a	EtOH/DMF	200	75%	$C_{23}H_{19}O_5N_3S$	449.11	449	
13b	EtOH/DMF	210	73%	$\mathrm{C}_{23}\mathrm{H}_{18}\mathrm{O}_{6}\mathrm{N}_{3}\mathrm{S}$	465.1	465	
13c	EtOH/DMF	215	70%	$\mathrm{C}_{23}\mathrm{H}_{18}\mathrm{O}_{7}\mathrm{N}_{4}\mathrm{S}$	494.1	494	
14a	EtOH/DMF	220	78%	$\mathrm{C}_{24}\mathrm{H}_{20}\mathrm{O}_5\mathrm{N}_4\mathrm{S}$	476.11	478(M+2)	
14b	EtOH/DMF	225	73%	$\mathrm{C}_{24}\mathrm{H}_{20}\mathrm{O}_{6}\mathrm{N}_{4}\mathrm{S}$	492.11	492	
14c	EtOH/DMF	227	70%	$\mathrm{C}_{24}\mathrm{H}_{19}\mathrm{O}_{7}\mathrm{N}_{5}\mathrm{S}$	521.1	521	
15a	EtOH/DMF	215	75%	$C_{24}H_{20}O_4N_4S_2$	491.1	491	
15b	EtOH/DMF	220	75%	$C_{24}H_{20}O_5N_4S_2$	507.1	507	
15c	EtOH/DMF	220	73%	$C_{24}H_{19}O_6N_5S_2$	537.1	537	
16a	EtOH/DMF	240	76%	$\mathrm{C}_{22}\mathrm{H}_{19}\mathrm{O}_{4}\mathrm{N}_{3}\mathrm{S}$	421.1	423(M+2)	
16b	EtOH/DMF	242	74%	$\mathrm{C}_{22}\mathrm{H}_{18}\mathrm{O}_{4}\mathrm{N}_{3}\mathrm{SCl}$	455.1	455	
16c	EtOH/DMF	245	73%	$\mathrm{C}_{22}\mathrm{H}_{18}\mathrm{O}_6\mathrm{N}_4\mathrm{S}$	466.1	466	
17a	EtOH/DMF	210	72%	$\mathrm{C}_{24}\mathrm{H}_{20}\mathrm{O}_5\mathrm{N}_3\mathrm{SCl}$	497.1	497	
17b	EtOH/DMF	215	70%	$\mathrm{C}_{24}\mathrm{H}_{19}\mathrm{O}_5\mathrm{N}_3\mathrm{SCl}_2$	531.04	531	
17c	EtOH/DMF	212	70%	$\mathrm{C}_{24}\mathrm{H}_{19}\mathrm{O}_{7}\mathrm{N}_{4}\mathrm{SCl}$	542.1	542	
18a	EtOH/DMF	230	75%	$C_{24}H_{21}O_5N_3S_2$	495.1	495	
18b	EtOH/DMF	235	72%	$\mathrm{C}_{24}\mathrm{H}_{20}\mathrm{O}_5\mathrm{N}_3\mathrm{S2Cl}$	529.05	529	
18c	EtOH/DMF	240	70%	$\mathrm{C}_{24}\mathrm{H}_{19}\mathrm{O}_{7}\mathrm{N}_{4}\mathrm{S}_{2}$	540.1	540	

**TABLE I** Characterization of Compounds

#### Synthesis of New $\beta$ -Lactam Derivatives 8a–d

A solution of 7a-d (5.01 g, 0.01 mol) chloroacetylchloride (1.13 g, 0.01mol) in a mixture of ethanol and dimethylformamide in the

Compound number	<sup>1</sup> H-NMR d6 (DMSO)				
4	δ1.77(s,CH3), δ 3.74 (s,COCH3), δ 4.5 (s,NH), δ 7.01–9 (m,7H, Aromatic protons, CH,NH,OH)				
7a	$\delta1.63(\rm s,CH3),\delta$ 3.54 (s,CH), $\delta$ 6.01 (s,NH), $\delta$ 7.11–8.07 (m,13H, Aromatic protons, CH,NH,OH), $\delta$ 9.673 (s, Carboxylic OH)				
7b	$\delta1.60(\rm s,CH3),\delta$ 3.50 (s,CH), $\delta$ 6.02 (s,NH), $\delta$ 7.0–8.2 (m,13H, Aromatic protons, CH,NH,OH), $\delta$ 9.723 (s, Carboxylic OH)				
7c	δ1.4(s,CH3), δ 3.7 (s,CH), δ 5.8 (s,NH), δ 7.2–8.3 (m,11H, Aromatic protons, CH,NH,OH), δ 9.9 (s, Carboxylic OH)				
7d	$ \begin{split} &\delta 1.5({\rm s,CH3}),  \delta \; 2.34 \; ({\rm s,6H},2{\rm CH}_3),  \delta \; 3.56 \; ({\rm s,CH}),  \delta \; 5.6 \; ({\rm s,NH}),  \delta \; 7.1{-}8.0 \\ &({\rm m,10H},  {\rm Aromatic \ protons},  {\rm CH}, {\rm NH}),  \delta \; 9.92 \; ({\rm s, \ Carboxylic \ OH}) \end{split} $				
8a	$ \begin{split} &\delta 1.64(s,CH_3),  \delta  2.92  (s,CH),  \delta  3.41  (s,CH),  \delta  5.23  (s,NH),  \delta  7.05 - 8.7 \\ &(m,13H,  Aromatic  protons,  CH,NH,OH),  \delta  8.85  (s,  Carboxylic  OH) \end{split} $				
8b	$\begin{split} &\delta 1.45(s,CH_3), \delta \; 2.83\;(s,CH), \delta \; 3.35\;(s,CH), \delta \; 5.41\;(s,NH), \delta \; 7.1-8.8\;(m,13H, \\ &Aromatic\; protons,\; CH,NH,OH), \delta \; 8.9\;(s,\; Carboxylic\; OH) \end{split}$				
8c	$ \begin{split} &\delta 1.53(s,CH_3),  \delta  2.9  (s,CH),  \delta  3.3  (s,CH),  \delta  5.52  (s,NH),  \delta  78.3  (m,11H, \\ &Aromatic  protons,  CH,NH,OH),  \delta  8.98  (s,  Carboxylic  OH) \end{split} $				
8d	$ \begin{split} &\delta 1.56({\rm s}{\rm ,CH}_3), \delta \ 2.3 \ ({\rm s}{\rm ,6H}{\rm ,2CH}_3), \delta \ 2.98 \ ({\rm s}{\rm ,CH}), \delta \ 3.54 \ ({\rm s}{\rm ,CH}), \delta \ 5.64 \ ({\rm s}{\rm ,NH}), \\ &\delta \ 7-8 \ ({\rm m}{\rm ,10H}{\rm ,Aromatic \ protons, \ CH}{\rm ,NH},), \delta \ 9.5 \ ({\rm s}{\rm ,Carboxylic \ OH}) \end{split} $				
9a	$\delta 1.62$ (s,CH <sub>3</sub> ), $\delta 2.9$ (s,CH), $\delta 3.42$ (s,CH <sub>2</sub> ), $\delta 5.24$ (s,NH), $\delta 7.5$ –8.19 (m,13H, Aromatic protons, CH,NH,OH), $\delta 8.21$ (s, Carboxylic OH)				
9b	δ1.62(s,CH <sub>3</sub> ), δ 2.89 (s,CH), δ 3.42 (s,CH <sub>2</sub> ), δ 5.23 (s,NH), δ 7.5–8.2 (m,13H, Aromatic protons, CH,NH,OH), δ 8.32 (s, Carboxylic OH)				
9c	$\delta 1.63(s, CH_3), \delta 2.9(s, CH), \delta 3.43(s, CH_2), \delta 5.23(s, NH), \delta 7.5-8.18(m, 11H, Aromatic protons, CH, NH, OH), \delta 8.4(s, Carboxylic OH)$				
90	δ1.66(s,CH <sub>3</sub> ), δ 2.65 (s,6H,2CH <sub>3</sub> ), δ 2.95 (s,CH), δ 3.45 (s,CH <sub>2</sub> ), δ 5.4 (s,NH), δ 7.1–8.3 (m,10H, Aromatic protons, CH,NH,), δ 8.5 (s,Carboxylic OH)				
10a	$ \begin{split} &\delta 1.61(s,CH_3),  \delta \; 3.41\;(s,NH),  \delta \; 5.99\;(s,CH),  \delta \; 6.49\;(s,CH),  \delta \; 7.37\;(s,SCH),  \delta \\ & 7.45-82\;(m,10H,  Aromatic\; protons,  NH),  \delta \; 10.05\;(s,OH) \end{split} $				
10b	$\delta$ 1.6(s,CH <sub>3</sub> ), $\delta$ 3.8 (s,NH), $\delta$ 6.31 (s,CH), $\delta$ 6.89 (s,CH), $\delta$ 7.35 (s,SCH), $\delta$ 7.43–8 (m,10H, Aromatic protons, NH,OH), $\delta$ 10.24 (s,Carboxylic OH)				
10c	$ \begin{split} &\delta 1.8({\rm s},{\rm CH}_3),\delta\;4.2\;({\rm s},{\rm NH}),\delta\;6.65\;({\rm s},{\rm CH}),\delta\;6.9\;({\rm s},{\rm CH}),\delta\;7.3\;({\rm s},{\rm SCH}),\delta\\ &7.4{-}8.01\;({\rm m},{\rm 9H},{\rm Aromatic}\;{\rm protons},{\rm NH}),\delta\;10.47\;({\rm s},{\rm OH}) \end{split} $				
11a	$ \begin{split} &\delta 1.45(s,CH_3),  \delta  1.65  (s,COCH_3),  \delta  1.7  (s,CH_2),  \delta  2.4  (s,CH),  \delta  3.7(s,NH),  \delta \\ &7.03\text{-}8.54  (m,11H,Aromatic  protons,  CH,NH),  \delta  10.12  (s,OH) \end{split} $				
11b	$ \begin{split} &\delta 1.53({\rm s,CH_3}),  \delta  1.54  ({\rm s,COCH_3}),  \delta  1.8  ({\rm s,CH_2}),  \delta  2.45  ({\rm s,CH}),  \delta  3.73  ({\rm s,NH}), \\ &\delta  7.32-8.57  ({\rm m,11H},  {\rm Aromatic \ protons},  {\rm CH}, {\rm NH}, {\rm OH}),  \delta  10.2 \\ & ({\rm s,Carboxylic \ OH}) \end{split} $				
11c	$ \begin{split} &\delta 1.6(s,CH_3), \ \delta \ 1.7 \ (s,COCH_3), \ \delta \ 1.8 \ (s,CH_2), \ \delta \ 2.5 \ (s,CH), \ \delta \ 3.8 \ (s,NH), \ \delta \\ &7.3-8.6 \ (m,10H, \ Aromatic \ protons, \ CH,NH), \ \delta \ 10.3 \ (s, \ OH) \end{split} $				
12a	$ \begin{split} &\delta 1.47(s,CH_3),  \delta  1.9  (s,CH_2),  \delta  2.6  (s,CH),  \delta  4.01(s,NH),  \delta  7.18.7  (m,11H, \\ Aromatic protons,  CH,NH),  \delta  10.01  (s,  OH) \end{split} $				
12b	$ \begin{split} &\delta 1.46(s,CH_3), \delta  1.95  (s,CH_2), \delta  2.75  (s,CH), \delta  4.3  (s,NH), \delta  7.3-8.87 \\ &(m,11H,  Aromatic  protons,  CH,NH,OH), \delta  10.12  (s,  Carboxylic  OH) \end{split} $				
12c	$ \begin{split} &\delta 1.48(s,CH_3),  \delta  1.98(s,CH_2),  \delta  2.8(s,CH),  \delta  4.5(s,NH),  \delta  7.48.89(m,10H, \\ Aromatic  protons,  CH,NH),  \delta  10.3(s,OH) \end{split} $				

### TABLE II Selected HNMR Spectra Data for the New Compounds Listed in Table I

Compound number	<sup>1</sup> H-NMR d6 (DMSO)
13a	δ1.23(s,CH <sub>3</sub> ), δ 1.43(s,CH <sub>2</sub> ), δ 1.8 (s,CH), δ 3.5 (s,NH), δ 7.2–8.5 (m,11H, Aromatic protons, CH,NH), δ 10.2 (s,OH)
13b	$ \begin{split} &\delta 1.3({\rm s,CH_3}), \delta  1.45({\rm s,CH_2}), \delta  1.9  ({\rm s,CH}), \delta  3.7  ({\rm s,NH}), \delta  7.58.7  ({\rm m,11H}, \\ {\rm Aromatic \ protons,NH,OH}), \delta  10.  24  ({\rm s, \ Carboxylic \ OH}) \end{split} $
13c	$ \begin{split} &\delta 1.42(s,CH_3),  \delta  1.48(s,CH_2),  \delta  2.2(s,CH),  \delta  3.85(s,NH),  \delta  7.78.75 \\ &(m,10H,Aromaticprotons,CH,NH),  \delta  10.26(s,OH) \end{split} $
14a	$ \begin{split} &\delta 1.25({\rm s}{\rm ,CH}_3), \delta \; 1.6 \; ({\rm s}{\rm ,CH}_2), \delta \; 2.03 \; ({\rm s}{\rm ,CH}), \delta \; 3.6 \; ({\rm s}{\rm ,NH}), \delta \; 3.85 \; ({\rm s}{\rm ,NH}), \delta \; 7.23 &= 8.47 \; ({\rm m}{\rm ,11H}, \; {\rm Aromatic \ protons}{\rm ,CH}{\rm ,NH}), \; \delta \; 10.4 \; ({\rm s}{\rm ,OH}) \end{split} $
14b	$\delta$ 1.27(s,CH <sub>3</sub> ), $\delta$ 1.65 (s,CH <sub>2</sub> ), $\delta$ 2.12 (s,CH), $\delta$ 3.67 (s,NH), $\delta$ 3.89 (s,NH), $\delta$ 7.4–8.7 (m,11H, Aromatic protons, CH,NH,OH), $\delta$ 10.45 (s,Carboxylic OH)
14c	
15a	$\delta$ 1.05(s,CH <sub>3</sub> ), $\delta$ 1.5 (s,CH <sub>2</sub> ), $\delta$ 1.9 (s,CH), $\delta$ 3.5 (s,NH), $\delta$ 3.8 (s,NH), $\delta$ 7.4–8.6 (m,11H, Aromatic protons, CH,NH), $\delta$ 10.32 (s,OH)
15b	$\delta$ 1.08(s,CH <sub>3</sub> ), $\delta$ 1.56 (s,CH <sub>2</sub> ), $\delta$ 1.95 (s,CH), $\delta$ 3.84 (s,NH), $\delta$ 4.2 (s,NH), $\delta$ 7.65–8.85 (m,11H, Aromatic protons, CH,NH,OH), $\delta$ 10.45 (s, Carboxylic OH)
15c	
16a	$\delta 1.2 (\rm s, CH_3),  \delta  1.85  (\rm s, CH_3),  \delta  3.54  (\rm s, NH),  \delta  7.34-8.23  (\rm m, 11H,  Aromatic protons,  CH, NH),  \delta  9.57  (\rm s, OH)$
16b	$\delta 1.25(\rm s, CH_3),  \delta  1.9  (\rm s, CH_3),  \delta  3.98  (\rm s, NH),  \delta  7.6-8.5  (m, 10H,  Aromatic protons,  CH, NH),  \delta  9.7  (\rm s, OH)$
16c	δ1.4(s,CH <sub>3</sub> ), δ 2.3 (s,CH <sub>3</sub> ), δ 4.24 (s,NH), δ 7.7–8.5 (m,10H, Aromatic protons, CH,NH), δ 10.15 (s,OH)
17a	$ \begin{split} &\delta 1.6(s, CH_3),  \delta  1.75  (s, CH_3),  \delta  2.8  (s, CH),  \delta  3.7  (s, NH),  \delta  7.43 - 8.45  (m, 11H, \\ Aromatic  protons,  CH,  NH),  \delta  9.85  (s, OH) \end{split} $
17b	$ \begin{split} &\delta 1.7({\rm s},{\rm CH}_3),\delta \; 1.85 \;({\rm s},{\rm CH}_3),\delta \; 2.9 \;({\rm s},{\rm CH}),\delta \; 3.9 \;({\rm s},{\rm NH}),\delta \; 7.58.6 \;({\rm m},10{\rm H},\\ &{\rm Aromatic \ protons,\ CH,\ NH}),\delta \; 9.9 \;({\rm s},{\rm OH}) \end{split} $
17c	$ \begin{split} &\delta 1.8(s, CH_3), \ \delta \ 1.9 \ (s, CH_3), \ \delta \ 2.98 \ (s, CH), \ \delta \ 3.98 \ (s, NH), \ \delta \ 7.6-8.634 \\ &(m, 10H, \ Aromatic \ protons, \ CH, \ NH), \ \delta \ 9.95 \ (s, OH) \end{split} $
18a	$ \begin{split} &\delta 1.68(s,CH_3),  \delta  1.832  (s,CH_3),  \delta  2.04  (s,CH_2),  \delta  4.02  (s,NH),  \delta  7.45 - 8.23 \\ &(m,11H,  Aromatic  protons,  CH,NH),  \delta  10.2  (s,OH) \end{split} $
18b	$ \begin{split} &\delta 1.8(s,CH_3),\delta \; 1.867\;(s,CH_3),\delta \; 2.34\;(s,CH_2),\delta \; 4.21\;(s,NH),\delta \; 7.58.3\\ &(m,10H,Aromatic\;protons,CH,NH),\delta \; 10.25\;(s,OH) \end{split} $
18c	$\delta1.85(\rm{s,CH}_3),\delta$ 1.92 (s,CH_3), $\delta$ 2.5 (s,CH_2), $\delta$ 4.3 (s,NH), $\delta$ 7.6–8.5 (m,10H, Aromatic protons, CH,NH), $\delta$ 10.28 (s,OH)

### TABLE II Selected HNMR Spectra Data for the New Compounds Listed in Table I (Continued)

presence of a few drops of piperidine was refluxed for 10-12 h. The filtrate was evaporated under reduced pressure, poured into ice water where the product was separated, filtered, washed, several times with water, and crystallized from ethanol and dimethylformamide (c.f., Table I).

Comp.		Calculated % Found % $% \label{eq:calculated}$				2
no.	IR $v_{\rm max}$ Cm $^{-1}$	С	Η	Ν	S	Cl
4	1591 (2CO), 1665 (NHCO), 1727	55.65	4.28	8.89	9.85	_
	(COOH), 3069-3450 (NH,OH)	55.45	4.15	8.65	9.50	
7a	1589 (2CO), 1628 (NHCO), 1665	62.54	3.82	8.39	6.40	_
	(COOH), 3061-3400 (NH,OH)	62.17	3.25	8.11	6.23	_
7b	1590 (2CO), 1630 (NHCO), 1667	62.32	3.82	8.39	6.40	_
	(COOH), 3063-3403 (NH,OH)	62.10	3.34	8.15	6.25	_
7c	1593 (2CO), 1632 (NHCO), 1670	58.58	3.80	9.32	7.12	_
	(COOH), 3063–3405 (NH,OH)	58.18	3.26	9.10	6.85	—
7d	1559 (2CO), 1638 (NHCO), 1673	60.29	4.64	11.72	6.71	—
	(COOH), 3065–3410 (NH,OH)	59.99	4.28	11.22	6.11	—
8a	1590 (2CO), 1631 (NHCO), 1667	58.28	3.49	7.28	5.56	6.14
	(COOH), 2921 (CHCl), 3066–3450 (NH,OH)	57.98	3.15	7.05	5.35	5.74
8b	1591 (2CO), 1632 (NHCO), 1668	58.28	3.49	7.28	5.56	6.14
	(COOH), 2922 (CHCl), 3066–3450 (NH,OH)	58.08	3.15	7.15	5.40	5.74
8c	1592 (2CO), 1632 (NHCO), 1670	54.38	3.23	7.93	6.05	6.69
	(COOH), 2923 (CHCl), 3067–3452 (NH,OH)	54.17	2.85	7.55	5.95	6.50
8d	1594 (2CO), 1634 (NHCO), 1670	56.36	4.18	10.11	5.79	6.40
	(COOH), 2925 (CHCl), 3070–3455 (NH,OH)	55.97	3.99	9.98	5.56	5.99
9a	1590 (2CO),1630 (NHCO), 1667	58.48	3.68	7.31	11.15	—
	(COOH), 3063–3445(NH,OH)	57.99	3.38	6.92	10.75	—
9b	1590 (2CO),1631 (NHCO), 1668	58.48	3.68	7.31	11.15	—
	(COOH), 3063–3445(NH,OH)	57.99	3.38	6.92	10.75	—
9c	1591 (2CO),1632 (NHCO), 1669	54.90	3.65	8.00	12.21	—
	(COOH), 3064–3447(NH,OH)	54.69	3.45	7.74	11.15	
9d	1592 (2CO),1632 (NHCO), 1670	56.56	4.38	10.15	11.61	—
	(COOH), 3065–3450(NH,OH)	56.24	3.99	9.85	11.41	—
10a	1588 (2CO),1662 (NHCO), 1723	63.64	4.18	6.45	7.39	—
_	(COOH), 3064–3440 (NH,OH)	63.38	3.98	6.36	7.16	—
10b	1588 (2CO),1661 (NHCO), 1723	61.38	4.03	6.22	7.12	—
	(COOH), 3060–3440 (NH,OH)	61.08	3.94	5.85	6.86	_
10c	1589 (2CO),1663 (NHCO), 1725	59.66	3.58	8.77	6.69	
	(COOH), 3066–3442 (NH,OH)	59.45	3.26	8.52	6.39	
11a	1590 (2CO), 1685 (COCH3), 1730	61.16	4.52	11.41	6.53	_
111	(COOH), 3060–3430 (NH,OH)	59.96	4.22	11.09	6.25	_
110	1591 (2CO), 1686 (COCH3), 1731	59.23	4.37	11.05	6.32	_
	(COOH), 3063–3432 (NH,OH)	58.94	4.07	10.85	5.96	_
11c	1592 (2CO), 1688 (COCH3), 1733	56.08	3.95	13.08	5.99	_
10-	(COOH), 3003–3439 (NH,OH) 1590 (900), 1797 (000H), 2055	00.00 CC 45	3.73	12.70	0.79 C 10	_
12a	1589 (200), 1727 (000п), 3055– 2460 (NH OH)	00.40 66 15	4.01	10.69	6.1Z	_
19h	3400 (NH, OH) 1590 (200) 1799 (COOH) 2050	64 40	4.01	10.44	5.00	_
120	1589 (200), 1728 (000п), 3050- 2460 (NH OH)	04.49 64.91	4.48	10.37	0.94 5.65	_
190	3400 (NH, OH) 1500 (200) 1720 (COOH) 2056	04.21 61.90	4.13 1 00	10.03	0.00 5.69	_
140	2465 (NH OH)	60.96	4.00 3.79	12.31 19.11	5.03 5.39	_
190	1588 (9CO) 1791 (COOH) 2060	61 15	J.10 196	0.26	0.00 7 14	
10a	3465 (NH OH)	60.82	4.40 2.05	9.50 9.05	6.81	
	0100 (111,011)	00.00	0.00	0.00	0.04	

TABLE III Selected IR Spectra, Elemental Analysis for the New Compounds Listed in Table I

Comn		Calculated % Found %				
no.	IR $\upsilon_{\rm max}{\rm Cm}^{-1}$	С	Η	Ν	S	Cl
13b	1589 (2CO), 1723 (COOH), 3063-	59.40	3.67	11.34	6.89	_
	3465 (NH,OH)	59.26	3.35	11.14	6.66	_
13c	1589 (2CO), 1725 (COOH), 3060-	55.91	3.67	11.34	6.49	
	3460 (NH,OH)	55.75	3.36	10.99	6.25	
14a	1592 (2CO),1667 (NHCO), 1732	60.55	4.23	11.77	6.73	—
	(COOH), 3065–3435 (NH,OH)	60.32	3.84	11.52	6.44	
14b	1591 (2CO),1666 (NHCO), 1731	58.58	4.10	11.39	6.51	_
	(COOH), 3064–3435 (NH,OH)	58.24	3.78	11.16	6.25	
14c	1593 (2CO),1668 (NHCO), 1733	55.32	3.67	13.44	6.15	_
	(COOH), 3067–3437 (NH,OH)	54.98	3.35	13.24	5.75	_
15a	1589 (2CO),1245 (CS), 1729	58.70	4.10	11.41	13.06	_
	(COOH), 3061–3451 (NH,OH)	58.46	3.83	11.24	12.96	_
15b	1589 (2CO),1244 (CS), 1730	56.85	3.98	11.05	12.64	_
	(COOH), 3062-3453 (NH,OH)	56.65	3.58	10.85	12.34	_
15c	1590 (2CO),1242 (CS), 1731	53.67	3.57	13.04	11.94	_
	(COOH), 3063-3454 (NH,OH)	53.47	3.27	12.82	11.76	_
16a	1589 (2CO),1443(CN), 1667	62.75	4.55	9.98	7.61	_
	(COOH), 3063-3405 (NH,OH)	62.32	4.36	9.65	7.36	_
16b	1588 (2CO),1445(CN), 1669	58.06	3.99	9.23	7.04	7.79
	(COOH), 3063-3410 (NH,OH)	57.87	3.74	8.99	6.82	7.24
16c	1592 (2CO),1446(CN), 1670	56.69	3.89	12.56	6.88	_
	(COOH), 3065-3415 (NH,OH)	56.48	3.63	11.89	6.45	_
17a	1591 (2CO), 1668 (COOH), 1720	57.99	4.06	8.45	6.44	7.14
	(CO), 2923 (CHCl), 3066–3450 (NH,OH)	57.76	3.79	8.16	6.25	6.95
17b	1593 (2CO), 1670 (COOH), 1722	54.28	3.61	7.91	6.04	13.35
	(CO), 2925 (CHCl), 3068–3455 (NH,OH)	53.97	3.35	7.63	5.85	13.15
17c	1594 (2CO), 1671 (COOH), 1725	53.18	3.53	10.34	5.91	6.54
	(CO), 2925 (CHCl), 3068–3460 (NH,OH)	52.88	3.35	10.15	5.71	6.24
18a	1589 (2CO), 1676 (COOH,CO),	58.22	4.28	8.49	12.95	_
	3060–3443 (NH,OH)	57.88	3.96	8.29	12.66	_
18b	1592 (2CO), 1678 (COOH,CO),	54.49	3.81	7.94	12.12	6.70
	3065–3445 (NH,OH)	54.17	3.61	7.65	11.82	6.36
18c	1594 (2CO), 1679 (COOH,CO),	53.37	3.55	10.37	11.87	_
	3066–3446 (NH,OH)	53.16	3.25	10.16	11.58	_

TABLE III Selected IR Spectra, Elemental Analysis for New Compounds Listed in Table I (*Continued*)

#### Synthesis of New Thiazolidinone Derivatives 9a-d

A solution of **7a–d** (5.01 g, 0.01 mol) and thioglycolic acid (0.92 g, 0.01 mol) in a mixture of ethanol and dimethylformamide in the presence of a few drops of piperidine was refluxed for 8–10 h. The filtrate was evaporated under reduced pressure, poured into ice water where the product was separated, filtered, washed several times with water, and crystallized from the proper solvent to give **9a–d** (c.f., Table I).

# Synthesis of 2-Arylidenoamino-2-methyl, 2-H-naphtho[2, 3-b][1, 4]thiazine-5, 10-dione, 3-carboxylic Acid Derivatives 10a–c

A solution of 4 (3.46 g, 0.01 mol) and aromatic aldehydes (1.06 g, 1.22 g, 1.51 g, 0.01 mol) in a mixture of ethanol and dimethylformamide in the presence of two drops of piperidine as catalyst was refluxed for 7–8 h. The mixture was concentrated and then left to cool and pour onto ice water acidified by concentrated hydrochloric acid. The solid product so formed was collected by filtration and crystallized from the proper solvent to give **10a–c** (c.f., Table I).

#### Synthesis of Acetyl Pyrazolino Derivatives 11a-c

To a solution of arylidino **10a–c** (4.34 g, 4.50 g, 4.79 g, 0.01 mol) in a mixture of ethanol and dimethylformamide as solvent, hydrazine hydrate (0.5 g, 0.01) was added followed by glacial acetic acid (10 ml). The reaction mixture was refluxed for 10–12 h. The reaction mixture was concentrated and filtered. These substances were triturated with water and the precipitates were separated, filtered, washed several times with water, dried, collected, and crystallized from the proper solvent to give **11a–c** (c.f., Table I).

#### Synthesis of 3-aryl-4-phenylpyrazolino Derivatives 12a-c

To a solution of arylidino **10a–c** (4.34 g, 4.50 g, 4.79 g, 0.01 mol) in a mixture of ethanol and dimethylformamide as solvent, added phenyl hydrazine (1.08 g, 0.01 mol) in the presence of 0.5 ml of piperidine as catalyst. The reaction mixture was refluxed for 9–11 h. The reaction mixture was concentrated, cooled, poured onto water acidified by concentrated hydrochloric acid. The solid substances were collected by filtration, washed with water for several times, and crystallized from the proper solvent to give **12a–c** (c.f., Table I).

#### Synthesis of 3-arylisoxazolo Derivatives 13a-c

A solution of arylidino **10a–c** (4.34 g, 4.50 g, 4.79 g, 0.01 mol) was refluxed with hydroxylamine hydrochloride (0.69 g, 0.01 mol) in the presence of sodium hydroxide as catalyst and a mixture of ethanol and dimethylformamide as solvent for 6–8 h. The reaction mixture was filtered hot, the filtrate was concentrated and poured onto ice water, where the products were separated, filtered, washed several times with water, and crystallized from the proper solvent to give **13a–c** (c.f., Table I).

#### Synthesis of 3-arylpyrimidino Derivatives 14a-c

A solution of arylidino **10a–c** (4.34 g, 4.50 g, 4.79 g, 0.01 mol) was refluxed with urea (0.6 g, 0.01 mol) in the presence of sodium hydroxide as catalyst and a mixture of ethanol and dimethylformamide as solvent for 8–10 h. The reaction mixture was filtered from unreacted materials, and the filtrate was concentrated and poured on to ice water, and the products were separated, filtered, washed several times with water, and crystallized from the proper solvent to give **14a–c** (c.f., Table I).

#### Synthesis of 3-arylthiopyrimidino Derivatives 15a-c

A solution of arylidino **10a–c** (4.34 g, 4.50 g, 4.79 g, 0.01 mol) was refluxed with thiourea (0.67 g, 0.01 mol) in the presence of hydrochloric acid as catalyst and a mixture of ethanol and dimethylformamide as solvent for 8–10 h. The reaction mixture was filtered hot, cooled, the filtrate was concentrated, poured onto ice/water, and the products were separated, filtered, washed several times with water, and crystallized from the proper solvent to give **15a–c** (c.f., Table I).

#### Synthesis of New Schiff Base Derivatives 16a-c

Compound 4 (3.46 g, 0.01 mol) and aromatic amine (0.93 g, 1.27 g, 1.38 g, 0.01 mol) in an equimolar ratio were dissolved in ethanol and dimethylformamide and 0.5 ml of piperidine as catalyst was added. The mixture was refluxed about 10 h. The reaction mixture was allowed to cool at room temperature, then filtered, washed several times with water, dried, collected, and crystallized from the proper solvent to give **16a–c** (c.f., Table I).

#### Synthesis of New $\beta$ -Lactam Derivatives 17a–c

A solution of (4.21 g, 4.55 g, 4.66 g, 0.01 mol) of chloroacetylchloride (1.13 g, 0.01 mol) in a mixture of ethanol and dimethylformamide in the presence of a few drops of piperidine was refluxed for 12–14 h. The filtrate was evaporated under reduced pressure, poured into ice/water where the product was separated, filtered, washed several times with water and crystallized from the proper solvent to give **17a–c** (c.f., Table I).

#### Synthesis of New Thiazolidinone Derivatives 18a-c

A solution of **16a–c** (4.21 g, 4.55 g, 4.66 g, 0.01 mol) and thioglycolic acid (0.92 g, 0.01 mol) in a mixture of ethanol and dimethylformamide

in the presence of a few drops of piperidine was refluxed for 11-13 h. The filtrate was evaporated under reduced pressure, poured into ice/water where the product was separated, filtered, washed several times with water, and crystallized from the proper solvent to give **18a-c** (c.f. Table I).

#### REFERENCES

- J. B. Patrick, R. P. Williams, W. E. Meyer, W. Fulmor, D. B. Cosulich, R. W. Broschard, and J. S. Webb, J. Amer. Chem. Soc., 86, 1889 (1964).
- [2] W. Durckheimer, J. Blumhach, R. Lattrll, and K. H. Scheunemann, Angew. Chem., 97, 183 (1985).
- [3] S. R. Woulfe and M. Miller, Tetrahedron Lett., 25, 3293 (1984).
- [4] S. R. Woulfe and M. Miller, J. Med. Chem., 28, 1447 (1985).
- [5] S. R. Woulfe and M. Miller, J. Org. Chem., 51, 3133 (1986).
- [6] H. Breure, H. Straub, U. Treuner, J. M. Drooard, H. Hoehnh, and K. J. Lindler, Antibiot., 83, 813 (1985).
- [7] M. J. Miller, A. Biswas, and M. A. Krook, Tetrahedron, 39, 2571 (1983).
- [8] W. O. Foye, Principles of Medicinal Chemistry (Lea and Febiger, Philadephia, 1981), 2nd ed.
- [9] B. Pirotte, J. Delarge, J. Coyette, and J. M. J. Frere, Antibiot., 44, 844 (1991).
- [10] M. Aoki, T. Ohtsuka, Y. Itezono, K. Yokose, K. Furihata, and H. Seto, *Tetrahedron Lett.*, 32, 217 (1991).
- [11] X. F. Ren, M. I. Konaklieva, H. Shi, Dickeys, D. V. Lim, J. Gonzalez, E. Turos, J. Org. Chem., 63, 8898 (1998).
- [12] A. K. Khalafallah, R. M. Abd El-Aal, and N. A. A. El Kanzi, J. Chin. Chem. Soc., 49, 387–396 (2002).
- [13] A. K. Khalafallah, R. M. Abd El-Aal, and N. A. A. El Kanzi, J. Chin. Chem. Soc., 51, 769–774 (2004).
- [14] A. K. Khalafallah, R. M. Abd El-Aal, and N. A. A. El Kanzi, J. Heterrocyclic. Comm., 8(4) (2002).