of the crude product from Me₂CO-hexane. Recrystallization from EtOAc-hexane gave the analytical specimen: mp 209-212°; $[\alpha]^{25}D - 78.9^{\circ}$; ir (CHCl₃) 3620 and 1720 cm⁻¹. Anal. (C₂₀H₂₉ClO₂) C, H, Cl.

7-Chloro- 5α , 6α -epoxy- 3β , 17β -dihydroxypregn-7-en-20-one 3, 17-Diacetate (11a). A solution of 2a (0.9 g, 2 mmol) and *m*-chloroperbenzoic acid (0.42 g) in CH₂Cl₂ (10 ml) was stirred at room temperature for 30 min, diluted with CH₂Cl₂, and washed in turn with 5% NaHSO₃ solution, 5% NaHCO₃ solution, and H₂O. The dried (Na₂SO₄) organic layer was evaporated *in vacuo* and crystallization of the resulting solid from CH₂Cl₂-Et₂O gave 0.68 g (72%) of 11a, mp 224-225°. Two additional crystallizations from the same solvent system afforded the analytical sample: mp 226-227°; $[\alpha]D - 169.2°$; nmr (CDCl₃) δ 3.20 (s, 1 H, C_eH). Anal. (C₂₅H₃₃ClO₆) C, H, Cl.

6β,7-Dichloro-3β,5α,17α-trihydroxypregn-7-en-20-one 3,17-Diacetate (12). A suspension of 11a (0.9 g, 1.93 mmol) in CCl₄ (20 ml) was cooled to 0°. HCl (g) was bubbled through the stirred mixture until the solid dissolved (~4 min). The solvent was removed at low temperature (<15°) under reduced pressure and the resulting solid was crystallized from Et₂O-hexane to give 0.629 g (65%) of 12, mp 197-198°. Recrystallization from the same solvent system gave the analytical sample: mp 198-199°; [α]D -124.0°; ir (CHCl₃) 3700 (sharp), 3400 (br), and 1735 cm⁻¹; nmr (CDCl₃) δ 4.33 (s, 1 H, C₆H) and 1.23 (s, 3 H, C₁₉H₃). Anal. (C₂₅H₃₄Cl₂O₆) C, H, Cl.

6α,7-Dichloro-3β,5α,17α-trihydroxypregn-7-en-20-one 3,17-Diacetate (13). A solution of 11a (1.0 g, 2.15 mmol) in CH₂Cl₂ (20 ml) was cooled to 0°. HCl (g) was bubbled through the solution for 4 min and the solvent was then removed *in vacuo*. The residue was crystallized from CH₂Cl₂-Et₂O to furnish (0.41 g, 37%) of 13, mp 209-210° (mmp with 12, 183-185°). A second crop (0.1 g, 9%, mp 210-211°) was obtained from the mother liquors. Recrystallization from the same solvent mixture afforded the analytical specimen: mp 211-212°; $[α]^{25}D - 104.5°$; ir (CHCl₃) 3650 and 1735 cm⁻¹; nmr (CDCl₃) δ 4.57 (t, 1 H, C₆H) and 1.01 (s, 3 H, C₁₉H₃). *Anal.* (C₂₃H₃₄Cl₂O₆) C, H, Cl.

7-Chloro-5 α , 6 α -epoxy-3 β , 17 α -dihydroxypregn-7-en-20-one 17-Acetate (11b). To a solution of 2b (2.9 g, 7.2 mmol) in CH₂Cl₂ (30 ml) was added *m*-chloroperbenzoic acid (1.8 g). The solution was allowed to stand at room temperature for 1 hr and was worked up as before. The crude product was crystallized from CH₂Cl₂-Et₂O to give 1.9 g (63%) of 14. A second crop (0.4 g, 13%) was recovered from the mother liquors. Recrystallization from the same solvents furnished the analytical sample: mp 213° dec; [α]²⁵D -179.5°; ir $(CHCl_{3})$ 3700, 1730, and 1710 cm $^{-1}$; nmr $(CDCl_{3})$ δ 3.20 (s, 1 H, C_6H). Anal. $(C_{23}H_{31}ClO_{5})$ C, H, Cl.

Acknowledgment. We wish to thank the following members of our physical chemistry department: Dr. V. Toome, Mr. S. Traiman, and Dr. T. Williams for the ultraviolet, infrared, and nmr spectra, respectively. Thanks are also due Dr. F. Scheidl for the microanalyses.

References

- (1) R. E. Schaub and M. J. Weiss, J. Org. Chem., 26, 3915 (1961).
- (2) J. A. Campbell and J. Babcock, J. Amer. Chem. Soc., 81, 4069 (1959).
- (3) J. A. Campbell, S. C. Lyster, G. W. Duncan, and J. C. Babcock, *Steroids*, 1, 317 (1963).
- (4) P. Wieland and G. Anner, Helv. Chim. Acta, 50, 1453 (1967).
- (5) P. A. Diassi, S. D. Levine, and R. M. Palmere, J. Med. Chem., 10, 551 (1967).
- (6) R. Deghenghi and R. Gaudry, Can. J. Chem., 40, 818 (1962).
- (7) G. W. Moersch and W. A. Neuklis, *ibid.*, 41, 1627 (1963).
- (8) C. W. Marshall, R. E. Ray, I. Laos, and B. Riegel, J. Amer. Chem. Soc., 79, 6303 (1957).
- (9) K. Bowden, I. Heilbron, E. R. H. Jones, and B. Weedon, J. Chem. Soc., 39 (1946).
- (10) G. I. Poos, G. E. Arth, R. E. Beyler, and L. H. Sarett, J. Amer. Chem. Soc., 75, 422 (1953).
- (11) K. E. Pfitzner and J. G. Moffatt, *ibid.*, 87, 5661 (1965).
- (12) J. D. Albright and L. Goldman, *ibid.*, **89**, 2416 (1967).
- (13) J. Eastman and R. Teranishi, "Organic Syntheses," Collect. Vol. IV, Wiley, New York, N. Y., 1963, p 192.
- (14) K. Yasuda, Chem. Pharm. Bull., 11, 1167 (1963).
- (15) C. W. Marshall, U. S. Patent 3,227,124 (1966).
- (16) C. W. Greenhalgh, H. B. Henbest, and E. R. H. Jones, J. Chem. Soc., 2375 (1952).
- (17) A. Windhaus and A. Luttringhaus, Justus Liebigs Ann. Chem., 481, 119 (1930).
- (18) P. Mayor and C. D. Meakins, J. Chem. Soc., 2792 (1960).
- (19) R. A. LeMahieu, A. Boris, M. Carson, and R. W. Kierstead,
- J. Med. Chem., 14, 629 (1971).
- (20) A. Boris, Steroids, 11, 681 (1968)
- (21) L. G. Hershberger, E. G. Shipley, and R. K. Meyer, Proc. Soc. Exp. Biol. Med., 83, 175 (1953).

Antimalarials. 4. 4-Pyridinemethanols with Styryl and Benzoyl Substituents[†]

M. P. LaMontagne

Ash Stevens Inc., Detroit, Michigan 48202. Received July 13, 1972

A series of seven α -alkylaminomethyl-4-pyridinemethanols containing styryl substituents in the 2 and/or 6 positions of the pyridine ring has been synthesized. One compound was curative against *Plasmodium* berghei in mice at a dosage of 20 mg/kg, three were curative at 40 mg/kg, two were curative at 80 mg/kg, and one was inactive through 160 mg/kg. In addition, one compound bearing a 4-trifluoromethylbenzoyl substituent in the 2 position of the pyridine ring was prepared. It was curative at 80 mg/kg and showed marginal activity at 40 mg/kg.

In the two preceding papers^{1,2} in this series, we reported the preparation and antimalarial activity against *Plasmodium berghei* in mice[‡] of a series of α -alkylaminomethyl-2,6bis(phenyl)-4-pyridinemethanols bearing Cl, Br, F, OCH₃, and CF₃ substituents on the phenyl rings. The very encouraging antimalarial activity demonstrated by several compounds in these series prompted us to investigate the effect of replacing one or both of the phenyl substituents with styryl and benzoyl groups.

Chemistry. The four structural types of compounds described in the present work are shown in Chart I. Compounds of type I were synthesized by the procedure shown in Scheme I.

Ethyl 2,6-dimethylisonicotinate was prepared from 4cyano-2,6-lutidine⁴ via hydrolysis and esterification under usual conditions. Condensation of the ester with excess aldehyde afforded the requisite 2,6-bis(styryl)isonicotinic acid esters in approximately 50% yield. Hydrolysis of the esters to the isonicotinic acid, followed by introduction of the amino alcohol side chain by the procedure developed

[†]This work was supported by the U. S. Army Medical Research and Development Command under Contract No. DADA17-69-C-9065. This is Contribution No. 1087 from the Army Research Program on Malaria.

[‡]The antimalarial tests were performed by Dr. Leo Rane of the University of Miami (see ref 3). See footnote *a*, Table III. Testing results were supplied through the courtesy of Drs. Thomas R. Sweeney and Richard E. Strube of the Walter Reed Army Institute of Research.

Scheme I



by Lutz and coworkers,⁵ afforded the desired bis(styryl)-4pyridinemethanols **3a-d**.

The two examples of type II were prepared as shown in Scheme II. The 2-methyl-6-(4-trifluoromethylphenyl)isonicotinic acid (1c) was prepared *via* the modified Zecher-Krohnke ring-closure reaction previously described.¹ The remainder of the sequence was identical with that used to prepare compounds 3a-d.

The analog of type III was prepared as shown in Scheme III. The starting 2-picoline was also prepared by the modified Zecher-Krohnke cyclization procedure.¹ Conversion to the 2-pyridylcarbinol acetate was by the procedure of Boekelheide and Linn.⁶ Hydrolysis with a catalytic amount of sodium ethoxide, followed by oxidation with selenium dioxide, afforded the required aldehyde. The styryl function was introduced in good yield by the reaction of the 2pyridinecarboxaldehyde with 4-chlorophenyltriphenylphosphonium methylide. Hydrolysis to the isonicotinic acid,

Chart I



followed by the introduction of the side chain by the usual sequence, afforded the target compound. The benzoyl analog synthesized (3h) was prepared as shown in Scheme IV (see Tables I and II)

Biological Activity. Antimalarial activity data against *P. berghei* in mice, as measured by the Rane test, ‡,3 are presented in Table III. All compounds prepared were curative at a dosage of 80 mg/kg or less, except **3d** which was inactive through 160 mg/kg (the highest dose level tested); three compounds were curative at 40 mg/kg and one was active at 20 mg/kg.

In preceding papers, we have reported results for the substituted 2,6-bis(phenyl)-4-pyridinemethanols.^{1,2} If we compare the 2,6-bis(trifluoromethylphenyl) analogs² with the present styryl and benzoyl counterparts, we observe a decrease in activity; on the other hand, α -(di-*n*-butylaminomethyl)-2,6-bis(4-chlorostyryl)-4-pyridinemethanol is one dosage level more active than the corresponding 2,6-bis(4chlorophenyl) compound,¹ but the data are too limited to generalize that Cl substituents are more effective than CF₃ substituents in the styryl series. Again, while examples are





limited, the effects of the amino alcohol side-chain configuration upon activity differ with styryl vs. phenyl substituents. Particularly striking is that α -(*n*-butylaminomethyl)-2,6-bis(4-trifluoromethylstyryl)-4-pyridinemethanol (3d) is inactive through 160 mg/kg, whereas the di-*n*butyl analog 3b is curative at 80 mg/kg. Compare also 3e and 3f. None of the compounds were toxic to the host at the dosages tested.

Experimental Section

Melting points were taken in open capillary tubes using a Thomas-Hoover melting point apparatus and are uncorrected.

Scheme III



Elemental analyses were performed by Midwest Microlab, Ltd., Indianapolis, Ind. Analyses indicated by element symbols agree with calculated values within $\pm 0.4\%$.

Preparation of Styryl-Containing Isonicotinic Acids. Ethyl 2,6-Dimethylisonicotinate. A solution of 4-cyano-2,6-lutidine⁴ (15 g, 0.115 mol) in H₂O (300 ml) containing NaOH (50 g) was heated at reflux until NH₃ evolution ceased. The solution was acidified with HCl and concentrated to dryness at reduced pressure. The residue was dissolved in EtOH (500 ml) containing H₂SO₄ (20 g) and the solution was heated at reflux 6 hr. The reaction mixture was carefully poured into H₂O containing NaHCO₃ (80 g) and the product was extracted with Et₂O. The Et₂O was dried (MgSO₄) and concentrated and the residue distilled to yield the ester (13.5 g, 75%), bp 95-97° (2 mm) [lit.⁷ bp 115-118° (5 mm)].

Ethyl 2,6-Di-*p*-chlorostyrylisonicotinate. A solution of ethyl 2,6-dimethylisonicotinate (5.4 g, 0.03 mol) and *p*-chlorobenzaldehyde (15.1 g, 0.12 mol) in Ac₂O (25 g) containing ZnCl₂ (0.2 g) was heated to reflux for 19 hr. The HOAc and Ac₂O were distilled until the pot temperature reached 215°. The viscous oil crystallized upon cooling. The solid was triturated with petroleum ether (two times) and with dilute K₂CO₃. The solid was crystallized from C₆H₆ to afford the product (7.0 g, 55%), mp 158-160°

Anal. $(C_{24}H_{19}Cl_2NO_2) C, H, Cl, N.$

2,6-Di-*p*-chlorostyrylisonicotinic Acid (1a). The above ester (6.5 g, 15 mmol) was heated at reflux in 5% ethanolic KOH (150 ml) for 2 hr. Water was added and the precipitate dissolved. The solution was acidified to pH 2 with dilute HCl. The solid was filtered and recrystallized from HOAc to yield the acid (5.5 g, 92%), mp $283-286^{\circ}$

Anal. (C₂₂H₁₅Cl₂NO₂) C, H, Cl.

The same procedure was used to prepare 1b.

2-Methyl-6-(4-trifluoromethylphenyl)isonicotinic Acid (1c). A solution of 3-(4-trifluoromethylbenzoyl)acrylic acid² (50 g, 0.2 mol), acetylmethylpyridinium bromide (44 g, 0.2 mol), and NH₄OAc (165 g) in MeOH was heated at reflux 6 hr. MeOH was removed and the solid was slurried in dilute AcOH. The mixture was extracted with CHCl₃ (two times), dried (MgSO₄), and concentrated. The crude



Table I. Isonicotinic Acids and Bromo Ketone Intermediates

$R_3 \longrightarrow R_1$									
Compd	R ₁	R ₂	R ₃	Mp, °C (solvent)	Yield, %	Formula	Analyses		
1a	4-CIC,H_CH=CH-	СООН	4-ClC ₆ H ₄ CH=CH-	283-286 (HOAc)	50 ^a	C ₂₂ H ₁₅ Cl ₂ NO ₂	C, H, Cl		
1b	4-CF ₃ C ₆ H ₄ CH=CH-	COOH	4-CF ₃ C ₄ H ₄ CH=CH-	282-285 (HOAc)	47 ^a	C ₂₄ H ₁₅ F ₆ NO ₂	C, H, F		
1c	CH,	COOH	4-CF ₃ C ₆ H ₄ -	$162-164 (C_6H_6)$	74 ^b	$C_{14}H_{10}F_{3}NO_{2}$	C, H, N, F		
1d	4-CF ₃ C ₆ H₄CH=CH−	COOH	4-CF ₃ C ₄ H ₄ -	234-237 (C,H,CH,)	38 <i>°</i>	C ₂₂ H ₁₃ F ₆ NO ₂	C, H, F, N		
1e	4-ClČ₅H₄CH=CH-	COOH	CF,	268-270 (HOAc)	84^d	C ₁ , H ₂ ClF ₃ NO ₂	C, H, Cl		
1f	4-CF ₃ C ₄ H ₄ C(O)-	СООН	4-CF ₃ C ₆ H ₄ -	215-217 (C ₆ H ₆ CH ₃)	64 ^d	C ₂₁ H ₁₁ F ₆ NO ₃	C, H, F, N		
2a	4-ClČ ₆ H ₄ CH=CH-	C(O)CH ₂ Br	4-CIČ,H₄CH=CH-	183-185 (CHCl, - <i>i</i> -PrOH)	62 ^e	C ₂₃ H ₁₆ BrCl ₂ NO	N		
2 b	4-CF ₃ C ₆ H ₄ CH=CH-	C(O)CH ₂ Br	4-CF ₃ C ₆ H ₄ CH=CH-	f					
2c	4-CF ₃ C ₆ H ₄ CH=CH-	C(O)CH,Br	4-CF ₃ C ₄ H ₄ -	132-134 (<i>i</i> -PrOH)	90 ^e	C ₂₃ H ₁₄ BrF ₆ NO	C, H, F, N		
2d	4-CIC ₆ H ₄ CH=CH−	C(O)CH ₂ Br	CF,	115-117 (<i>i</i> -PrOH-H ₂ O)	64 ^e	C ₁₆ H ₁₀ BrClF ₃ NO	C, H, N		
2e	4-CF ₃ C ₆ H ₄ -	C(O)CH ₂ Br	$4-CF_{3}C_{6}H_{4}C(O)-$	140-142 (<i>i</i> -PrOH)	79 ^e	C ₂₂ H ₁₂ BrF ₆ NO ₂	C, H, Br		

R₂

^{*a*}From ethyl 2,6-dimethylisonicotinate. ^{*b*}From cyclization reaction. ^{*c*}From 1c. ^{*d*}From pyridinecarboxaldehyde. ^{*e*}From isonicotinic acid. ^{*f*}Isolated as a mixture of base and HBr salt.

Table II. 4-Pyridinemethanols

$ \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array}\\ \end{array}\\ \end{array}\\ \end{array}\\ \end{array}\\ \end{array}\\ \end{array} \end{array} \begin{array}{c} \begin{array}{c} \end{array}\\ \end{array}\\ \end{array} \begin{array}{c} \end{array} \end{array} \begin{array}{c} \end{array} \end{array} \begin{array}{c} \end{array} $								
Compd	R ₁	R ₂	R ₃	R4	Mp, °C (solvent)	Yield, ^a %	Formula	Analyses
	4-CIC,H,CH=CH-	4-CIC_H_CH=CH-	1-Bu	1-Bu	195-197 (i-PrOH)	28	C ₃₁ H ₃₇ Cl ₃ N ₂ O	C, H, Cl, N
3b	4-CF ₄ C ₄ H ₄ CH=CH-	4-CF ₄ C ₄ H ₄ CH=CH-	1-Bu	1-Bu	188-189 (CH ₃ CN)	45	C ₃₃ H ₃₇ ClF ₆ N ₂ O	C, H, Cl, F, N
3c	4-CF ₄ C ₄ H ₄ CH=CH-	4-CF ₄ C ₆ H ₄ CH=CH-	Н	4-Heptyl	110–112 ^b (<i>i</i> -PrOH)	34	$C_{32}H_{34}F_{6}N_{2}O$	C, H, F, N
3d	4-CF ₄ C ₆ H ₄ CH=CH-	4-CF ₃ C ₆ H ₄ CH=CH-	Н	1-Bu	137–139 ^b (<i>i</i> -PrOH)	51	$C_{29}H_{28}F_{6}N_{2}O$	C, H, F, N
3e	4-CF C, H, CH=CH-	4-CF ₃ C ₆ H ₄ -	1-Bu	1-Bu	194-196 (CH ₃ CN)	42	C ₃₁ H ₃₅ ClF ₆ N ₂ O	C, H, F, N
3f	4-CF ₃ C ₆ H ₄ CH=CH-	$4 - CF_3C_6H_4 -$	Н	1-Bu	111-113 ^b (CH ₃ CN)	56	C ₂₇ H ₂₆ F ₆ N ₂ O	C, H, F, N
3g	4-ClC ₆ H ₄ CH=CH-	CF ₃	1-Bu	1-Bu	$162-163.5 (C_6 H_6)$	44	C ₂₄ H ₃₁ Cl ₃ F ₃ N ₂ O	C, H, N
3ĥ	$4-CF_{3}C_{6}H_{4}C(0)-$	4-CF₃C₅H₄-	1-Bu	1-Bu	90.5-92 (C ₆ H ₅ ČH ₃)	43	C ₃₀ H ₃₃ F ₆ ClN ₂ O ₂	C, H, Cl, N

^aFrom bromo ketone precursor. ^bFree base.

Table III. Antimalarial Activity against P. berghei

	$\Delta MST \text{ or } C^{a,b}$ Dosage, mg/kg						
Compd	10	20	40	80	160		
3a	1.1	4.9	3C	5C	5C		
3b		0.2	2.9	2C	5C		
3c	1.1	3.1	4C	5C	5C		
3d			Inactive				
3e	3.7	13.5	4C	5C	5C		
3f		0.5	13.6	4C	5C		
3g	4.5	2C	4C	5C	5C		
3ĥ		1.3	6.7	3C	5C		

^aThe test method, described in ref 3, is a highly standardized procedure in which the *P. berghei* causes death of control mice at essentially 6 days. An increase in the mean survival time of five mice by more than 2.5 days beyond this time is statistically significant. Mice surviving more than 60 days are regarded as cured (C). A candidate drug is considered active (A) at a given dosage if one or more mice are alive on day 14. ^bAll compounds listed, with the exception of 3d, were nontoxic and curative (5C) through 640 mg/kg. Data for dosages above 160 mg/kg for compound 3d are not available.

product was crystallized from C_6H_6 (500 ml) to yield the acid (42 g, 74%), mp 162-164°.

Anal. (C14H10F3NO2) C, H, F, N.

Methyl 2-Methyl-6-(4-trifluoromethylphenyl)isonicotinate. A solution of the isonicotinic acid (42 g, 0.15 mol) in dry MeOH (600 ml) containing concentrated H₂SO₄ (35 ml) was heated at reflux 17 hr. The solution was concentrated to one-third volume under reduced pressure. H₂O was added and the slurry was extracted with CHCl₃ (two times). The CHCl₃ was dried (Na₂SO₄) and concentrated under reduced pressure. The residue was crystallized from EtOH-H₂O to yield the title ester (33 g, 75%), mp 164-166°.

Anal. (C15H12NF3O2) C, H, N.

2-(4-Trifluoromethylstyryl)-6-(4-trifluoromethylphenyl)isonicotinic Acid (1d). A solution of the above ester (8.85 g, 0.03 mol), 4-trifluoromethylbenzaldehyde (11 g, 0.06 mol), and ZnCl₂ (0.2 g) in Ac₂O (35 ml) was heated at reflux 18 hr. The solvents were distilled to an internal temperature of 220°. The oily residue was placed onto a silica gel column and eluted with C₆H₆-petroleum ether (1:1). The fast-moving material was collected and the solvent was removed under reduced pressure. The material crystallized on standing. Recrystallization from C₆H₆-petroleum ether gave the ester (5.5 g, 41%). The ester was hydrolyzed as described for 1a. Crystallization from toluene gave the acid (5.0 g, 92%), mp 234-237°.

Anal. (C22H13F6NO2) C, H, F, N.

Ethyl 2-Methyl-6-trifluoromethylisonicotinate. A solution of trifluoromethylacylpyridinium bromide (25.5 g, 0.1 mol), ethyl β -acetylacrylate (14.2 g, 0.1 mol), and NH₄OAc (85 g) in EtOH (150 ml) containing HOAc (50 ml) was heated at reflux for 6 hr. The solvent was removed under reduced pressure. H₂O was added and the mixture was extracted with CHCl₃ (three times). The CHCl₃ was washed with H₂O, dried (MgSO₄), and concentrated. The dark residual liquid was distilled *in vacuo* to afford the title ester (6 g, 32%), bp 65-70° (0.5 mm).

Anal. (C10H10F3NO2) C, H.

Ethyl 2-Methyl-6-trifluoromethylisonicotinate N-Oxide. A solution of the ester (19 g, 0.08 mol) in CF₃COOH (150 ml) containing 35% H₂O₂ (36 cc) was refluxed 2.25 hr. The solution was diluted with H₂O and extracted with CHCl₃ (three times). The CHCl₃ was washed with H₂O and aqueous K₂CO₄, dried (MgSO₄), and concentrated. The residual solid was crystallized (petroleum ether) to yield the product (11 g, 57%), mp 71-73°.

Anal. (C10H10F3NO3) C, H, F, N.

4-Carbethoxy-6-trifluoromethyl-2-pyridylcarbinol Acetate. A solution of the above N-oxide (11.8 g, 0.05 mol) in Ac₂O (120 ml) was heated at reflux 65 hr. The Ac₂O was removed under reduced pressure and the residue was distilled *in vacuo* to yield the product (7.5 g, 55%), bp 95-100° (0.25 mm).

Anal. (C12H12F3NO4) C, H, N.

4-Carbethoxy-6-trifluoromethyl-2-pyridylcarbinol. To a solution of the above ester (5.0 g, 0.018 mol) in anhydrous EtOH was added a catalytic amount of NaOEt in EtOH. The solution was stirred 15 min and neutralized with HOAc. The solvent was removed at reduced pressure. The residue was dissolved in Et₂O and washed with H₂O (two times). The Et₂O was dried (MgSO₄) and concentrated to give the product (4.0 g), an oil.

Anal. (C10H10F3NO3) C, H, N.

4-Carbethoxy-6-trifluoromethyl-2-pyridinecarboxaldehyde. A solution of the carbinol (3.6 g, 0.015 mol) and SeO₂ (1.8 g) in dioxane (70 ml) was heated for 17 hr. The selenium was removed *via* filtration (Celite) and the solvent was removed *in vacuo*. The residue was chromatographed over a silica gel column and eluted with C_6H_6 . The product was collected (2.9 g, 81%), an oil.

Anal. $(C_{10}H_8F_3NO_3)C, H.$

4-Chlorobenzyltriphenylphosphonium Chloride. A solution of triphenylphosphine (5 g) and $\alpha_{,p}$ -dichlorotoluene (3 g) in p-xylene (50 ml) was heated at reflux 5 hr. The solution was cooled and filtered to afford the phosphonium salt (3.9 g), mp 285-287°.

2-(4-Chlorostyryl)-4-carbethoxy-6-trifluoromethylpyridine. To a slurry of the above phosphonium salt (4.8 g, 0.01 mol) in anhydrous Et_2O (20 ml) was added *n*-BuLi in hexane (2.56 *M*, 4.6 cc). The orange slurry was stirred 0.5 hr. To this slurry was added 4carbethoxy-6-trifluoromethyl-2-pyridinecarboxaldehyde (2.85 g, 0.011 mol). The resulting slurry was stirred an additional hour. The triphenylphosphine oxide and LiCl were filtered and the filtrate was dried (MgSO₄) and concentrated to afford the product (4.0 g, 99%), an oil. A sample was purified by chromatography over a silica gel column and elution with C₆H₆.

Anal. $(C_{17}H_{13}ClF_3NO_2)C, H, N.$

2-(4-Chlorostyryl)-6-trifluoromethylisonicotinic Acid (1e). A solution of the above ester (4 g, 0.011 mol) in 5% KOH-EtOH (60 ml) was stirred at room temperature for 0.5 hr. H_2O was added until all the salt dissolved. Most of the EtOH was removed under reduced pressure. Dilute aqueous K_2CO_3 (100 ml) was added and the mixture was extracted with Et_2O (two times). The aqueous layer was acidified to pH 2 and extracted with Et_2O (three times) to afford the crude acid which was crystallized from HOAc to yield pure product (3.1 g, 84%), mp 268-270°.

Anal. $(C_{15}H_9ClF_3NO_2)$ C, H, Cl.

Styryl-Containing 4-Pyridinemethanols. The above isonicotinic acids were converted to the amino alcohols *via* the procedure described previously.¹ The analytical data are shown in Table II.

4-Carboxymethyl-6-(4-trifluoromethylphenyl)-2-pyridinecarboxaldehyde. A mixture of methyl 2-methyl-6-(4-trifluoromethylphenyl)isonicotinate (15 g, 0.05 mol) and I_2 (12.58 g) was heated at 125° for 20 min. The solution was diluted with DMSO (30 ml) and added to DMSO (60 ml) preheated to 130°. This solution was heated at 165° (bath) under N₂ for 4 hr and cooled and CHCl₃ was added. The solution was washed successively with H₂O, dilute sodium thiosulfate, and dilute KOH. The organic layer was dried (MgSO₄) and concentrated at reduced pressure. The crude aldehyde was dissolved in C₆H₆ (25 ml) and added to sodium bisulfite solution (75 ml). The addition compound was filtered and stirred with saturated aqueous NaHCO₃ (650 ml) and Et₂O (850 ml). The Et₂O layer was dried and concentrated under reduced pressure to yield the title aldehyde (7.2 g, 48%), mp 118–120°. Crystallization from acetonitrile gave mp 118–120°

Anal. (C15H10F3NO3) C, H, N.

4-(Carboxymethyl)-6-(4-trifluoromethylphenyl)-2-pyridyl-(4trifluoromethylphenyl)carbinol. A slurry of the above aldehyde (6.6 g) in anhydrous Et₂O (150 ml) was treated with a solution of 4trifluoromethylphenylmagnesium bromide (from 5.4 g of 4-bromobenzotrifluoride and 0.58 g of Mg) in Et₂O. The solution was stirred at room temperature for 1 hr. The mixture was poured into dilute aqueous NH₄OAc (150 ml). The Et₂O layer was dried (MgSO₄) and concentrated. The crude carbinol was crystallized from C₆H₆ – petroleum ether to yield the title compound (6.9 g, 73%), mp 132– 134°.

Anal. $(C_{22}H_{15}F_6NO_3)C, H, F, N.$

Methyl 2-(4-Trifluoromethylbenzoyl)-6-(4-trifluoromethylphenyl)isonicotinate. The title compound was prepared according to the procedure described for 4-carbethoxy-6-trifluoromethyl-2pyridinecarboxaldehyde. A sample, crystallized from *i*-PrOH, had mp $167-169^{\circ}$.

Anal. (C22H23F6NO3) C, H, F, N.

2-(4-Trifluoromethylbenzoyl)-6-(4-trifluoromethylphenyl)isonicotinic Acid (1f). A solution of the above ester (5.0 g, 11 mmol) in AcOH (140 ml) and concentrated HCl (27 ml) was refluxed 3.5 hr. Additional HCl (30 ml) was added and the solution was heated an additional 90 min. Filtration gave the title acid (4.4 g, 90%), mp 215-217°. A sample, recrystallized from toluene, had mp 215-217° Anal. (C₂₁H₁₁F₆NO₃) C, H, F, N.

This acid was converted to the bromomethyl ketone 2e as previously described.

2-(4-Trifluoromethylbenzoyl)-6-(4-trifluoromethylphenyl)-4pyridylethylene Oxide. The bromo ketone was reduced with NaBH₄ as previously described.¹ The resulting epoxycarbinol was oxidized with SeO₂ as described for 4-carbethoxy-6-trifluoromethyl-2-pyridinecarboxaldehyde. Crystallization from *i*-PrOH (10 ml) gave the title compound (2.2 g, 67%), mp 104–106°. A sample, recrystallized from *i*-PrOH, had mp 110–112°.

Anal. $(C_{22}H_{13}F_6NO) N$.

The ethylene oxide was converted to the amino alcohol 3h by the procedure previously described.¹

Acknowledgment. The advice and timely suggestions of Drs. T. R. Sweeney and R. E. Strube of the Walter Reed Army Institute of Reserach are gratefully acknowledged.

References

- P. Blumbergs, M. P. LaMontagne, A. Markovac, J. G. Moehring, A. B. Ash, and C. L. Stevens, J. Med. Chem., 15, 808 (1972).
- (2) A. Markovac, M. P. LaMontagne, P. Blumbergs, A. B. Ash, and C. L. Stevens, *ibid.*, 15, 918 (1972).
- (3) T. S. Osdene, P. B. Russell, and L. Rane, ibid., 10, 431 (1967).
- (4) W. E. Feeley and E. M. Beavers, J. Amer. Chem. Soc., 81, 4004 (1959).
- (5) R. E. Lutz, P. S. Bailey, M. T. Clark, J. F. Codington, A. J. Deinet, J. A. Freek, G. H. Harnest, N. H. Leake, T. A. Martin, R. J. Rowlett, Jr. J. M. Salsbury, N. H. Shearer, Jr., J. D. Smith, and J. W. Wilson, III, *ibid.*, 68, 1813 (1946).
- (6) V. Boekelheide and W. J. Linn, ibid., 76, 1286 (1954).
- (7) N. Sperber, M. Sherlock, D. Papa, and D. Kender, *ibid.*, 81, 704 (1959).

Structure and Antischistosomal Activity in the Nitrofuran Series. Requirement for a 5-Nitro-2-furyl-Vinyl Moiety Based on Comparison of 3-(5-Nitro-2-furyl)-Substituted Propionic, Acrylic, and Propiolic Acid Derivatives[†]

P. B. Hulbert, E. Bueding, and C. H. Robinson*

Department of Pharmacology and Experimental Therapeutics, School of Medicine, and Department of Pathobiology. School of Hygiene and Public Health, The Johns Hopkins University, Baltimore, Maryland 21205. Received May 24, 1972

The structural features required for antischistosomal activity in the nitrofuran series have been postulated to include linkage of the nitrofuran *via* a vinyl group to a nitrogen atom of low basicity. This proposal has been examined by the synthesis and testing against *Schistosoma mansoni* of a series of amides of 3-(5-nitro-2-furyl)propionic acid (Ia) and 3-(5-nitro-2-furyl)propiolic acid (IIIa) for comparison with a corresponding series of amides derived from 3-(5-nitro-2-furyl)acrylic acid (IIa). A direct comparison of active amides derived from IIa with exact analogs differing only in the substitution of a C=C or CH₂CH₂ grouping for a *trans*-CH=CH group has been made. The results show that the vinyl bridge is required for manifestation of antischistosomal activity by such nitrofuran derivatives. Differences in activity among the various amides of IIa are attributed to differences in lipophilicity. Additional studies using *Schistosoma japonicum* revealed that whereas two of the most active amides, IId and IIg, effected few parasitological cures at tolerated doses, the nitrofuran derivative VII was highly effective using large but tolerated doses.

Schistosomiasis, a parasitic disease afflicting approximately 200 million of the world's population, remains a great challenge to the medicinal chemist. (For a recent review, see ref 2.) It has been estimated that over a quarter of a million compounds have been tested for antischistosomal activity;³ yet no drug has yet emerged which has gained general acceptance and widespread use.

We have already described⁴ the general structural and conformational features present in those few nitrofurans and nitrothiazoles which demonstrate good antischistosomal activity and indicated that these features are rather specific. While it is well recognized that nitrofurans may be chemotherapeutically useful against a wide spectrum of organisms,⁵ very few of these compounds exhibit antischistosomal activity, suggesting that the structural features necessary for nitrofurans to show this anthelmintic activity are far more precise.

We have reported that *trans*-5-amino-3-[(5-nitro-2-fury])vinyl]-1,2,4-oxadiazole (VII)⁴ and some close analogs⁶

 $[\]dagger$ This investigation was supported in part by Research Grants GM 16492 and AI 08022 from the National Institutes of Health, as well as Research Grant GA BMS 7011 from the Rockefeller Foundation. Presented¹ in part at the Sixth Middle Atlantic Regional Meeting of the American Chemical Society, Baltimore, Md., Feb 1971.