

HETEROCYCLES, Vol. 78, No. 6, 2009, pp. 1569 - 1571. © The Japan Institute of Heterocyclic Chemistry  
Received, 19th January, 2009, Accepted, 19th February, 2009, Published online, 20th February, 2009  
DOI: 10.3987/COM-09-11659

## A FACILE AND CONVENIENT METHOD TO THE ONE-POT SYNTHESIS OF 2-MERCAPTO-4(3*H*)-QUINAZOLINONES


Farzad Nikpour,\* Roya Mozafari, and Touraj Paibast

Department of Chemistry, Faculty of Sciences, University of Kurdistan, P.O. Box 66315-416, Sanandaj, Iran

**Abstract** – A facile and convenient method to the one-pot synthesis of 2-mercapto-4(3*H*)-quinazolinones is described from the reaction of anthranilic acid derivatives with thiourea in PEG.

2-Mercapto-4(3*H*)-quinazolinone derivatives represent one of the most active classes of heterocycles possessing a wide spectrum of industrial<sup>1</sup> and biological activities.<sup>2-5</sup> Due to their diverse and remarkable applications, considerable attention has been focused on their synthesis.<sup>3,6</sup> The simplest and most accessible method has proven to be the one based on the reaction of anthranilic acid or its functional derivatives with a source of NH and thiocarbonyl such as thiocyanate salts and isothiocyanates.<sup>6a, 6c, 6f, 6g</sup> Although some of these synthesis methods are very useful, most of them are associated with one or the other limitations, such as involving two or more steps, use of very toxic chemicals, employment of harsh reaction conditions, low reaction yields and moreover the low availability of starting materials. Certainly, exploitation of simple reagents with good reaction yields and development of convenient methods to synthesize or to modify such compounds is a worthwhile effort in heterocycles synthesis, and could be of interest.

Here, we wish to report a convenient and simple procedure for the one-pot synthesis of 2-mercapto-4(3*H*)-quinazolinones **2** by direct cyclocondensation reaction of anthranilic acid derivatives **1** with thiourea in polyethyleneglycol (PEG). The reaction proceeds in neutral media and without use of any catalyst (Scheme 1).



Scheme 1



dissolves the polar anthranilic acid derivatives and thiourea. Also, it is miscible with water; therefore, it is simple to remove the solvent and the residual thiourea in work-up process by washing of the reaction mixture in water. In similar conditions, the reactions do not proceed in DMF and in EtOH.

In conclusion, the present procedure develops a simple and convenient method for the one-pot synthesis of 2-mercapto-4(3*H*)-quinazolinones in neutral media and without need to use any catalyst. Also, this procedure provides good yields of products with easy work-up of the reaction mixtures.

## ACKNOWLEDGEMENTS

We are thankful to the University of Kurdistan Research Council for the partial support of this work.

## REFERENCES AND NOTES

1. J. Kono and K. Ogi, JP Patent 60,170,843 (*Chem. Abstr.*, 1984, **104**, 119889c).
2. G. Kulcsár, T. Kálai, E. Ösz, C. P. Sár, J. Jekő, B. Sümegi, and K. Hideg, *Arkivoc*, 2003, 121.
3. A. G. A. El-Helbi and M. H. Abdel-Wahab, *Acta Pharm.*, 2003, **53**, 127; and references cited therein.
4. K. Kalcher, J. Lintschinger, and C. Neuhold, *Mikrochimica Acta*, 1993, **112**, 135.
5. C. Y. Lin, A. C. Ho, H. C. Chiang, J. S. Sun, and S. Y. Sheu, *Anticancer Res.*, 2002, **22**, 937.
6. (a) V. K. Tiwari, D. D. Singh, H. A. Hussain, B. B. Mishra, and A. Singh, *Monatsh. Chem.*, 2008, **139**, 43; and references cited therein; (b) N. I. Mukarramov, B. A. Urakov, and K. M. Shakhidoyatov, *Chem. Heterocyclic Compounds*, 2006, **42**, 540; and references cited therein; (c) K. M. Pritchard, J. M. Al-Rawi, and A. B. Hughes, *Synth. Commun.*, 2005, **35**, 1601; (d) V. N. Britsun, A. N. Esipenko, A. A. Kudryavtsev, and M. O. Lozinskii, *Russ. J. Org. Chem.*, 2005, **41**, 1333.; (e) G. Calestani, L. Capella, R. Leardini, M. Minozzi, R. Papa, D. Nanni, and G. Zanardi, *Tetrahedron*, 2001, **57**, 7221; (f) J. W. Chern, F. J. Shish, C. H. Chan, and K. C. Liu, *Heterocycles*, 1987, **26**, 3193; and references cited therein; (g) G. R. Dave, G. S. Mewada, and G. C. Amin, *J. Indian Chem. Soc.*, 1950, **37**, 595.
7. W. P. Flieck, *Chem. Ber.*, 1973, **106**, 3149.
8. S. L. Shapiro, I. M. Rose, and L. Freadman, *J. Am. Chem. Soc.*, 1957, **79**, 281.
9. J. Chen, S. K. Spear, J. G. Huddleston, and R. D. Rogers, *Green Chem.*, 2005, **7**, 64.
10. *General reaction procedure*: In a 25 mL round flask a mixture of anthranilic acids **1a-f** (5 mmol), 0.38 g (5 mmol) of thiourea, and 1 mL PEG was heated at 140-150 °C for the times as indicated in Table 1. The reaction mixture was cooled to rt; then, H<sub>2</sub>O (5 mL) was added and mixed with the contents and decanted (three times). The raw products were washed with 10% aqueous NaHCO<sub>3</sub> to remove the residual of anthranilic acid derivatives. The products collected and dried first in air then in oven (100 °C). They may be recrystallized from EtOH or EtOH/EtOAc, if needed.