


This article was downloaded by: [University of Hong Kong Libraries]
On: 21 February 2013, At: 07:15
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for
authors and subscription information:

<http://www.tandfonline.com/loi/lcyc20>

DIRECT SYNTHESIS OF γ - BUTYROLACTONES VIA γ - PHENYL SUBSTITUTED BUTYRIC ACIDS MEDIATED BENZYL RADICAL CYCLIZATION

N. O. Mahmoodi^a & M. Jazayri^a

^a University of Guilan, P.O. Box 1914, Rasht, Iran
Version of record first published: 02 Aug 2010.

To cite this article: N. O. Mahmoodi & M. Jazayri (2001): DIRECT SYNTHESIS OF γ -
BUTYROLACTONES VIA γ -PHENYL SUBSTITUTED BUTYRIC ACIDS MEDIATED BENZYL
RADICAL CYCLIZATION, *Synthetic Communications: An International Journal for
Rapid Communication of Synthetic Organic Chemistry*, 31:10, 1467-1475

To link to this article: <http://dx.doi.org/10.1081/SCC-100104057>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study
purposes. Any substantial or systematic reproduction, redistribution,
reselling, loan, sub-licensing, systematic supply, or distribution in any form
to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

DIRECT SYNTHESIS OF γ -BUTYROLACTONES VIA γ -PHENYL SUBSTITUTED BUTYRIC ACIDS MEDIATED BENZYL RADICAL CYCLIZATION

N. O. Mahmoodi* and M. Jazayri

Organic Research Laboratory, Department of Chemistry,
University of Guilan, Rasht, P.O. Box 1914, Iran


ABSTRACT

Synthesis of several γ -butyrolactones with aromatic substitution at carbon 5 from comparative γ -aryl acids with 25–85% yield are covered. The straight oxidation in the presence of peroxydisulphate-copper(II)chloride system in aqueous medium was applied. The reaction is highly regioselective and leads exclusively to γ -butyrolactone, through stable benzylic radical intermediate.

Oxidative addition of carboxylic acids and alkenes was used as a mild methodology for formation of γ -butyrolactones.^{1–4} The oxidative systems such as $S_2O_8^{2-}$ - Ag^+ , $S_2O_8^{2-}$ - Ag^+ - Cu ,^{5–7} different oxidants' derivatives of Pb(IV), Co(III), Ag(II), other polyvalent metals,^{8–10} and $KMnO_4$ ¹¹ can interact with carboxylic acids, resulting usually in their decarboxylation and creation of alkyl radicals. Direct oxidative systems such as oxidative lactonization of carboxylic acids in the presence of

* Corresponding author.

peroxydisulphate-containing systems have been published. However, the yields of conversion are either very low or the reaction is not selective, e.g., 3.3–9.5% and 4–10%, respectively, for mixtures that contain 5- and 6-member ring lactones together with some decarboxylated compounds.¹² The objectives of this study are to convert 4-substituted aryl acids in the presence of an oxidative system such as $S_2O_8^{2-}-Cu^{2+}$ from mild to high yields, and to monitor the conversion to exclusively 5-member lactone ring. Recently, we synthesized several mono, and di-substituted γ -butyrolactones with aryl and aliphatic substitution at carbon 5, 3,5 and 3,4 as an essence,¹³ an anti-glaucoma, and an anti-tumor,^{14–17} respectively. The direct oxidation system in the presence of peroxydisulphate-copper chloride, and in the aqueous solution at 90°C was applied. The reaction is regioselective and leads mainly to γ -butyrolactone (Scheme 1).




Scheme 1.

Table 1. Yield of Lactonization from 4-Aryl Acids

Lactone	R ¹	R ²	R ³	Yield %
10a	H	H	H	41
10b	H	H	Me	51
10c	H	H	OH	84
10d	Me	Me	H	25
10e	H	H	PH	35
10f	OMe	OMe	H	0

Several 4-aryl carboxylic acids for one-pot direct conversion to the corresponding butyrolactones were elected. These acids were prepared from available starting materials by using straightforward procedures. The lactonization was entirely regioselective and only the γ -butyrolactones are isolated with 25–84% yields (Table 1).

Utilization of this oxidation reagent for intramolecular reaction of aryl acids of **11** (Scheme 2) to expected 5- and 6-members fused lactones **12** failed, and, in one case, lead to ester **13a**. Spectroscopic data for all compounds are shown in (Table 2).

*Scheme 2.*

Formation of lactone **10c** in high yield (Table 1) with hydroxy substitution instead of methoxy group of starting aryl acids **4c** at the para position, implies that the reaction has involved some extent of “through conjugation”¹⁸ (Scheme 3).

For the same reason, the 4-(2,5-dimethoxy phenyl) butyric acid **4f** was not converted to the corresponding lactone **10f**; instead, that entirely led to the formation of comparable p-benzoquinone, 4-(2,5-cyclohexadiene-1,4-dione) butanoic acid **19**, probably through stable radical **17** and biradical **18** (Scheme 4).


Table 2. Physical Property of All Synthesized Compounds

Entry	IR cm ⁻¹	¹ H & ¹³ C NMR δ	Yield (%)	M.p.°C & MS
3a	(KBr): 3200(b), 1690(b)	(CDCl ₃): 2.4(t) 2H, 3.4(t) 2H, 7.5(m) 3H, 8(m) 2H	90	114–116 (lit. 115) ¹⁹
4a	(KBr): 3200(b), 1700(s)	(CDCl ₃): 2.1(m) 2H, 2.5(t) 2H, 2.7(t) 2H, 7.3(m) 5H, 11.1(s) 1H	89	47–48 (lit. 47–48) ¹⁹
10a	(neat): 1770(s)	(CDCl ₃): 2.5(m) 4H, 5.4(t) 1H, 7.3(s) 5H, 13C (CDCl ₃): 32, 82, 125, 128, 129, 130, 165	41	Exact mass(M ⁺): calcd. = 162.0681, found 162.0675
3b	(KBr): 3200(b), 1680(sb)	(CDCl ₃): 2.5(s) 3H, 2.9(t) 2H, 3.3(t) 2H, 7.4(m) 2H, 7.9(m) 2H, 9.9(s) 1H	83	122–125
4b	(KBr): 3200(b), 1690(s)	(CDCl ₃): 1.9(m) 2H, 2.3(s) 3H, 2.4(t) 2H, 2.6(t) 2H, 7.1(s) 4H, 11.3(s) 1H	—	52–55 (lit. 57–48) ²⁰
10b	(CCl ₄): 3005(m), 1780(s)	(CDCl ₃): 2.3(m) 2H, 2.6(t) 3H, 5.5(t) 1H, 7.2(s) 4H 13C(CDCl ₃): 21, 29, 30, 81, 125, 126, 128, 129, 165	51	62–64 Exact mass(M ⁺):calcd. 176.0837, found 176.0836
3c	(KBr): 3200(b), 1700(s)	(CDCl ₃): 2.7(t) 2H, 3.2(t) 2H, 3.8(s) 3H, 6.9(d) 2H, 7.9(d) 2H	82	144–152 (lit. 148–150) ²¹
4c	(KBr): 3200(b), 1700(s)	(CDCl ₃): 1.9(t) 2H, 2.3(m) 2H, 2.6(m) 2H, 3.8(s) 3H, 7.0(m) 4H	71	56–58 (lit. 56–59) ²¹
10c	(neat): 3010(m), 1765(s)	(CDCl ₃): 2.5(m) 4H, 5.4(t) 1h, 7.3(m) 4H, 8.7(s) 1H 13C(CDCl ₃): 29, 30, 81, 126, 128, 176	84	Exact mass: (M ⁺): calcd. 178.0630, found 178.0627
3d	(KBr): 3200(b), 1680(sb)	(CDCl ₃): 2.3(s) 3H, 2.5(s) 3H, 2.7(t) 2H, 3.1(t) 2H, 7.0(s) 2H, 7.6(m) 1H, 11.3(s) 1H	85	100–102
4d	(KBr): 3200(b), 1685(s)	(CDCl ₃): 2(m) 2H, 2.5(s) 6H, 2.8(t) 2H, 5.8(t) 1H, 7.2(m) 3H	82	62–66


(continued)

Table 2. Continued

Entry	IR cm^{-1}	^1H & ^{13}C NMR δ	Yield (%)	M.p. $^\circ\text{C}$ & MS
10d	(CCl_4): 3000(m), 1775(s)	(CDCl_3): 2.5(m) 4H, 5.4(t) 1H, 7.3(s) 5H, 13C (CDCl_3): 32, 82, 125, 128, 129, 130, 165	25	Exact mass(M^+): calcd. 190.0994, found 190.0992
3e	(KBr): 3200(b), 1680(sb)	(CDCl_3): 2.8(t) 2H, 3.3 (t) 2H, 7.5(m) 7H, 8(m) 2H	62	180–182
4e	(KBr): 3200(b), 1680(s)	(CDCl_3): 1.9(m) 2H, 2.3(s) 3H, 2.4(t) 2H, 2.6(t) 2H, 7.1(s) 4H, 11.3(s) 1H	80	115–118
10e	(CCl_4): 3005(m), 1785(s)	(CDCl_3): 2.5(m) 4H, 5.5(t) 1H, 7.5(m) 9H	35	Exact mass: (M^+) calcd. 238.0996, found 238.0989
3f	(KBr): 3200(b), 1700(s)	(CDCl_3): 2.8(t) 2H, 3.4(t) 2H, 3.8(s) 3H, 3.9(s) 3H, 7.0(m) 2H, 7.3(m) 1H, 8.8(s) 1H	88	100–102 (lit. 102) ²²
4f	(KBr): 3200(b), 1700(s)	(CDCl_3): 1.9(m) 2H, 2.4(t) 2H, 2.6(t) 2H, 3.8(s) 6H, 6.7(s) 2H, 7.2(s) 1H	56	64–68 (lit. 68–69) ²²
19	(CCl_4): 3000(b), 1700(s)	(CDCl_3): 2(m) 2H, 2.5(t) 2H, 2.8(t) 2H, 6.8(m) 3H, 9(s) 1H	—	Exact mass: (M^+) calcd. 194.0595, found, 194.0592
13a	(neat): 3010(s), 3015(s), 1736(s)	(CDCl_3): 3.7(s) 2H, 5.11(s) 2H, 7.3(m) 10H	—	Exact mass: (M^+) calcd. 226.0994, found, 226.0989



Scheme 3.



Scheme 4.

EXPERIMENTAL

General

Yields refer to isolated pure center cut from column chromatography or scratched from preparative TLC. Products were characterized by comparison with authentic samples (IR, NMR, GC, TLC, and m.p.). Melting points are uncorrected and determined by Mettler Fp5 melting point apparatus. The ^1H and ^{13}C NMR spectra were recorded on a Bruker spectrometer, generally with TMS as internal standard. The IR spectra were recorded with a Shimadzu model 470. The high-resolution mass spectra were obtained from a Fisons Trio-1000 instrument.

Synthesis of 4-Phenyl-4-oxobutanoic Acids 3a: A Typical Procedure

Twenty mL (17.5 g, 0.225 mol) benzene (Na-dried and free of thiophene) and 3.4 g (0.034 mol) succinic anhydride were combined in a 100-mL flask equipped with a reflux condenser connected through a Y-junction to a single efficient gas absorption device. The succinic anhydride dissolved in the benzene, 10 g (0.075 mol) of powdered anhydrous aluminum chloride was added and stirred in the mixture solution. The reaction started immediately; HCl evolved and the mixture became hot. The resulting mixture was then allowed to warm up at room temperature and then again allowed to warm up for 30 m at gentle reflux. The solution was cooled in cold water. Then 15 mL water and 5 mL conc. HCl were added. Excess benzene was removed by steam distillation. The residue was cooled and filtered on a Buchner funnel and washed with 10 mL (1:3) dilute HCl and

1 mL H₂O. The crystals were boiled in Na₂CO₃ (4 g in 25 mL H₂O) for 15 min. The filtrate was cooled and acidified by conc. HCl at 0°C, 5.4 g (90%) crystals m.p. 114°–116°C (lit.,¹⁹ 115°C) was separated. The IR and NMR were recorded and shown in Table 2.

Preparation of 4-Phenyl Butanoic Acids 4a: A Typical Procedure

Zn powder (4.3 g), 0.43 g (1.58 mmol) mercury (II) chloride, 0.2 mL conc. HCl, and 5.5 mL water, were combined in a 50-mL flask. The mixture was stirred at room temperature for several minutes to produce a homogenous solution. After homogenization was completed, the stirring was stopped and the liquid was decanted as completely as possible. In a flask equipped with a reflux condenser, 2.7 mL water, 0.65 mL conc. HCl, 3.6 mL toluene (as solvent) and 1.8 g (0.01 mol) 4-phenyl-4-oxobutanoic acid **3a** were combined. The flask was refluxed vigorously for 30 h. During this period, 1.8 mL conc. HCl was added to the flask at approximately 6-h intervals during the refluxing period to maintain the conc. of HCl. After cooling, two layers were separated water (7.2 mL) was added to the aqueous layer, and was extracted with 3×30 mL ether. The extracted layer was added to toluene, washed with water, and dried over MgSO₄. The solvent was evaporated and the residue was distilled 185°C _{20 mmHg} (1.49 g, 89%), (m.p. 47°–48°C, lit.²⁰ 47°–48°C). The IR and NMR spectra were recorded (Table 2).

Preparation of 5-Phenyl- γ -butyrolactone 10a: A Typical Procedure

4-Phenyl butanoic acid **4a** (4.92 g, 30 mmol), 27 mL water, and 5.1 g (30 mmol) copper (II) chloride 2H₂O were combined in a 250-mL, two-neck, round-bottom flask equipped with a reflux condenser and an additional

Table 3. Red. Condition for Conversion of Keto Acids to 4-Aryl Acids

Keto Acids	Reagent & Method	Time of Red. (h)	Yield (%)	Aryl Acid
3a	amalgamated zinc ¹⁶	30	89	4a
3b	amalgamated zinc ¹⁶	45	—	4b
3c	amalgamated zinc ¹⁶	40	71	4c
3d	amalgamated zinc ¹⁶	42	82	4d
3e	amalgamated zinc ¹⁶	40	80	4e
3f	NH ₂ -NH ₂ , KOH ¹⁷	7	56	4f

funnel. A solution of 8.5 g (30 mmol) $\text{Na}_2\text{S}_2\text{O}_8$ and 15 mL water was added to the additional funnel. The reaction mixture was allowed to reflux by vigorous stirring while the temperature of solution was adjusted to $85^\circ\text{--}90^\circ\text{C}$. The solution from the additional funnel was added dropwise to a flask during 40 m, and the flask was refluxed for 3.5 h. After this time, the reaction was stopped. The flask was cooled and extracted with $3\times 30\text{ mL}$ ether and dried with MgSO_4 . The solvent was removed and 2 g (41%) of **10a** was collected as a pure center cut from silica gel column chromatography. The solvent system used was 5–10% EtOAc:ligroin. The IR, ^1H , ^{13}C NMR spectra and Exact mass (M^+) were recorded (Table 2).

Synthesis of 4-(4-Methoxyphenyl)-4-oxobutanoic Acid **3c**

Nitrobenzene (60 mL redistilled and dried) as a solvent and 2.5 g (0.025 mol) succinic anhydride were combined in a 250-mL, 2-neck, round-bottom flask equipped with a reflux condenser connected through a y-junction to a single efficient gas absorption device and an additional funnel. Then 6.67 g (0.05 mol) anhydrous AlCl_3 powder and 15 mL nitrobenzene (redistilled) from the additional funnel dropwise was added to the solution during 45 m. The resulting mixture was then stirred at room temperature for an additional 6 h. After this time, the reaction mixture was transferred to the solution of 300 mL HCl (20%) and 200 g ice, stirred vigorously, and extracted with $4\times 30\text{ mL}$ ether. The ether layers were combined and washed with $3\times 30\text{ mL}$ water, and extracted with $5\times 30\text{ mL}$ conc. NaHCO_3 . The aqueous layers were washed with $3\times 20\text{ mL}$ ether, and acidified with HCl. The white precipitate (3.4 g, 82% yield), m.p. $144^\circ\text{--}152^\circ\text{C}$ lit.²¹ $148^\circ\text{--}150^\circ\text{C}$. The IR, and NMR are recorded (Table 2)

Synthesis of 4-(4-Methoxy phenyl)butanoic Acid **4c**

A similar procedure as used for **4a** was applied, but instead of reflux for 30 h, the solution was refluxed for a 40-h period and 1.8 mL conc. HCl was added to the flask at 8-h intervals.

Preparation of 5-(4-Hydroxy phenyl)- γ -butyrolactone **10c**

A similar procedure for **10a** was used. The spectral data were recorded (Table 2).

ACKNOWLEDGMENTS

We appreciate the Research Committee of Guilan University for the partial support given to this study.

REFERENCES

1. Corey, E.J.; Kang, M. J. Am. Chem. Soc. **1984**, *106*, 5384.
2. Snider, B.B.; Mohan, R.; Kates, S.A. J. Org. Chem. **1985**, *50*, 3661.
3. Fristad, W.E.; Hershberger, S.S. J. Org. Chem. **1985**, *50*, 1026.
4. Kraus, G.A.; Landgrebe, K. Tetrahedron Lett. **1984**, *25*, 3939.
5. Ogibin, Y.N.; Rakhmatulline, L.K.; Nikishin, G.I. Izu. An SSSR, Ser. Khim. **1975**, 2723.
6. Anderson, J.M.; Kochi, J.K. J. Am. Chem. Soc. **1970**, *92*, 1651.
7. Ogibin, Y.N.; Troyansky, E.I.; Nikishin, G.I.; Kadentsev, V.I.; Lubuzh, E.D.; Chizhov, O.S. Izu. An SSSR, Ser. Khim. **1976**, 2518.
8. Bacha, J.D.; Kochi, J.K. Tetrahedron **1968**, *24*, 2215.
9. Anderson, J.M.; Kochi, K.J. J. Org. Chem. **1970**, *35*, 986.
10. Lande, S.S.; Kochi, J.K. J. Am. Chem. Soc. **1968**, *90*, 5196.
11. Kenyon, J.; Symons, M.C.R. J. Chem. Soc. **1953**, *2129*, 3580.
12. Nikishin, G.I.; Svitanko, I.V.; Troyansky, E.I. J. Chem. Soc., Perkin Trans. 2 **1983**, 595.
13. Mahmoodi, N.O.; Mirmohammadi, S.H., Submitted for Publication, *in press*.
14. Belletire, J.L.; Mahmoodi, N.O. Tetrahedron Lett. **1989**, *30*, 4363.
15. Belletire, J.L.; Mahmoodi, N.O. Synth. Commun. **1989**, *19*, 3371.
16. Belletire, J.L.; Mahmoodi, N.O. J. Nat. Prod. **1992**, *55*, 194.
17. Mahmoodi, N.O., *Dissertation Abstracts International*, University of Michigan, MI, USA 1992, *52*, 8.
18. Brown, H.C.; Okamoto, Y. J. Am. Chem. Soc. **1958**, *80*, 4979.
19. Brown, P.M.; Russell, J.; Thomsoon, R.H.; Wylile, A.G. J. Chem. Soc. (c) **1968**, 842.
20. *Dictionary of Organic Compounds*, 5th Ed.; Chapman and Hall: Pennsylvania, 1982.
21. *Aldrich Catalog-Handbook of Fine Chemicals*, 1993–1994.
22. Shoekravi, A. Iran J. Chem. and Chem. Eng. **1995**, *14*, 23.
23. Clemmensen, E. Ber. **1913**, *46*, 1837.
24. Cannizzaro, S. Ann. **1835**, *88*, 129.

Received in the UK September 21, 1999

