This article was downloaded by: [University of Calgary] On: 13 March 2013, At: 05:16 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

DEALKYLATION OF PHOSPHONATE ESTERS WITH CHLOROTRIMETHYLSILANE

A. J. Gutierrez ^a , E. J. Prisbe ^a & J. C. Rohloff ^a ^a Department of Chemical Development, Gilead Sciences, Inc., 333 Lakeside Dr., Foster

City, California, 94404, U.S.A. Version of record first published: 07 Feb 2007.

To cite this article: A. J. Gutierrez, E. J. Prisbe & J. C. Rohloff (2001): DEALKYLATION OF PHOSPHONATE ESTERS WITH CHLOROTRIMETHYLSILANE, Nucleosides, Nucleotides and Nucleic Acids, 20:4-7, 1299-1302

To link to this article: http://dx.doi.org/10.1081/NCN-100002541

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

NUCLEOSIDES, NUCLEOTIDES & NUCLEIC ACIDS, 20(4-7), 1299-1302 (2001)

DEALKYLATION OF PHOSPHONATE ESTERS WITH CHLOROTRIMETHYLSILANE

A. J. Gutierrez,* E. J. Prisbe, and J. C. Rohloff

Department of Chemical Development, Gilead Sciences, Inc., 333 Lakeside Dr., Foster City, California 94404

ABSTRACT

Chlorotrimethylsilane completely dealkylates phosphonate esters at elevated temperature in a sealed reaction vessel. These conditions are tolerated by a variety of functional groups and lead to high conversions of dimethyl, diethyl and disopropyl phosphonates to their corresponding phosphonic acids.

Phosphonic acids are generated from their dialkyl esters by reaction with conc. HCl or HBr, however the conditions are too harsh for many functional groups (1,2). Bromotrimethylsilane (TMSBr) selectively cleaves PO dialkyl esters yielding bis(trimethylsilyl) esters which are easily hydrolyzed with water under neutral conditions (3). In contrast, chlorotrimethylsilane (TMSCl) has been used mainly for deprotection of the more labile dimethyl phosphonates (4). For diethyl phosphonates, the low reactivity of TMSCl results in long reaction times and unsatisfactory yields (5). The rate of diester cleavage is accelerated by addition of sodium or lithium iodide (6). Here we report a procedure using TMSCl to completely dealkylate diethyl and diisopropyl phosphonates in high yield. The procedure is amenable to laboratory and industrial scale preparations.

To assess the ability of TMSCl to cleave dialkyl phosphonates, 1.0 M solutions of **1** in chlorobenzene were mixed with TMSCl in sealed glass pressure tubes and heated at temperatures ranging from $130-140^{\circ}$ C (Fig. 1). As substitution to **2** progressed, modest increases in the internal pressure of the reaction vessels were observed due to the formation of volatile alkyl chloride. The pressure

^{*}Corresponding author.

ORDER		REPRINTS
-------	--	----------

1300



Figure 1. Cleavage of phosphonate diesters to phosphonic acids using TMSCI.

reached a maximum at reaction completion, then returned to atmospheric pressure on cooling back to room temperature. The conversions to phosphonic acids 3 were monitored by evaporating an aliquot of the reaction mixture, adding D_2O and observing the resulting formation of phosphonic acids 3 by 1 H and 31 P NMR (3) (Table 1).

Dialkyl phosphonates (1a-g) containing several functional groups were evaluated for ease of diester cleavage and compatibility to the reaction conditions. Heating 1a-g with TMSCl in chlorobenzene at 130-140°C, followed by hydrolysis, lead to complete conversions to phosphonic acids 3a-g (Table 1). In most cases these deprotections went to completion in 8 to 12 hours. The high percent conversions to 3a-g show that a variety of functional groups including carboxylic esters, ethers and alkenes are compatible with these conditions. Among the compounds evaluated, the more labile dimethyl phosphonate 1a was the most easily deprotected by TMSCl in chlorobenzene. However, diethyl phosphonates 1b, 1d-g were also completely cleaved after slightly longer heating times and even the more hindered isopropyl ester groups of 1c were removed if a greater excess of TMSCl was used. These rates of cleavage are consistent with their rates of cleavage by TMSBr (3). The heating time required for diester cleavage of diethyl phosphonates **1e** and **1f** with TMSCl in chlorobenzene was 10 h. By comparison, heating 1e and 1f with TMSCl, in the absence of solvent, afforded their phosphonic acids 3e and 3f after 4

Phosphonate 1	R	R′	TMSCl (equiv.)	Temp. (°C)	Time (h)	% Conv. ^b 3
1a	CH ₃ OOCCH ₂	CH ₃	3	130	8	98
1b	C ₂ H ₅ OOCCH ₂	C_2H_5	4	140	18	98
1c	C ₂ H ₅ OOCCH ₂	$(CH_3)_2CH$	6	140	36	98
1d	C ₆ H ₅ OCCH ₂	C_2H_5	4	140	12	98
1e	C ₆ H ₅ CH ₂	C_2H_5	4	140	10	99
1f	CH ₂ =CH	C_2H_5	4	140	10	98
1g	CH ₃ OCH ₂ CH ₂ OCH ₂ ^c	C_2H_5	4	140	8	>99

Table 1. Cleavage of Dialkyl Phosphonates 1^a with Chlorotrimethylsilane in Chlorobenzene

^aFor descriptions of R and R', see Figure 1.

^bPercent conversions of 1 to 3 were estimated by measuring their ¹H and ³¹P NMR absorbances in D_2O

^cFor preparation of this phosphonate see [7].

Marcel Dekker, Inc.

270 Madison Avenue, New York, New York 10016

ORDER		REPRINTS
-------	--	----------

DEALKYLATION OF PHOSPHONATE ESTERS



Figure 2. Preparation of PMEA and PMPA from their diethylesters using TMSCI.

and 5 days respectively (4). The deprotections with TMSCl also work well in DMF, acetonitrile, and dichloroethane. However the latter two solvents produce higher reaction vessel pressures.

The deprotection of dialkyl phosphonates using TMSCl is potentially very useful in the large scale synthesis of phosphonic acid derivatives as a replacement for TMSBr which is considerably more expensive and more difficult to handle. For instance, 9-[2-(phosphonomethoxy)ethyl]adenine (PMEA) **6** and (R)-9-[2-(phosphonomethoxy)propyl]adenine (PMPA) **7** continue to attract widespread attention as broad spectrum antivirals including potent and selective activity against human immunodefficiency virus (8).

One reported synthesis of **6** and **7** utilizes TMSBr for cleavage to the phosphonic acids from the diethyl esters **4** and **5** (9). By comparison, the same deprotections were performed with TMSCl (10,11). (Fig. 2) affording **6** and **7** in equivalent yields and purities as those obtained with TMSBr. The process is amenable to large scale synthesis. Equivalent results were obtained for preparing **6** and **7** when the process was scaled to 5 kg in a 30 gal glass-lined Pfaudler reactor. The ease in which **4** and **5** are deprotected with TMSCl in chlorobenzene is in striking contrast to deprotection by heating with TMSCl in the absence of solvent, which is not effective in deprotecting adenosine diethyl phosphonates (12).

In summary, TMSCl in chlorobenzene readily cleaves dialkyl phosphonates in high yield in the presence of a wide variety of functional groups. These conditions also efficiently cleave the ethyl ester groups of adenine containing diethylphosphonates. TMSCl should be a useful and economical reagent for laboratory and industrial scale preparation of phosphonic acid derivatives.

REFERENCES

- Kosolopoff, G. M. in "Organophosphorus Compounds", John Wiley and Sons, Inc., New York, 1950.
- 2. For the deprotection of diethyl PMPA **5** with conc. HCl, we observe 10–16% acid catalyzed deamination of adenine to hypoxanthine. Data not shown.

Copyright @ Marcel Dekker, Inc. All rights reserved

Marcel Dekker, Inc.

1301

ORDER		REPRINTS
-------	--	----------

GUTIERREZ, PRISBE, AND ROHLOFF

- a) McKenna, C. E.; Schmidhauser, J. J. C. S. Chem. Commun. 1979, 739. b) McKenna, C. E.; Higa, M. T.; Cheung, M. T.; McKenna, M.-C. Tetrahedron Lett. 1977, 155–158.
 c) Rudinskas, A. J.; Hullar, T. L. J. Org. Chem. 1977, 42, 2771–2776.
- 4. a) Rabinowitz, R.; J. Org. Chem. 1963, 28, 2975–2978. b) Paulsen, H.; Thiem, J. Chem. Ber. 1973, 106, 3850–3876.
- Baer, E.; Robinson, R. Can. J. Chem. 1973, 51, 104–110. b) Paulsen, H.; Bartsch, W. Chem. Ber. 1975, 108, 1732–1744.
- 6. a) Machida, Y.; Nomoto, S.; Saito, I. *Synthetic Comm.* **1979**, *9*, 97–102. b) Morita, T.; Okamoto, Y.; Sakurai, H. *Tetradedron Lett.* **1978**, 2523–2526.
- 7. Kluge, A. F.; Cloudsdale, I. S. J. Org. Chem. 1979, 44, 4847–4852.
- a) Tsai, C.-C.; Folis, K. E.; Sabo, A.; Beck, T. W.; Grant, R. F.; Bischofberger, N.; Benveniste, R. E.; Black, R. Science 1995, 270, 1197–1199. b) Naesens, L.; Balzarini, J.; De Clercq, E. Rev. Med. Virol. 1995, 4, 147. c) Kim, C. U.; Luh, B. Y.; Misco, P. F.; Bronson, J. J.; Hitchcock, M. J. M.; Ghazzouli, I.; Martin, J. C. J. Med. Chem. 1990, 33, 1207–1213. d) Bronson, J. J.; Kim, C. U.; Ghazzouli, I.; Hitchcock, M. J. M.; Kern, E.; Martin, J. C. Nucleosides and Antiviral Agents; Martin, J. C., Ed.; American Chemical Society, Washington, DC, 1989, 72–87. e) Starret, J. E. Jr.; Tortolani, D. R.; Russell, J.; Hitchcock, M. J. M.; Whiterock, V.; Martin, J. C.; Mansuri, M. M. J. Med. Chem. 1994, 37, 1857–1864. f) Balzarini, J.; Aquaro, S.; Perno, C. F.; Witvrouw, M.; Holy, A.; DeClercq, E. Biochem. Biophys. Res. Commun. 1996, 219, 337.
- Schultz, L. M.; Chapman, H. H.; Dupree, N. J.; Jones, R. J.; Kent, K. M.; Lee, T. T.; Louie, M. J.; Postich, M. J.; Prisbe, E. J.; Rohloff, J. C.; Yu, R. H. *Tetrahedron Lett.* **1998**, *39*, 1853–1856.
- 10. General procedure for cleavage of dialkyl phosphonates preparation of **6**: Diethyl PMEA 4 (9) (41.1 g, 0.125 mol) and chlorobenzene (0.125 L) were charged to a 1 L ACETM glass pressure reactor containing a thermowell and a pressure gauge. The contents were stirred while TMSCI (56 mL, 0.44 mol) was added slowly. The reaction vessel was purged with N₂, sealed, and heated to 125°C. The internal reactor pressure increased incrementally to a maximum of 30 psi after 9 h. The contents were cooled to r.t. at which time the pressure returned to ambient pressure and water (160 mL) was added with vigorous stirring. The layers were separated and the lower aqueous layer containing the product was collected. The aqueous layer was adjusted to pH = 3.2 with 25% NaOH (~36 g), the resulting slurry cooled to 0°C and the solid collected by vacuum filtration. Water (40 mL) was added to the wet cake, the resulting slurry was heated to 70°C for 1 h, and then cooled to 0°C. The solid product was collected by vacuum filtration and dried under vacuum (50°C/28 mmHg) affording 33.0 g (0.120 mol, 96%) PMEA **6**. NMR data for PMEA **6** was consistent with literature values. Holy, A.; Rosenberg, I. *Collect. Czech. Chem. Commun.* **1987**, *52*, 2801–2809.
- In the same manner as above 48.3 g (0.140 mol) diethyl PMPA 5 (9) were deprotected with 4.5 equiv. TMSCl and yielded 30.3 g (0.105 mol, 75%) of PMPA 7 (99% pure by quantitative HPLC assay versus an external standard). ¹H NMR (300 MHz, D₂O, δ): 8.31 (s, 2H, Adenine–2H, –8H), 4.39 (dd, J = 14, 3 Hz, 1H, –CH₂N), 4.20 (dd, J = 14, 7 Hz, 1H, –CH₂N), 3.89 (m, 1H, –CH–), 3.58 (dd, J = 13, 9 Hz, 1H, –OCH₂P), 3.38 (dd, J = 13, 9 Hz, 1H, –OCH₂P), 1.07 (d, J = 6 Hz, 3H, –CH₃) ³¹P NMR (121 MHz, D₂O, δ): 15.79.
- 12. Hampton, A.; Sasaki, T.; Paul, B. J. Am. Chem. Soc. 1973, 95, 4404-4414.

Copyright © Marcel Dekker, Inc. All rights reserved

1302



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the <u>U.S. Copyright Office</u> for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on <u>Fair Use in the Classroom</u>.

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our <u>Website</u> User Agreement for more details.

Order now!

Reprints of this article can also be ordered at http://www.dekker.com/servlet/product/DOI/101081NCN100002541