

Tetrahedron Letters 41 (2000) 3509-3512

TETRAHEDRON LETTERS

A norbornyl route to cyclohexitols: stereoselective synthesis of conduritol-E, *allo*-inositol, MK 7607 and gabosines

Goverdhan Mehta* and Sripada Lakshminath

Department of Organic Chemistry, Indian Institute of Science, Bangalore 560 012, India

Received 14 February 2000; accepted 7 March 2000

Abstract

A novel fragmentation sequence within the norbornane system, involving C_1-C_7 bond scission, provides convenient access to a highly functionalized and versatile cyclohexenoid building block which has been further elaborated to a range of cyclohexitols such as, conduritol E, *allo*-inositol and gabosine B. Our synthesis of the structure corresponding to gabosine K indicates that the structure of this natural product needs to be revised. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: fragmentation reactions; cyclitols; hydroxylation.

Molecular entities having cyclohexitol (polyhydroxylated cyclohexanoids) or amino cyclohexitol core structures have attracted a great deal of attention in recent years because of diverse biological activities exhibited by them, ranging from herbicidal and anti-microbial on the one hand to glycosidase inhibition and mediation of cellular communication on the other.¹ Cyclohexitols with varied levels of oxygenation and stereochemical patterns are known and some typical examples are zeylenol **1**, conduritol-E **2**, *allo*-inositol **3**, valienamine **4** and MK 7607 **5**. While several of the known cyclohexitols have been encountered in nature, many more have been accessed through syntheses to evaluate their biological potential. The syntheses of cyclohexitols remains an active area of current research and new approaches continue to be explored to achieve greater preparative efficiency and higher levels of stereo- and regiocontrol.^{1,2} In the accompanying communication,³ we have outlined a novel Groblike 'top-to-bottom' fragmentation sequence in a norbornyl derivative **6** to furnish a building block **7** for polyoxygenated cyclohexenoids and demonstrated its utility in the synthesis of carbasugars, e.g. fucopyranose **8**, Scheme 1.



* Corresponding author. E-mail: gm@orgchem.iisc.ernet.in (G. Mehta)

0040-4039/00/\$ - see front matter © 2000 Elsevier Science Ltd. All rights reserved. *PII:* S0040-4039(00)00409-3



Scheme 1.

The structural similarity between 7 and the *cis*-dihydroxylated cyclohexenoids obtained from the microbial oxygenation^{2b} of arenes spurred us to further explore the potential of 7 in a variety of cyclohexitol syntheses and these endeavours form the subject matter of this letter.

Our first objective was to elaborate 7 to conduritol and inositol type cyclohexitols. Towards this end, 7 was transformed to the bis-acetonide tosylate 9 and further elimination via the iodide gave the olefin 10.⁶ Ozonolysis to cyclohexanone 11, reduction and mesylation furnished 12.⁶ Base mediated elimination to 13⁶ and acetonide deprotection led to the conduritol-E 2, Scheme 2.^{4,6}



Scheme 2. Reagents and conditions: (a) i. NaI, Me₂CO, Δ , 30 h, 92%; ii. ¹BuO⁻K⁺, ¹BuOH, Δ , 20 h, 70%; (b) O₃, NaHCO₃, DCM, -78°C, 5 min, DMS, 90%; (c) i. NaBH₄, MeOH, 0°C, 30 min, 89%; ii. MsCl, Et₃N, DCM, -10°C, 30 min, 90%; (d) ¹BuO⁻K⁺, DMSO, rt, 2 h, 75%; (e) Amberlyst-15, aq. MeOH, rt, 20 h, 93%

The bis-acetonide 13 also served as the precursor of *allo*-inositol 3. Catalytic osmium tetraoxide dihydroxylation of 13 was understandably slow but did lead to 14^6 in reasonable yield. Deprotection in 14 furnished *allo*-inositol 3, which was conveniently characterised as the hexa-acetate 15, Scheme $3.^{5,6}$



Scheme 3. Reagents and conditions: (a) OsO_4 , NMMO, $Me_2CO:H_2O$ (4:1), rt, 7 days, 70%; (b) 5% HCl, $H_2O:Et_2O$ (4:1), rt, 30 h; (c) Ac_2O , Py, 80°C, 6 h, 65% (two steps)

Recently, structure elucidation of nearly a dozen novel secondary metabolites, named gabosines A–K, bearing the C_7 skeleton of carbasugars has been reported from the various strains of *Streptomycetes*.⁷ We recognised that our building block **7** and particularly the olefin **10** derived from it were well suited for the synthesis of gabosines. We first chose gabosine B **17** as the target and its synthesis was achieved in

a straightforward manner. Exposure of **10** to rhodium trichloride resulted in isomerisation of the double bond to the desired tetra-substituted position and simple hydrolysis of 16^6 furnished gabosine B **17**, identical in all respects with the natural product, Scheme 4.⁶ Stereoselective protonation in **16** during hydrolysis to furnish **17** as a single diastereomer is notable.



Scheme 4. Reagents and conditions: (a) RhCl₃, NaHCO₃, EtOH, Δ , 20 h, 60%; (b) 5% HCl, H₂O:Et₂O (4:1), rt, 90%

We next turned our attention to the more embellished gabosine K **18** and the patented herbicide MK 7607 **5** from *Curvularia* sp.,⁸ closely related to the former. OsO₄-mediated dihydroxylation of **10** led to readily separable diastereomeric diols **19** and **20** (70:30) (Scheme 5).⁶ Selective acetylation of the primary hydroxyl group in the major isomer **19** and dehydration furnished the olefins **21** and **22** (1:2) in which the latter could be recycled.⁶ A one-pot acetonide deprotection and acetate hydrolysis in **21** led to MK 7607.⁶ On the other hand, only acetonide deprotection in **21**, under controlled conditions, resulted in **18**,⁹ the structure assigned to gabosine K. However, we found that the spectroscopic data (¹H and ¹³C NMR) reported for gabosine K were different from our synthetic **18** and a direct comparison of the spectra provided by the authors confirmed this surmise. The structure of gabosine K therefore needs to be revised.⁹



Scheme 5. Reagents and conditions: (a) OsO₄, NMMO, Me₂CO:H₂O (4:1), rt, 2 days, 95%; (b) i. Ac₂O, DMAP, DCM, 0°C, 30 min, 100%; ii. SOCl₂, Py, DCM, rt, 6 h, 45% for **21** and **22**; (c) Amberlyst-15, THF:H₂O (2:3), rt, 48 h, 85%; (d) 5% HCl, H₂O:Et₂O (4:1), rt, 2 days, >95%

In short, we have amplified the utility of the cyclohexenoid building block 7, obtained via a novel 'top-to-bottom' Grob-like fragmentation, in diverse cyclohexitol syntheses. Besides conduritol E 2 and *allo*-inositol 3, synthesis of these members of the gabosine family has been accomplished for the first time.

Acknowledgements

We are grateful to Professor A. Zeeck for the copies of the spectra and helpful correspondence. S.L. thanks CSIR for the award of a Research Fellowship. Part of this work was initiated at the University of Hyderabad. We thank SIF at I.I.Sc. for high field NMR data.

References

- Reviews: (a) Heightman, T. D.; Vasella, A. T. Angew. Chem., Int. Ed. Engl. 1999, 38, 750. (b) Suami, T.; Ogawa, S. Adv. Carbohydr. Chem. Biochem. 1990, 48, 21. (c) Suami, T. Top. Curr. Chem. 1990, 154, 257. (d) Berridge, M. J.; Irvine, R. F. Nature 1989, 341, 197.
- (a) Balci, M.; Sutbeyaz, Y.; Secen, H. *Tetrahedron* 1990, 46, 3715. (b) Hudlicky, T.; Entwistle, D. A.; Pitzer, K. K.; Thorpe, A. J. *Chem. Rev.* 1996, 96, 1195 and references cited therein.
- 3. Mehta, G.; Mohal, N.; Lakshminath, S. Tetrahedron Lett. 2000, 41, 3505.
- 4. For some of the recent syntheses of Conduritol-E, see: (a) Landais, Y.; Angelaud, R. J. Org. Chem. 1996, 61, 5202. (b) Honzumi, M.; Hiroya, K.; Taniguchi, T.; Ogasawara, K. J. Chem. Soc., Chem. Commun. 1999, 1985. (c) Liu, P.; Dobbelaere, J.; Eycken, V.; Vandewalle, M. Synlett 1992, 243. (d) Hudlicky, T.; Luna, H.; Olivo, H. F.; Anderson, C.; Nugent, T.; Price, J. D. J. Chem. Soc., Perkin Trans. 1 1991, 2907.
- (a) Motherwell, W. B.; Williams, A. S. Angew. Chem., Int. Ed. Engl. 1995, 34, 2031.(b) Tschamber, T.; Backenstrass, F.; Fritz, H.; Streith, J. Helv. Chim. Acta 1992, 75, 1052. (c) Mandel, M.; Hudlicky, T.; J. Chem. Soc., Perkin Trans. 1 1993, 741.
- 6. All new compounds reported here were racemic and characterized on the basis of spectroscopic data (¹H and ¹³C NMR, *J* in Hz) and elemental analyses. Selected spectroscopic data **10**: $\delta_{\rm H}$ (300 MHz, CDCl₃): 5.19 (1H, br s), 5.06 (1H, br s), 4.58 (2H, ABq, *J* 7), 4.47 (2H, br s), 2.56 (1H, br d, *J* 13), 2.16 (1H, d, *J* 13), 1.48 (3H, s), 1.37 (6H, s), 1.28 (3H, s); $\delta_{\rm C}$ (75 MHz, CDCl₃): 139.61, 116.77, 108.36, 107.74, 76.22, 74.39, 73.74, 72.68, 31.84, 26.25, 26.03, 24.26, 23.88. **13**: $\delta_{\rm H}$ (300 MHz, CDCl₃): 5.73 (2H, s), 4.58–4.52 (4H, m), 1.38 (6H, s), 1.37 (6H, s); $\delta_{\rm C}$ (75 MHz, CDCl₃): 126.93 (2C), 109.14 (2C), 73.24 (2C), 70.23 (2C), 27.83 (2C), 26.43 (2C). **2**: $\delta_{\rm H}$ (300 MHz, D₂O, ref. acetone): 5.74 (2H, m), 4.17 (2H, m), 3.78 (2H, m); $\delta_{\rm C}$ (75 MHz, D₂O, ref. acetone) 130.23 (2C), 69.63 (2C), 67.14 (2C). **15**: $\delta_{\rm H}$ (300 MHz, CDCl₃): 5.52–5.42 (3H, m), 5.36–5.3 (3H, m), 2.08 (18H, s); $\delta_{\rm C}$ (100 MHz, CDCl₃): 169.71 (3C), 169.45 (3C), 67.60 (3C), 67.21 (3C), 20.71 (3C), 20.64 (3C). **17**: $\delta_{\rm H}$ (400 MHz, CD₃OD): 4.41 (1H, d, *J* 10), 4.11–4.09 (1H, m), 3.45 (1H, dd, *J* 10, 3), 2.97–2.87 (1H, m), 2.11 (1H, ddd, *J* 14, 6, 3.5), 1.42 (1H, dt, *J* 14, 2), 1.03 (3H, d, *J* 6.5); $\delta_{\rm C}$ (100 MHz, CD₃OD): 211.52, 78.26, 78.19, 69.78, 38.18, 37.88, 13.75. **5**: $\delta_{\rm H}$ (400 MHz, D₂O): 5.46 (1H, br s), 4.12 (1H, br s), 4.05 (1H, br s), 3.95 (2H, br s), 3.68 (2H, br s); $\delta_{\rm C}$ (100 MHz, D₂O): 141.07, 124.93, 69.34, 69.00, 67.40, 66.72, 62.84.
- 7. Bach, G.; Breiding-Mack, S.; Grabley, S.; Hammann, P.; Hutter, K.; Thiericke, R.; Uhr, H.; Wink, J.; Zeeck, A. Liebigs. Ann. Chem. 1993, 241.
- 8. Yoshikawa, N.; Chiba, N.; Mikawa, T.; Ueno, S.; Harimaya, K.; Iwata, M. Chem. Abstr. 1995, 122, 185533e.
- 9. Spectroscopic data for **18**: $\delta_{\rm H}$ (400 MHz, CD₃OD): 5.85–5.75 (1H, m), 4.70 (1H, $\frac{1}{2}$ ABq, *J* 13.5), 4.60 (1H, $\frac{1}{2}$ ABq, *J* 13.5), 4.27–4.23 (2H, m), 3.90–3.87 (2H, m), 2.07 (3H, s); $\delta_{\rm C}$ (75 MHz, CD₃OD): 172.51, 137.15, 128.21, 70.93, 70.67, 67.96, 67.40, 65.75, 20.78. Spectroscopic data reported⁷ for gabosine K: $\delta_{\rm H}$ (200 MHz, CD₃OD): 2.06 (s, 9-H₃), 3.39 (m, 2H, *J* 8, 6.5, 3-H and 4-H), 4.04–4.18 (m, 2H, 1-H and 2-H), 4.53/4.73 (2 d, *J*_{AB} 13.5, m, 7-H₂), 5.59 (m, 6-H); $\delta_{\rm C}$ (50.3 MHz, CD₃OD): 172.5, 136.2, 128.8, 77.5, 77.1, 73.5, 73.0, 64.8, 20.8. These spectral data reveal that the two compounds though not identical, do share common structural features indicating that the natural product may have a stereoisomeric structure related to **18**.