This article was downloaded by: [Universitat Politècnica de València] On: 26 October 2014, At: 09:30 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lncn20

MILD, EFFICIENT, SELECTIVE AND "GREEN" BENZOYLATION OF NUCLEOSIDES USING BENZOYL CYANIDE IN IONIC LIQUID

Ashok K. Prasad^a, Vineet Kumar^a, Jyotirmoy Maity^a, Yogesh S. Sanghvi^b, Vasulinga T. Ravikumar^b & Virinder S. Parmar^a

^a Bioorganic Laboratory, Department of Chemistry, University of Delhi, Delhi, India

^b Isis Pharmaceuticals Inc., Carlsbad, California, USA Published online: 15 Nov 2011.

To cite this article: Ashok K. Prasad, Vineet Kumar, Jyotirmoy Maity, Yogesh S. Sanghvi, Vasulinga T. Ravikumar & Virinder S. Parmar (2005) MILD, EFFICIENT, SELECTIVE AND "GREEN" BENZOYLATION OF NUCLEOSIDES USING BENZOYL CYANIDE IN IONIC LIQUID, Nucleosides, Nucleotides and Nucleic Acids, 24:5-7, 747-751, DOI: <u>10.1081/NCN-200060065</u>

To link to this article: http://dx.doi.org/10.1081/NCN-200060065

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



MILD, EFFICIENT, SELECTIVE AND "GREEN" BENZOYLATION OF NUCLEOSIDES USING BENZOYL CYANIDE IN IONIC LIQUID

Ashok K. Prasad, Vineet Kumar, and Jyotirmoy Maity • Bioorganic Laboratory, Department of Chemistry, University of Delhi, Delhi, India

Yogesh S. Sanghvi* Isis Pharmaceuticals Inc., Carlsbad, California, USA

Vasulinga T. Ravikumar • Isis Pharmaceuticals Inc., Carlsbad, California, USA

Virinder S. Parmar Bioorganic Laboratory, Department of Chemistry, University of Delhi, Delhi, India

Use of benzoyl cyanide (BzCN) for benzoylation of nucleosides has been studied, both in pyridine and in ionic liquid. BzCN in 1-methoxyethyl-3-methylimidazolium methanesulfonate as ionic liquid has been found to be a "green" alternative compared to the pyridine-BzCN system. An efficient and selective benzoylation of nucleosides of both, the 2'-deoxy- and the ribo-series at ambient temperature was accomplished.

INTRODUCTION

One of the intrinsic problems, during the course of multistep synthetic protocols in nucleic acid chemistry, is the selective manipulation of hydroxyl and amino groups using labile protecting groups. This problem is further aggravated by poor solubility of nucleosides in common organic solvents. One of the most frequently used and preferred protecting groups for the protection of hydroxyl and amino groups in nucleoside and nucleotide chemistry is the benzoyl group. This is due to its less pronounced tendency to vicinal migration in comparison to acetyl group.^[1] The classical method for benzoylation of hydroxyl and amino groups is treatment of the substrate with benzoyl chloride/benzoic anhydride in the presence of pyridine, which usually requires extended time periods, elevated temperatures

This project was funded by ISIS pharmaceuticals Inc., CA. VK and JM thank CSIR, New Delhi, India, for the award of Junior Research Fellowships to them.

^{*}Current Address: Rasayan Inc., 2802 Crystal Ridge Road, Encinitas, CA 92008, USA.

Address correspondence to Ashok K. Prasad, Bioorganic Laboratory, Department of Chemistry, University of Delhi, Delhi 110 007, India.

A. K. Prasad et al.



FIGURE 1

or time-consuming work-up procedure. Holy and Soucek^[2] have demonstrated the application of benzoyl cyanide (BzCN) in acetonitrile/dimethylformamide as a mild benzoylating agent for the protection of nucleosides. However, these authors did not study this reagent on all the eight natural 2'-deoxy- and ribonucleosides. Our ongoing interest in acylation reactions on this class of compounds encouraged us to expand and thoroughly investigate the application of BzCN as a suitable benzoylation reagent.^[3-7] Herein, we report results of benzoylation studies on nucleosides of both, the 2'-deoxy- and the ribo-series using BzCN in pyridine and in 1-methoxyethyl-3-methylimidazolium methanesulfonate (MoeMIM.Ms) as an ionic liquid (IL).

Benzoylation of Nucleosides Using BzCN in Pyridine as Solvent

Since most of the unprotected nucleosides are soluble in pyridine at high concentrations, it was chosen as a solvent of choice. In a typical experiment, nucleoside (1 mmol) was dissolved in pyridine (5 mL) and DMAP (10 mg) was added, followed by the addition of BzCN (2.5-8.0 mmol). The reaction was stirred at 40°C (at 115°C in case of 2'-deoxyguanosine and guanosine) and the progress monitored by TLC. On completion, the reaction mixture was poured over crushed ice while stirring in a fume hood. The precipitate was filtered off and washed with water, followed by petroleum ether to afford benzoylated nucleosides as white crystalline compounds in 11.0-96.5% yields. The results of benzoylation are summarized in Figure 1.



FIGURE 2 1-Methoxyethyl-3-methylimidazolium methanesulphonate (MoeMIM.Ms).

TABLE 1							
Starting compd.	BzCN (equiv.)	Reaction time/temp	Product(s)	Yield (%)*	HRMS Calcd./Obs.		
la	2.5	5 h/40°C	3 a	94.0	[M + H] ⁺ 451.1505/451.1521		
1 b	2.1	6 h/40°C	3 b	93.0	$[M + H]^+$ 460.1621/460.1636		
1c	2.2	2 h/40°C	3 c	79.3*	$[M + H]^+$ 436.1509/436.1511		
			3 d	17.2*	$[M + H]^+$ 540.1771/540.1794		
1c	8.0	$2 \text{ h/}40^{\circ}\text{C}$	3 d	96.5*	$[M + H]^+$ 540.1771/540.1794		
1 d	6.0	4 h/115°C	3e	89.0	$[M + H]^+$ 580.1832/580.1886		
2 a	5.0	3 h/40°C	4a	89.0	$[M + H]^+$ 571.1717/571.1736		
2 b	6.0	5 h/40°C	4 b	71.5*	[M + Na] ⁺ 602.1652/602.131		
			4c	11.0*	[M + Na] ⁺ 682.1801/682.1901		
2 c	6.8	7 h/40°C	4 d	92.0	[M + Na] ⁺ 682.1801/682.1901		
2 d	7.0	7 h/115°C	4e	88.5*	[M + Na] ⁺ 722.1863/722.1981		

*The yields reported are on the basis of LC-MS analysis data.

Benzoylation of Nucleosides Using BzCN in Ionic Liquid as Solvent

In recent years, room-temperature ILs have attracted much attention as the "green" solvents of the future. This is because of their negligible vapor pressure, thermal stability, unprecedented ability to dissolve a broad range of compounds of organic and inorganic nature and easy reuse. We have investigated the applications of BzCN as benzoylating agent in the IL MoeMIM.Ms (Figure 2). This IL was chosen as reaction medium (or solvent) because of its ability to dissolve a variety of nucleosides^[5] (Table 1).

In a typical experiment, nucleoside (1 mmol) was dissolved in MoeMIM.Ms (1 ml) and catalytic amount of 4-DMAP was added, followed by the addition of BzCN (2.5-3.5 mmol). The reaction mixture was stirred at room temperature ($25-30^{\circ}$ C) and the progress monitored by TLC. On completion of the reaction, water



Starting compd.	BzCN (equiv.)	Reaction time	Product(s)	Yield (%)*	HRMS Calcd./Obs.
la	2.5	1.5 h	3 a	92	[M + Na] ⁺ 473.1325/473.1349
1 b	2.5	2.0 h	3 b	91	$[M + Na]^+$ 482.1440/482.1441
1 c	2.5	2.5 h	3 c	68	$[M + H]^+$ 436.1509/436.1511
			3 d	10	$[M + H]^+$ 540.1771/540.1804
1 d	3.0	2.5 h	3 f	65	$[M + H]^+$ 476.1570/476.1601
2 a	3.5	1.5 h	4a	96	[M + Na] ⁺ 593.1536/593.1604
2 b	3.0	1.5 h	4 b	85	$[M + H]^+$ 580.1832/580.1878
2c	3.0	2.0 h	4g	70	$[M + H]^+$ 556.1720/556.1751
			4d	12	$[M + Na]^+$ 682.1801/682.1901
2 d	3.5	2.5 h	4h	70	[M + Na] ⁺ 618.1601/618.1664

*Isolated yields.

(20 ml) was added and the product extracted with ethyl acetate (3×20 mL). The organic layer was separated, dried over sodium sulfate, concentrated, and the residue purified by silica gel column chromatography to afford the corresponding *O*-benzoylated nucleosides as white solids in 65–96% yields; minor amounts of perbenzoylated nucleosides were also obtained in case of 2'-deoxycytidine and cytidine (Figure 3).

The benzoylation of nucleosides of 2'-deoxy- and ribo-series with BzCN in IL is very selective and efficient as this combination exclusively/preferably benzoylated the hydroxyl groups of the nucleosides over their amino group (Figure 3). One of the major advantages of using IL in organic synthesis is that they are reusable. To verify the reusability, the IL was recovered from the aqueous portion of the reaction and used up to three cycles without any appreciable loss of its activity, except small loss of the IL in handling. The structures of all benzoylated nucleosides were unambiguously established on the basis of their spectral analysis (IR, ¹H and ¹³C NMR, and LC-MS) and comparison of their melting points and/or spectral data with those reported in the literature (Ref. [7] and references therein, and Ref. [8]) (Table 2).

CONCLUSION

The benzoylation reaction carried out on 2'-deoxy- and ribo-series of nucleosides with BzCN in pyridine was non-selective in most of the cases and required longer reaction times, elevated temperature and led to the formation of corresponding perbenzoylated products. Whereas, the BzCN in MoeMIM.Ms (IL) selectively/ preferentially benzoylated the hydroxyl groups over amino groups of nucleosides at ambient temperature (25–30°C). Also we have demonstrated that IL was recovered after benzoylation reactions. In conclusion, the ionic liquid MoeMIM.Ms is found to be an improved and "greener" alternative to the hazardous conventional solvents such as pyridine commonly employed in nucleoside chemistry.

TABLE 2

A. K. Prasad et al.

REFERENCES

- 1. Greene, T.W.; Wuts, P.G.M. Protective Groups in Organic Synthesis, 3rd Ed.; Wiley: New York, 1999; 173.
- Holy, A.; Soucek, M. Benzoyl cyanide–a new benzoylating agent in nucleoside and nucleotide chemistry. Tetrahedron Lett. 1971, 185–188.
- Garcia, J.; Fernandez, S.; Ferrero, M.; Sanghvi, Y.S.; Gotor, V.A. Mild, efficient and regioselective enzymatic procedure for 5'-O-benzoylation of 2'-deoxynucleosides. Tetrahedron Lett. 2004, 45, 1709–1712.
- Bhat, B.; Sanghvi, Y.S. A mild and highly selective N-benzoylation of cytosine and adenine bases in nucleosides with N-benzoyltetrazole. Tetrahedron Lett. 1997, 38, 8811–8814.
- Uzagare, M.C.; Sanghvi, Y.S.; Salunkhe, M.M. Application of ionic liquid 1-methoxyethyl-3-methylimidazolium methanesulphonate in nucleoside chemistry. Green Chem. 2003, 5, 370–372.
- Prasad, A.K.; Wengel, J. Enzyme-mediated protecting group chemistry on the hydroxyl groups of nucleosides. Nucleosides Nucleotides 1996, 15, 1347–1359.
- Kumar, V.; Maity, J.; Wang, Z.; Ravikumar, V.T.; Parmar, V.S.; Sanghvi, Y.S.; Prasad, A.K. Benzoyl cyanide: a mild and efficient reagent for benzoylation of nucleosides. Synth. Commun. 2005, 35, in press.
- Fox, J.J.; Wempen, I.; Hampton, A.; Doerr, I.L. Thiation of nucleosides. I. Synthesis of 2-amino-6-mercapto-9-β-D-ribofuranosylpurine ("Thioguanosine") and related purine nucleosides. J. Am. Chem. Soc. 1958, 80, 1669– 1675.