This article was downloaded by: [Korea University] On: 08 January 2015, At: 11:36 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <a href="http://www.tandfonline.com/loi/lncn20">http://www.tandfonline.com/loi/lncn20</a>

# Synthesis of Some New 2-a-L-Arabinopyranosyl-1,2,4triazines as Potential Antitumor Chemotherapeutics

Abdel Kader Mansour <sup>a c</sup> , Mohga M. Eid <sup>a</sup> & Nasser S. A. M. Khalil <sup>b</sup>

<sup>a</sup> Department of Chemistry, Faculty of Science, Cairo University, Giza, Egypt

<sup>b</sup> Central Laboratory for Food and Feed, Agricultural Research Center, Giza, Egypt

<sup>c</sup> Department of Chemistry , Faculty of Science , Cairo University , Giza, Egypt Published online: 07 Feb 2007.

To cite this article: Abdel Kader Mansour, Mohga M. Eid & Nasser S. A. M. Khalil (2003) Synthesis of Some New 2-a-L-Arabinopyranosyl-1,2,4-triazines as Potential Antitumor Chemotherapeutics, Nucleosides, Nucleotides and Nucleic Acids, 22:9, 1805-1823, DOI: <u>10.1081/NCN-120023274</u>

To link to this article: <u>http://dx.doi.org/10.1081/NCN-120023274</u>

# PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

# Synthesis of Some New 2-α-L-Arabinopyranosyl-1,2,4triazines as Potential Antitumor Chemotherapeutics

Abdel Kader Mansour,<sup>1,\*</sup> Mohga M. Eid,<sup>1</sup> and Nasser S. A. M. Khalil<sup>2</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science, Cairo University, Giza, Egypt <sup>2</sup>Central Laboratory for Food and Feed, Agricultural Research Center, Giza, Egypt

# ABSTRACT

Synthetic routes towards different 2- $\alpha$ -L-arabinopyranosyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-/ones or thiones were investigated. Primary human anticancer screening of two selected compounds resulted in an active compound against SF-268 (CNS) cell line.

Key Words: Synthesis; 1,2,4-Triazines; a-L-Arabinopyranosyls; Antitumors.

# **INTRODUCTION**

Many 1,2,4-triazine glycosides were reported to have pronounced biological activity.<sup>[1]</sup> Only little attention had been directed towards the synthesis of unnatural L-nucleosides until the finding that some  $\beta$ -L-nucleosides<sup>[2–5]</sup> were found to be more active and less toxic than their D-isomers<sup>[5,6]</sup> against HIV-1 and Hepatitis B virus

1805

DOI: 10.1081/NCN-120023274 Copyright © 2003 by Marcel Dekker, Inc.

Downloaded by [Korea University] at 11:36 08 January 2015

1525-7770 (Print); 1532-2335 (Online) www.dekker.com

<sup>\*</sup>Correspondence: Abdel Kader Mansour, Department of Chemistry, Faculty of Science, Cairo University, Giza, Egypt.

(HBV).<sup>[7,8]</sup> Biochemically, some L-nucleosides are substrates for cellular kinases<sup>[9,10]</sup> and also have greater stability for catabolizing enzymes such as cytidine and adenosine deaminase,<sup>[11]</sup> thereby providing higher anti HIV and anti HBV activities.<sup>[2]</sup> Recently, we have synthesized many 1,2,4-triazine galactosides<sup>[1]</sup> in order to find new antitumor agents. In this context we found that 2-(2,3,4,6-tetra-*O*-acetyl- $\beta$ -D-galactopyranosyl)-6-benzyl-1,2,4-triazin-3,5(2*H*,4*H*)-dithione (1) showed a primary an in vitro activity against both MCF7 (Breast) and SF-268 (CNS) cell lines. In continuation of our studies, it was of interest to synthesize some new  $\alpha$ -L-arabinopyranosides as promising antitumor chemotherapeutics. Therefore, herein we wish to report the synthesis of some new 2- $\alpha$ -L-arabinosyl derivaties of 3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-/ones or thiones and the primary human anticancer screening of two selected nucleosides.



### **RESULTS AND DISCUSSION**

Base induced glycosidation<sup>[1,12–17]</sup> of 4-amino-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-ones followed by deamination was reported to offer a convenient selective synthesis of the 2-glycosyl derivatives.<sup>[1,14–16]</sup> Also, glycosidation of the 4-aryl-3thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-ones was reported to give the corresponding 2-glycosyl derivatives.<sup>[1,17]</sup> In the present report we applied these strategies to selectively synthesize some new 2- $\alpha$ -L-arabinopyranosyl-6-substituted-3-thioxo-2,3dihydro-1,2,4-triazin-5(4*H*)-ones.

Glycosidation of the 4-amino-6-substituted-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-ones **4a-g** with 2,3,4-tri-*O*-acetyl- $\beta$ -L-arabinopyranosyl bromide **(5)** in acetonitrile containing triethylamine gave the corresponding 4-amino-2-(2,3,4-tri-*O*-acetyl- $\alpha$ -L-arabinopyranosyl)-6-substituted-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-ones**6a-g**. Among the different possible monoarabinosyl derivatives **6–8**, the structure of **6a-g** is clearly assigned by spectral and chemical evidences. Thus, the position of the anomeric proton at  $\delta$  6.82–6.67 with a coupling constant value of 8.7–9.4 Hz proves that the anomeric proton is in a trans position with respect to

#### Synthesis of Some New 2-a-L-Arabinopyranosyl-1,2,4-triazines

Downloaded by [Korea University] at 11:36 08 January 2015

the proton on position 2 of the L-arabinosyl ring which comes in agreement to the  $\alpha$ -*N*-arabinopyranosyls **6a-g** and not their isomeric  $\alpha$ -*S*-derivatives (this is consistent with reported positions and coupling constant values of the anomeric protons of *N*-type of different pyranosyls).<sup>[1,13,14,16,17]</sup> The presence of the NH<sub>2</sub> protons signal of  $\delta$  6.51–6.32 excludes the formation of the  $\alpha$ -4-NH-pyranosyls **8a-g**.

Analogous arabinopyrnosylation of the 4-aryl-6-substituted-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-ones **10a-i** gave the corresponding 4-aryl-2-(2,3,4-tri-*O*acetyl- $\alpha$ -L-arabinopyranosyl)-6-substituted-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)ones (**11a-i**). Assignment of the structure of compounds **11a-i** rather than their isomeric *S*-arabinopyranosyl derivatives **12a-i** was established spectroscopically. Thus, the <sup>1</sup>HNMR spectral data of compounds **11a-i** revealed the position of the anomeric protons at  $\delta$  6.78–6.69 ( $J_{H-1'-H-2'} = 8.4-9.4$  Hz) in agreement with that reported for analogous 4-aryl-2-(2,3,4,6-tetra-*O*-acetyl- $\beta$ -D-gluco<sup>[14,17]</sup>/or galactopyranosyl<sup>[1]</sup>)-6-substituted-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-ones.

Deamination of **6d-g** into the target compounds **13a-d** was achieved in almost quantitative yields by the action of nitrous acid in acetic acid. The structure of compounds **13a-d** was inferred from their chemical and spectral evidences. Thus, the <sup>1</sup>H NMR spectra of compounds **13a-d** showed the absence of NH<sub>2</sub> signal at  $\delta$  6.51–6.32 and appearance of NH signal at  $\delta$  10.76–9.32 (brs, 1H, exchangeable). Thiation of the latter compounds with phosphorous pentasulphide in pyridine afforded the corresponding 2-(2,3,4-tri-*O*-acetyl- $\alpha$ -L-arabinopyranosyl)-6-substituted-1,2,4-triazine-3,5(2*H*, 4*H*)-dithiones **14a-d**. Structure assignment of compounds **14a-d** was inferred from their correct analytical and spectral data. Thus, the IR spectra of these compounds showed the absence of the amide carbonyl function at 1716–1713 cm<sup>-1</sup>. Also, the <sup>1</sup>H NMR spectra of compounds **14a-d** revealed signals consistent with their structures.

Condensation of compounds **6d-g** with different aldehydes gave the corresponding 4-arylideneamino derivatives **15a-k**. The structure of **15a-k** was inferred from their analytical, chemical, and spectral data. Thus, the <sup>1</sup>H NMR spectra of these compounds revealed N=CH signal at  $\delta$  8.72–8.13 consistent with similar reported data.<sup>[1,14–16]</sup>

Heating of compound **15b** under reflux for 30 minutes with benzaldehyde afforded compound **13b** via the elimination of benzonitrile.

Treatment of the appropriate acetyl derivatives **6f**, **13a-d** with methanolic ammonia led to the formation of the corresponding free L-arabinopyranosyl nucleosides **16**, **17a-d** respectively.

# **BIOLOGICAL EVALUATION**

Compounds **6g**, and **15j** were tested for their human anticancer activity using an in vitro model through a 3-cell line, one-dose primary anticancer assay consisting of MCF7 (Breast), NCI-H460 (Lung), SF-268 (CNS). Only compound **15j** was found to be active against SF-268 (CNS) in this 3-cell line one dose primary human anticancer assay. The following table shows the activity of the previously mentioned compounds.

Compound	Concentration	Growth Percentage			
		(Lung) NCI-H460	(Breast) MCF7	(CNS) SF-268	Activity
6g 15j	1.00 E–0.04 M 1.00 E–0.04 M	67 69	71 52	48 28	Inactive Active

Table 1. Human anticancer activity of compounds 6g, 15j.

# **EXPERIMENTAL**

All melting points are uncorrected. IR spectra were recorded on a Perkin-Elmner 1430 spectrophotometer. <sup>1</sup>H NMR spectra were recorded at 200 MHz with a Varian GEMINI 2000 spectrometer. Element analyses were carried out at the microanalytical Center, Cairo University. Anticancer screening of compounds **6g**, **15j** was carried out at the National Cancer Institute – National Institutes of Health, Bethesda, Maryland, United States of America. The starting 4-amino/4-aryl-6substituted-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4H)-ones **4a-g**<sup>[18–20]</sup>/**10a-i**,<sup>[1,21]</sup> and 2,3,4-tri-*O*-acetyl- $\beta$ -L-arabinopyranosyl bromide (**5**)<sup>[22]</sup> were prepared as reported.

4-Amino-2-(2,3,4-tri-O-acetyl- $\alpha$ -L-arbinopyranosyl)-6-substituted-3-thioxo-2,3dihydro-1,2,4-triazin-5(4*H*)-ones (6a-g). General Procedure: 2,3,4-tri-O-acetyl- $\beta$ -Larabinopyranosyl bromide (5) (10 mmol) was added to a solution of each of compounds 4a-g (10 mmol) in acetonitrile (5 mL) and triethylamine (2 mL, 14 mmol). After shaking the reaction mixture for 20 minutes, it was left overnight at room temperature. The next day, the reaction mixture was cooled in an ice bath, acidified with acetic acid (1 mL) and diluted with cold water. The precipitate was then collected by filtration, dried at room temperature, and recrystallized from dichloromethane/ petroleum ether (bp. 40–60°C) as brown crystals of compounds 6a-c or from dichloromethane/methanol as yellow crystals of compounds 6d-g.

**4-Amino-2-(2,3,4-tri-***O***-acetyl**-*α***-L-arbinopyranosyl)-6-phenyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)<b>-one (6a).** Using the general procedure, **4a** gave **6a** (46%); mp. 132°C; IR (KBr) 3294, 3209 (NH<sub>2</sub>), 1747 (C=O acetate), 1689 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.2–7.44 (m, 5H, ArH'), 6.82 (d, 1H,  $J_{H-1'-H-2'} = 9$  Hz, H-1'), 6.51 (s, 2H, NH<sub>2</sub>, exchangeable), 6.08 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.5$  Hz, H-2'), 5.4 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.3 (dd, 1H,  $J_{H-3'-H-4'} = 3.4$  Hz,  $J_{H-5'} = 10$  Hz, H-3'), 4.16 (dd, 1H,  $J_{H-5'-H-4'} = 1.9$  Hz,  $J_{H-5''-H-5''} = 13.5$  Hz, H-5'), 3.94 (d, 1H,  $J_{H-5''-H-5''} = 13.4$  Hz, H-5''), 2.19, 2.06, 1.98 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>20</sub>H<sub>22</sub>N<sub>4</sub>O<sub>8</sub>S: C, 50.2; H, 4.63; N, 11.71. Found; C, 50.2; H, 4.7; N, 12.0.

4-Amino-2-(2,3,4-tri-*O*-acetyl-α-L-arbinopyranosyl)-6-benzyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one (6b). Using the general procedure, 4b gave 6b (62%); mp. 110°C; IR (KBr) 3308, 3215 (NH<sub>2</sub>), 1747 (C=O acetate), 1689 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.42–7.2 (m, 5H, ArH's), 6.67 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'),



Scheme 1.





6.32 (s, 2H, NH<sub>2</sub>, exchangeable), 6.01 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.38 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.25 (dd, 1H,  $J_{H-3'-H-4'} = 3.6$  Hz,  $J_{H-3'-H-2'} = 10$  Hz, H-3'), 4.13 (dd, 1H,  $J_{H-5'-H-4'} = 1.9$  Hz,  $J_{H-5'-H-5''} = 13.5$  Hz, H-5'), 4.14–3.85 (m, 2H, PhC<u>H</u><sub>2</sub>), 3.95 (d, 1H,  $J_{H-5''-H-5'} = 13.6$  Hz, H-5''), 2.22, 2.05, 1.91 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>21</sub>H<sub>24</sub>N<sub>4</sub>O<sub>8</sub>S Calcd.: C, 51.21; H, 4.91 Found: C, 51.3; H, 4.9.

4-Amino-2-(2,3,4-tri-*O*-acetyl-α-L-arbinopyranosyl)-6-(4-chlorobenzyl)-3-thioxo-2,3dihydro-1,2,4-triazin-5(4*H*)-one (6c). Using the general procedure, 4c gave 6c (68%); mp. 88°C; IR (KBr) 3113, 3219 (NH<sub>2</sub>), 1749 (C=O acetate), 1693 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.36–7.26 (2d, 4H, ArH's), 6.67 (d, 1H,  $J_{H-1'-H-2'} = 9.4$  Hz, H-1'), 6.33 (s, 2H, NH<sub>2</sub>, exchangeable), 5.94 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.5$  Hz, H-2'), 5.4 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.25 (dd, 1H,  $J_{H-3'-H-4'} = 3.4$  Hz,  $J_{H-3'-H-2'} = 10.0$  Hz, H-3'), 4.14 (dd, 1H,  $J_{H-5'-H-4'} = 1.7$  Hz,  $J_{H-5'-H-5''} = 13.2$  Hz, H-5'), 4.1–3.87 (m, 2H, 4-ClC<sub>6</sub>H<sub>4</sub>C<u>H<sub>2</sub></u>), 3.9 (d, 1H,  $J_{H-5''-H-5'} = 13.4$  Hz, H-5''), 2.2, 2.05, 1.92 (3s, 9H, CH<sub>3</sub>CHO).

Anal. Calcd. for C<sub>21</sub>H<sub>23</sub>N<sub>4</sub>O<sub>8</sub>SCL: C, 47.87; H, 4.4. Found: C, 47.9; H, 4.5.

**4-Amino-2-(2,3,4-tri-***O***-acetyl-α-L-arbinopyranosyl)-6-styryl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (6d). Using the general procedure, 4d gave 6d (64%); mp. 180°C; IR (KBr) 3302, 3202 (NH<sub>2</sub>), 1747 (C=O acetate), 1680 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.04, 7.16 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.65–7.36 (m, 5H, ArH's), 6.69 (d, 1H, J\_{H-1'-H-2'} = 9.2 Hz, H-1') 6.40 (s, 2H, NH<sub>2</sub>, exchangeable), 6.05 (t, 1H, J = (J\_{H-2'-H-1} + J\_{H-2'-H-3'})/2 = 9.6 Hz, H-2'), 5.42 (d, 1H, J\_{H-4'-H-3'} = 3.4 Hz, H-4'), 5.28 (dd, 1H, J\_{H-3'-H-4'} = 3.6 Hz, J\_{H-3'-H-2'} = 10.2 Hz, H-3'), 4.18 (dd, 1H, J\_{H-5'-H-4'} = 1.8 Hz, J\_{H-5'-H-5''} = 13.4 Hz, H-5'), 3.94 (d, 1H, J\_{H-5''-H-5'} = 13.4 Hz, H-5''), 2.25, 2.06, 1.92 (3s, 9H, CH<sub>3</sub>CO).** 

Anal. Calcd. for C<sub>22</sub>H<sub>24</sub>N<sub>4</sub>O<sub>8</sub>S: C, 52.37; H, 4.79. Found: C, 52.4; H, 4.8.

**4-Amino-2-(2,3,-tri-***O***-acetyI**-*α***-L-arabinopyranosyI)**-6-β-4-methylpheny**I)vinyI**-3-thioxo-2,3-dihydro-1,2,4,-triazin-5(4*H*)-one (6e). Using the general procedure, 4e gave 6e (77.0%); mp. 178°C; IR (KBr) 3306, 3209 (NH<sub>2</sub>), 1747 (C=O acetate), 1686 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.0, 7.11 (2d, 2H, J = 16.3 HZ, trans CH=CH), 7.51, 7.2 (2d, 4H, ArH's), 6.68 (d, 1H,  $J_{H-1'-H-2'} = 8.7$  Hz, H-1'), 6.43 (s, 2H, NH<sub>2</sub>,exchangeable), 6.05 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.3$  Hz, H-2'), 5.41 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H<sup>4</sup>), 5.27 (dd, 1H,  $J_{H-3'-H-4'} = 3.4$  Hz,  $J_{H-3'-H-2'} = 10.0$  Hz, H-3'), 4.16 (dd, 1H,  $J_{H-5''-H-4'} = 1.8$  Hz,  $J_{H-5''-H-5''} = 13.4$  Hz, H-5''), 23.8 (s, 3H, CH<sub>3</sub>), 2.24, 2.05, 1.95 (3s, 9H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>23</sub>H<sub>26</sub>N<sub>4</sub>O<sub>8</sub>S: C, 53.27; H, 5.05. Found: C, 53.3; H, 5.0.

**4-Amino-2-(2,3,4-tri-O-acetyl-α-L-arabinopyranosyl)-6-β-(4-methoxyphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4,-triazin-5(4H)-one (6f).** Using the general procedure, **4f** gave **6f** (74%); mp. 212°C; IR (KBr) 3308, 3209 (NH<sub>2</sub>), 1747 (C=O acetate), 1691 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.98, 7.03 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.57, 6.92 (2d, 4H, ArH's), 6.42 (s, 2H, NH<sub>2</sub>, exchangeable), 6.68 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 6.06 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.7$  Hz, H-2'), 5.41 (d, 1H,  $J_{H-4'-H-3'} = 3.6$  Hz, H-4'), 5.27 (dd, 1H,  $J_{H-3'-H-4'} = 3.6$  Hz,  $J_{H-3'-H-4'} = 3.6$  Hz,  $J_{\text{H-3'H-2'}} = 10.0 \text{ Hz}, \text{ H-3'}$ , 4.17 (dd, 1H,  $J_{\text{H-5'-H-4'}} = 1.6 \text{ Hz}, J_{\text{H-5'-H-5''}} = 14 \text{ Hz}, \text{ H-5'}$ ), 3.94 (d, 1H,  $J_{\text{H-5''-H-5'}} = 14 \text{ Hz}, \text{ H-5''}$ ), 3.85 (s, 3H, OCH<sub>3</sub>), 2.24, 2.05, 1.95 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>23</sub>H<sub>26</sub>N<sub>4</sub>O<sub>9</sub>S: C, 51.68; H, 4.9. Found: C, 51.7; H, 4.9.

**4-Amino-2-(2,3,4-tri-***O***-acety1**-*α***-L-arabinopyranosy1**)-6-β-(-4-chloropheny**1**)vinyl-**3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (6g). Using the general procedure, 4g gave 6g (76%); mp. 196°C; IR (KBr) 3319, 3214 (NH<sub>2</sub>), 1749, 1728 (C=O acetate), 1689 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.98, 7.12 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.37 (2d, 4H, ArH's), 6.69 (d, 1H.**  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 6.45 (s, 2H,NH<sub>2</sub>, exchangeable), 6.05 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'} = 3.5$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.5$  Hz.  $J_{H-3'-H-2'} = 10.1$  Hz, H-3'), 4.17 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5''-H-5''} = 13.4$  Hz, H-5'), 3.94 (d, 1H,  $J_{H-5''-H-5''} = 13.2$  Hz, H-5''), 2.25, 2.06, 1.96 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>22</sub>H<sub>23</sub>N<sub>4</sub>O<sub>8</sub>SCl: C, 49.03; H, 4.3. Found: C, 49.0; H, 4.2.

4-Aryl-6-substituted-2-(2,3,4-tri-*O*-acetyl- $\alpha$ -L-arabinopyranosyl-6-substituted-3thioxo-2,3-dihydro-1,2,4,-triazin-5(4*H*)-ones (11a-i). General Procedure: To a solution of each of 10a-i (10 mmol) in acetonitrile (5 mL) and triethylamine (2 mL, 14 mmol) was added 2,3,4-tri-*O*-acetyl- $\beta$ -L-arabino-pyranosyl bromide (5) (10 mmol). After shaking the reaction mixture for 20 minutes it was kept at room temperature overnight. The reaction mixture was then cooled, acidified with acetic acid (1 mL) and the formed preciptate was collected by filtration, washed with ater, and dried at room temperature. Compounds 11a-i were extracted from ethyl acetate and purified by preparative TLC plates (silica gel 60 GF<sub>254</sub>) using ethyl acetate as an eluent. The eluted compounds were then extracted from chloroform on a soxhlet extractor. The chloroform extracts were concentrated and diluted with petroleum ether (bp. 40–60°C). After collection of crude 11a-i by filtration, they were recrystallized from diethylether/petroleum ether (bp. 40–60°C). as pale crystals of 11a-g and orange yellow crystals of 11h,i.

**4-Phenyl-2-(2,3,4-tri-***O*-acetyl-α-L-arabinopyranosyl)-6-benzyl-3-thioxo-2,3-dihydro-1,2,4,-triazin-5(4*H*)-one (11a). Using the general procedure, 10a gave 11a (55%);  $R_f = 0.72$ ; mp. 104°C; IR (KBr) 1747 (C=O acetate), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.54–7.15 (m, 10H, ArH's), 6.76 (d, 1H,  $J_{H-1'-H-2'} = 9.0$  Hz, H-1'), 6.01 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.3$  Hz, H-2'), 5.38 (d, 1H,  $J_{H-4'-H-3'} = 3.0$  Hz, H-4'), 5.26 (dd, 1H,  $J_{H-3'-H-4'} = 3.0$  Hz,  $J_{H-3'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.2$  Hz, H-5'), 3.97 (s, 2H, PhC<u>H</u><sub>2</sub>), 3.89 (d, 1H,  $J_{H-5''-H-5'} = 13.2$  Hz, H-5''), 2.23, 2.07, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for  $C_{27}H_{27}N3O_8S$ : C, 58.58; H, 4.91;N,7.59. Found: C, 58.6; H, 5.0; N, 7.6.

**4-(4-Methylphenyl)-2-(2,3,4-tri-***O*-acetyl-α-L-arabinopyranosyl)-6-benzyl-3-thioxo-**2,3-dihydro-1,2,4-triazin-5(4H)-one (11b).** Using the general procedure, **10b** gave **11b** (68%);  $R_f = 0.74$ ; mp. 88°C; IR (KBr) 1747 (C=O acetate), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.45–6.98 (m, 9H, ArH's), 6.78 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, Hz, H-1'), 6.02 (t, 1H,  $J = (J_{H-2'-H-3'} + J_{H-2'-H-3'})/2 = 9.1$  Hz, H-2'), 5.39 (d, 1H, Synthesis of Some New 2-a-L-Arabinopyranosyl-1,2,4-triazines

Downloaded by [Korea University] at 11:36 08 January 2015

 $J_{\text{H-4'-H-3'}} = 3.3 \text{ Hz}, \text{ H-4'}$ , 5.26 (dd, 1H,  $J_{\text{H-3'-H-4'}} = 3.3 \text{ Hz}, J_{\text{H-3'-H-2'}} = 9.8 \text{ Hz}, \text{ H-3'}$ ), 4.11 (dd, 1H,  $J_{\text{H-5'-H-4'}} = 1.6 \text{ Hz}, J_{\text{H-5'-H-5''}} = 13.0 \text{ Hz}, \text{ H-5'}$ ), 3.98 (s, 2H, PhC<u>H</u><sub>2</sub>), 3.88 (d, 1H,  $J_{\text{H-5''-H-5'}} = 13.0 \text{ Hz}, \text{ H-5'}$ ), 2.42 (s,3H,4-C<u>H</u><sub>3</sub>-C<sub>6</sub>H<sub>4</sub>-N), 2.24, 2.08,2.00 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>28</sub>H<sub>29</sub>N<sub>3</sub>O<sub>8</sub>S: C, 59.25; H, 5.15. Found: C, 59.3; H, 5.2.

**4-Phenyl-2-(2,3,4-tri-***O*-acetyl-α-L-arabinopyranosyl)-6-(-4-methylbenzyl)-3-thioxo-**2,3-dihydro-1,2,4,-triazin-5(***4H***)-one (11c).** Using the general procedure, **10c** gave **11c** (82%);  $R_f = 0.7$ ; mp. 168°C; IR (KBr) 1747 (C=O acetate), 1709 (C=O amide) cm<sup>-1</sup>, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.2–7.07 (m, 9H, ArH's), 6.74 (d, 1H,  $J_{H-1'-H-2'} = 8.6$  Hz H-1'), 6.04 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 8.6$  Hz, H-2'), 5.41 (d, 1H,  $J_{H-4'-H-3'} = 3.0$  Hz, H-4'), 5.25(dd, 1H,  $J_{H-3'-H-4'} = 3.0$  Hz,  $J_{H-3'-H-2'} = 9.7$  Hz, H-3'), 4.14(dd, 1H,  $J_{H-5''-H-4'} = 1.6$  Hz,  $J_{H-5''-H-5''} = 13.0$  Hz, H-5'), 3.92 (s, 2H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.85 (d, 1H,  $J_{H-5''-H-5'} = 13.0$  Hz, H-5''), 2.32 (s, 3H, 4-C<u>H<sub>3</sub></u>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.22, 2.06, 1.99 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>28</sub>H<sub>29</sub>N<sub>3</sub>O<sub>8</sub>S: C, 59.25; H, 5.15. Found: C, 59.4; H, 5.1.

**4-(4-Methylphenyl)-2-(2,3,4-tri-***O***-acetyl**-*α*-L-**arabinopyranosyl)-6-(4-methylbenzyl)-3-thioxo-2,3-dihydro-1,2,4,-triazin-5(***4H***)-one (11d). Using the general procedure, <b>10d** gave **11d** (78%);  $R_f = 0.75$ ; mp. 118°C; IR (KBr) 1751 (C=O acetate), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.34–6.96 (m, 8H, ArH's), 6.76 (d, 1H,  $J_{H-1'-H-2'} = 8.7$  Hz, H-1'), 6.03 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 8.7$  Hz, H-2'), 5.38 (d, 1H,  $J_{H-4'-H-3'} = 2.9$  Hz, H-4'), 5.25(dd, 1H,  $J_{H-3'-H-4'} = 2.9$  Hz,  $J_{H-3'-H-2'} =$ 9.2 Hz, H-3'), 4.16(dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 12.75$  Hz, H-5'), 3.93 (s, 2H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.85 (d, 1H,  $J_{H-5''-H-5'} = 12.77$  Hz, H-5''), 2.4 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>-N), 2.33, (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.23, 2.07,2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>29</sub>H<sub>31</sub>N<sub>3</sub>O<sub>8</sub>S: C, 59.88; H, 5.37. Found: C, 60.0; H, 5.4.

**4-Phenyl-2-(2,3,4-tri-O-acetyl-α-L-arabinopyranosyl)-6-(4-methoxybenzyl)-3-thioxo-2,3-dihydro-1,2,4,-triazin-5(4***H***)-one (11e). Using the general procedure, 10e gave 11e (95%); R\_f = 0.73; mp. 138°C; IR (KBr) 1747 (C=O acetate), 1705 (C=O amide) cm<sup>-1</sup>, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.53–6.88 (m, 9H, ArH's), 6.77 (d, 1H, J\_{H-1'-H-2'} = 9.1 Hz, H-1'), 6.02 (t, 1H, J = (J\_{H-2'-H-1} + J\_{H-2'-H-3'})/2 = 9.8 Hz, H-2'), 5.38 (d, 1H, J\_{H-4'-H-3'} = 3.2 Hz, H-4'), 5.26 (dd, 1H, J\_{H-3'-H-4'} = 3.2 Hz, J\_{H-3'-H-2'} = 9.8 Hz, H-3'), 4.13 (dd, 1H, J\_{H-5'-H-4'} = 1.6 Hz, J\_{H-5'-H-5''} = 13 Hz, H-5'), 3.91 (s, 2H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.86 (d, 1H, J\_{H-5''-H-5'} = 13.1 Hz, 5″), 3.79 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>), 2.23, 2.07, 2.0 (3s, 9H, CH<sub>3</sub>CO).** 

Anal. Calcd. for C<sub>28</sub>H<sub>29</sub>N<sub>3</sub>O<sub>9</sub>S: C, 57.62; H, 5.01. Found: C, 57.6; H, 5.1.

**4-(4-Methylphenyl)-2-(2,3,4-tri-***O*-acetyl-α-L-arabinopyranosyl)-6-(4-methoxybenzyl)-**3-thioxo-2,3-dihydro-1,2,4,-triazin-5(4***H***)-one (11f). Using the general procedure, <b>10f** gave **11f** (85%);  $R_f = 0.71$ ; mp. 90°C; IR (KBr) 1747 (C=O acetate), 1705 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.35–6.85 (m, 8H, ArH's), 6.75 (d, 1H,  $J_{H-1'-H-2'} = 9.4$  Hz, H-1'), 6.01 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})$  /2 = 9.4 Hz, H-2'), 5.38 (d, 1H,  $J_{H-4'-H-3'} = 3.15$  Hz, H-4'), 5.25 (dd, 1H,  $J_{H-3'-H-4'} = 3.15$  Hz,  $J_{H-3'-H-2'} =$ 9.4 Hz, H-3'), 4.13 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.4$  Hz, H-5'), 3.88 (s, 2H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub><u>CH<sub>2</sub></u>), 3.77 (s, 3H, 4-C<u>H<sub>3</sub></u>OC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 3.86 (d, 1H,  $J_{H-5''-H-5'} = 13.4$  Hz, H-5''), 2.39 (s, 3H, 4-C<u>H<sub>3</sub></u>C<sub>6</sub>H<sub>4</sub>N), 2.21, 2.07, 2.09 (3s, 9H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>29</sub>H<sub>31</sub>N<sub>3</sub>O<sub>9</sub>S: C, 58.28; H, 5.23. Found: C, 58.3; H, 5.2.

**4-Phenyl-2-(2,3,4-tri***O*-acetyl-α-L-arabinopyranosyl)-6-(4-chlorobenzyl)-3-thioxo-**2,3-dihydro-1,2,4,-triazin-5(4***H***)-one (11g).** Using the general procedure, **10g** gave **11g** (81.3%);  $R_f = 0.73$ ; mp. 104°C; IR (KBr) 1751 (C=O acetate), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.54–7.15 (m, 9H, ArH's), 6.77 (d, 1H,  $J_{H-1'-H-2'} = 9.0$  Hz, H-1'), 5.91 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.4$  Hz, H-2'), 5.38 (d, 1H,  $J_{H-4'-H-3'} =$ 3.4 Hz, H-4'), 5.23 (dd, 1H,  $J_{H-3'-H-4'} = 3.4$  Hz,  $J_{H-3'-H-2'} = 10.0$  Hz, H-3'), 4.13(dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.4$  Hz, H-5'), 3.92 (s, 2H, 4-ClC<sub>6</sub>H<sub>4</sub>C<u>H</u><sub>2</sub>), 3.84 (d, 1H,  $J_{H-5''-H-5'} = 13.5$  Hz, H-5''), 2.2, 2.06, 1.99 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>27</sub>H<sub>26</sub>N<sub>3</sub>O<sub>8</sub>SCl: C, 55.15; H, 4.46. Found: C, 55.2; H, 4.4.

4-Phenyl-2-(2,3,4-tri-*O*-acetyl-α-L-arabinopyranosyl)-6-(4-*N*,*N*-dimethylaminobenzyl)-3-thioxo-2,3-dihydro-1,2,4,-triazin-5(4*H*)-one (11h). Using the general procedure, 10h gave 11h (60%);  $R_f = 0.70$ ; mp. 120°C; IR (KBr) 1747 (C=O acetate), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.52–6.7 (m, 9H, ArH's), 6.69 (d, 1H,  $J_{H-1'-H-2'} = 8.4$  Hz, H-1'), 6.04 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.48 (d, 1H,  $J_{H-4'-H-3'} = 3.6$  Hz, H-4'), 5.23 (dd, 1H,  $J_{H-3'-H-4'} = 3.0$  Hz,  $J_{H-3'-H-2'} = 10.2$  Hz, H-3'), 4.19 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5''-H-5''} = 12.6$  Hz, H-5'), 3.97 (d, 1H,  $J_{H-5''-H-5''} = 13.5$  Hz, H-5''), 3.86 (s, 2H, 4-NMe<sub>2</sub>C<sub>6</sub>H<sub>4</sub>C<u>H</u><sub>2</sub>), 2.92 (s, 6H, 4-N(C<u>H</u><sub>3</sub>) <sub>2</sub>-C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.14–1.95 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for  $C_{29}H_{32}N_4O_8S$ : C, 58.38; H, 5.40; N, 9.39. Found: C, 58.4; H, 5.4; N, 9.6.

**4-(4-Methylphenyl-2-(2,3,4-tri-***O*-acetyl-*α*-L-arabinopyranosyl)-6-(4-*N*,*N*-dimethylaminobenzyl)-3-thioxo-2,3-dihydro-1,2,4,-triazin-5(4*H*)-one (11i). Using the general procedure, **10i** gave **11i** (57.4%);  $R_f = 0.75$ ; mp. 100°C; IR (KBr) 1747 (C=O acetate), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.56–6.74 (m, 8H, ArH's), 6.72 (d, 1H,  $J_{H-1'-H-2'} = 8.7$  Hz, H-1'), 6.01 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.49 (d, 1H,  $J_{H-4'-H-3'} = 3.1$  Hz, H-4'), 5.23 (dd, 1H,  $J_{H-3'-H-4'} = 3.3$  Hz,  $J_{H-3'-H-2'} = 10.2$  Hz, H-3'), 4.17 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 12.5$  Hz, H-5'), 3.94 (d, 1H,  $J_{H-5''-H-5'} = 13.5$  Hz, H-5''), 3.86 (s, 2H, 4-NMe<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.93 (s, 6H, 4-N(CH<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>), 2.39 (s, 3H, 4-CH<sub>3</sub>-C<sub>6</sub>H<sub>4</sub>N),2.14–1.95 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>30</sub>H<sub>34</sub>N<sub>4</sub>O<sub>8</sub>S: C, 59.0; H, 5.61. N, 9.17. Found: C, 5.9; H, 5.9; N, 9.2.

2-(2,3,4-Tri-O-acetyl- $\alpha$ -L-arabinopyranosyl)-6-substituted-3-thioxo-2,3-dihydro-1,2,4,triazin-5(4H)-ones (13a-d). General procedure: To a solution of each of 6d-g (10 mmol) in acetic acid (100 mL) was added a solution of sodium nitrite (6g in 6 mL water) dropwise with stirring and cooling at 0°C over a period of one hour and the reaction mixture was then kept in the refrigerator overnight. The next day, the reaction mixture was diluted with an ice-water mixture (400 g) and the formed precipitate was collected by filtration and dried at room temperature. Crude 13a was recrystallized first from diluted ethanol then from (1:1) diethyl ether/petroleum petroleum ether (bp.  $40-60^{\circ}$ C). Crude **13b,d** were recrystallized from dichloromethane/ methanol, and crude **13c** was recrystallized from methanol. All compounds were isolated as yellow crystals of pure **13a-d**.

**2-(2,3,4-Tri-***O***-acetyl-α**-L-**arabinopyranosyl)-6-styryl-3-thioxo-2,3-dihydro-1,2,4,**triazin-5(4*H*)-one (13a). Using the general procedure, 6d gave 13a (59%); mp. 176°C; IR (KBr) 3230 (NH), 1749 (C=O acetate), 1716 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.32 (brs, 1H, NH, exchangeable),7.91, 7.09 (2d, 2H, J = 16.3 Hz, trans CH=CH), 7.58–7.35 (m, 5H, ArH's), 6.63 (d, 1H,  $J_{H-1'-H-2'} = 9.3$  Hz, H-1'), 5.96 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.9$  Hz, H-2'), 5.4 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.24 (dd, 1H,  $J_{H-3'-H-4'} = 3.3$  Hz,  $J_{H-3'-H-2'} = 9.3$  Hz, H-3'), 4.18 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5''-H-5''} = 13.2$  Hz, H-5'), 3.92 (d, 1H,  $J_{H-5''-H-5'} = 13.2$  Hz, H-5''), 2.25, 2.05, 1.95 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>22</sub>H<sub>23</sub>N<sub>3</sub>O<sub>8</sub>S:C, 53.98; H, 4.73; N, 8.58. Found: C, 54.1; H, 4.7; N, 8.6.

**2-(2,3,4-Tri-***O*-acetyl-α-L-arabinopyranosyl)-6-β-(4-methylphenyl)vinyl-3-thioxo-2,3dihydro-1,2,4,-triazin-5(4*H*)-one (13b). Using the general procedure, 6e gave 13b (77%); mp. 218°C; IR (KBr) 3215 (NH), 1745 (C=O acetate), 1715 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.2 (brs, 1H, NH, exchangeable), 8.0, 7.06 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.49, 7.24 (2d, 4H, ArH's), 6.64 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 5.98 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'} = 3.2$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.5$  Hz,  $J_{H-3'-H-2'} = 10.1$  Hz, H-3'), 4.19(dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.4$  Hz, H-5'), 3.93 (d, 1H,  $J_{H-5''-H-5'} = 13.2$  Hz, H-5'), 2.37 (s, 3H, CH<sub>3</sub>), 2.26, 2.06, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>23</sub>H<sub>25</sub>N<sub>3</sub>O<sub>8</sub>S: C, 54.86; H, 5.0. Found: C, 55.0; H, 5.0.

**2-(2,3,4-Tri-***O*-acetyl-α-L-arabinopyranosyl)-6-β-(4-methoxyphenyl)vinyl-3-thioxo-**2,3-dihydro-1,2,4,-triazin-5(4***H***)-one (13c).** Using the general procedure, **6f** gave **13c** (85.6%); mp. 138°C; IR (KBr) 3212 (NH), 1749 (C=O acetate), 1713 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.76 (s, 1H, NH, exchangeable), 7.87, (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.52, 6.89 (2d, 4H, ArH's), 6.62 (d, 1H,  $J_{H-1'-H-2'} = 9.4$  Hz, H-1'), 5.96 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.4$  Hz,  $J_{H-3'-H-2'} = 10.1$  Hz, H-3'), 4.16 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.8$  Hz, H-5'), 3.84 (s, 3H, OCH<sub>3</sub>), 2.25, 2.06, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>23</sub>H<sub>25</sub>N<sub>3</sub>O<sub>9</sub>S: C, 53.17; H, 4.84. Found: C, 53.2; H, 4.8.

**2-(2,3,4-Tri-***O*-acetyl-α-L-arabinopyranosyl)-6-β-(4-chlorophenyl)vinyl-3-thioxo-2,3dihydro-1,2,4,-triazin-5(4*H*)-one (13d). Using the general procedure, 6g gave 13d (75%); mp. 254°C; IR (KBr) 3165 (NH), 1743 (C=O acetate), 1713 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.75 (s, brs, 1H, NH, exchangeable), 7.99, 7.07 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.53, 7.37 (2d, 4H, ArH's), 6.62 (d, 1H,  $J_{H-1'-H-2'} = 9.4$  Hz, H-1'), 5.96 (t, 1H,  $J = (J_{H-2'-H-1} + J_{H-2'-H-3'})/2 = 9.5$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.27 (dd, 1H,  $J_{H-3'-H-4'} = 3.6$  Hz,  $J_{H-3'-H-2'} = 10.2$  Hz, H-3'), 4.19(dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.4$  Hz, H-5'), 3.92 (d, 1H,  $J_{H-5''-H-5'} = 13.8$  Hz, H-5″), 2.25, 2.06, 2.0 (3s, 9H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>22</sub>H<sub>22</sub>N<sub>3</sub>O<sub>8</sub>SCl: C, 50.43; H, 4.23;N,8.02. Found: C, 50.5; H, 4.2;N,7.9.

2-(2,3,4-Tri-O-acetyl- $\alpha$ -L-arabinopyranosyl)-6-substituted-1,2,4-triazine-3,5 (2H,4H)dithiones (14a-d). General procedure: To a solution of each of 13a-d (1 mmol) in dry pyridine (5 mL) was added phosphorous pentasulfide (0.45 g, 2 mmol). The reaction mixture was then heated under reflux for 6 hours. After cooling, ethanol (10 mL) was added, and the supernatent solution was decaned, acidified with acetic acid (0.5 mL), concentrated, and diluted with water. The precipitate was collected by filtration, dried at room temperature, dissolved in diethyl ether, and charcoal (0.5 g) was added, filtered, and after evaporation of the ether, the residue obtained was recrystallized from diethylether/petroleum ether (bp. 40–60°C) as yellow crystals of 14a-d.

**2-(2,3,4-Tri-***O***-acetyI**-*a***-L-arabinopyranosyI)**-6-styry**I-1,2,4-triazine-3,5(2***H***,4***H***)-dithione (14a). Using the general procedure, <b>13a** gave **14a** (51%); mp. 124°C; IR (KBr) 3209 (NH), 1749 (C=O acetate) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  8.04, 7.1 (2d, 2H, *J* = 16.4 Hz, trans CH=CH), 7.64–7.36 (m, 6H, NH, ArH's), 6.7 (d, 1H, *J*<sub>H-1'-H-2'</sub> = 9.4 Hz, H-1'), 6.1 (t, 1H, *J* = (*J*<sub>H-2'-H-1'</sub> + *J*<sub>H-2'-H-3'</sub>)/2 = 9.6 Hz, H-2'), 5.42 (d, 1H, *J*<sub>H-4'-H-3'</sub> = 3.4 Hz, H-4'), 5.28 (dd, 1H, *J*<sub>H-3'-H-4'</sub> = 3.4 Hz, *J*<sub>H-3'-H-2'</sub> = 10 Hz, H-3'), 4.17 (dd, 1H, *J*<sub>H-5'-H-4'</sub> = 1.6 Hz, *J*<sub>H-5'-H-5''</sub> = 13.4 Hz, H-5'), 3.92 (d, 1H, *J*<sub>H-5''-H-5'</sub> = 13.4 Hz, H-5''), 2.26, 2.07, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for  $C_{22}H_{23}N_3O_7S_2$ : C, 52.27; H, 4.58;N,8.31. Found: C, 52.3; H, 4.5; N, 8.4.

**2-(2,3,4-Tri-***O*-acetyl-α-L-arabinopyranosyl)-6-β-(4-methylphenyl)vinyl-1,2,4-triazine-3,5(2*H*,4*H*)-dithione (14b). Using the general procedure, 13b gave 14b (50%); mp. 136°C; IR (KBr) 3165 (NH), 1749 (C=O acetate), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.7–7.16 (m, 7H, NH, CH=CH, ArH's), 6.54 (d, 1H,  $J_{H-1'-H-2'}=9.2$  Hz, H-1'), 6.05 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.8$  Hz, H-2'), 5.4 (d, 1H,  $J_{H-4'-H-3'} =$ 3.4 Hz, H-4'), 5.23 (dd, 1H,  $J_{H-3'-H-4'} = 3.5$  Hz,  $J_{H-3'-H-2'} = 10.1$  Hz, H-3'), 4.17 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5'} = 13.4$  Hz, H-5'), 3.92 (d, 1H,  $J_{H-5'-H-5''} = 13.4$  Hz, H-5''), 2.37 (s, 3H, CH<sub>3</sub>), 2.23, 2.07, 2.01 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>23</sub>H<sub>25</sub>N<sub>3</sub>O<sub>7</sub>S<sub>2</sub>: C, 53.17; H, 4.85. Found: C, 53.2; H, 4.8.

**2-(2,3,4-Tri-***O*-acetyl-α-L-arabinopyranosyl)-6-β-(4-methoxyphenyl)vinyl-1,2,4-triazine-3,5(2*H*,4*H*)-dithione (14c). Using the general procedure, 13c gave 14c (52%); mp. 142°C; IR (KBr) 3200 (NH), 1749 (C=O acetate), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.64–6.88 (m, 7H, NH, CH=CH, ArH's),6.55 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 6.09 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.23 (dd, 1H,  $J_{H-3'-H-4'} = 3.4$  Hz,  $J_{H-3'-H-2'} = 10.1$  Hz, H-3'), 4.18 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.5$  Hz, H-5'), 3.93 (d, 1H,  $J_{H-5''-H-5'} = 13.4$  Hz, H-5''), 3.84 (s, 3H, OCH<sub>3</sub>), 2.23, 2.07, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C23H25N3O8S2: C, 51.58; H, 4.70. Found: C, 51.4; H, 4.8.

2-(2,3,4-Tri-*O*-acetyl- $\alpha$ -L-arabinopyranosyl)-6-β-(4-chlorophenyl)vinyl-1,2,4-triazine-3,5(2*H*,4*H*)-dithione (14d). Using the general procedure, 13d gave 14d (51%); Downloaded by [Korea University] at 11:36 08 January 2015

mp. 134°C; IR (KBr) 3165 (NH), 1749 (C=O acetate), cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  8.75 (brs, 1H, NH), 7.69–7.46 (m, 6H, CH=CH, ArH's), 6.72 (d, 1H,  $J_{\text{H-1'-H-2'}} = 9.2 \text{ Hz}$ , H-1'), 5.96 (t, 1H,  $J = (J_{\text{H-2'-H-1'}} + J_{\text{H-2'-H-3'}})/2 = 9.6 \text{ Hz}$ , H-2'), 5.47 (d, 1H,  $J_{\text{H-4'-H-3'}} = 3.4 \text{ Hz}$ , H-4'), 5.28 (dd, 1H,  $J_{\text{H-3'-H-4'}} = 3.3 \text{ Hz}$ ,  $J_{\text{H-3'-H-2'}} = 10.2 \text{ Hz}$ , H-3'), 4.08 (dd, 1H,  $J_{\text{H-5'-H-4'}} = 1.6 \text{ Hz}$ ,  $J_{\text{H-5''-H-5''}} = 13.6 \text{ Hz}$ , H-5'), 3.98 (d, 1H,  $J_{\text{H-5'-H-5'}} = 8.8 \text{ Hz}$ , H-5'), 2.16–1.92 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>22</sub>H<sub>22</sub>N<sub>3</sub>O<sub>7</sub>S<sub>2</sub>C1: C,48.93; H, 4.11. Found: C, 49.0; H, 4.1.

4-Arylideneamino-2-(2,3,4-tri-O-acetyl- $\alpha$ -L-arabinopyranosyl)-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4H)-ones (15a-k). General procedure: A mixure of each of 6d-g (1 g) and the appropriate aldehyde (1 mL or 1 g) was heated at 150–160°C in an oil bath for 15 minutes. This mixure was then heated under reflux in methanol for further 30 minutes, cooled, and the product was collected by filtration. Compounds 15a-k were recrystallized from methanol as yellow crystals.

**4-Benzylidineamino-2-(2,3,4-tri-***O***-acetyl-***α*-**L-arabinopyranosyl)-6-styryl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (15a). Using the general procedure, 6d gave 15a (87%); mp. 208°C; IR (KBr) 1743 (C=O acetate), 1697 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.39 (s, 1H, N=CH), 8.04, 7.19 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.97–7.37 (m, 10 H, ArH's),6.85 (d, 1H, J\_{H-1'-H-2'} = 9 Hz, H-1'), 6.09 (t, 1H, J = (J\_{H-2'-H-1'} + J\_{H-2'-H-3'})/2 = 9.8 Hz, H-2'), 5.42 (d, 1H, J\_{H-4'-H-3'} = 3.4 Hz, H-4'), 5.28 (dd, 1H, J\_{H-3'-H-4'} = 3.5 Hz, J\_{H-3'-H-2'} = 10.1 Hz, H-3'), 4.17 (dd, 1H, J\_{H-5''-H-4'} = 1.8 Hz, J\_{H-5''-H-5''} = 13 Hz, H-5'), 3.94 (d, 1H, J\_{H-5''-H-5'} = 13 Hz, H-5''), 2.26, 2.06, 2 0 (3s, 9H, CH<sub>3</sub>CO).** 

Anal. Calcd. for C<sub>29</sub>H<sub>28</sub>N<sub>4</sub>O<sub>8</sub>S: C,58.77; H, 4.76; N, 9.45. Found: C, 58.8; H, 4.6; N, 9.3.

**4-Benzylidineamino-2-(2,3,4-tri-O-acetyl-α-L-arabinopyranosyl)-6-β-(4-methylphenyl) vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(***4H***)-one (15b).** Using the general procedure, **6e** gave **15b** (78%); mp. 216°C; IR (KBr) 1747 (C=O acetate), 1693 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.38 (s, 1H, N=CH), 8.0, 7.14 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.97–7.11 (m, 9H, ArH's), 6.85 (d, 1H,  $J_{H-1'-H-2'} =$ 9.3 Hz, H-1'), 6.09 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.41 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.4$  Hz,  $J_{H-3'-H-2'} = 10$  Hz, H-3'), 4.17 (dd, 1H,  $J_{H-5'-H-4'} = 1.8$  Hz,  $J_{H-5'-H-5''} = 13.2$  Hz, H-5'), 3.94 (d, 1H,  $J_{H-5''-H-5'} =$ 13.2 Hz, H-5''), 2.37 (s, 3H, CH<sub>3</sub>), 2.26, 2.07, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>30</sub>H<sub>30</sub>N<sub>4</sub>O<sub>8</sub>S: C, 59.4; H, 4.98. Found: C, 59.4; H, 5.0.

**4-(4-Methylbenzylidineamino-2-(2,3,4-tri-***O***-acetyl-***α***-L-arabinopyranosyl)-6-β-(4methylphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (15c). Using the general procedure, <b>6e** gave **15c** (62%); mp. 238°C; IR (KBr) 1743 (C=O acetate), 1705 (C=O amide), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.32 (s, 1H, N=CH), 8.01, 7.14 (2d, 2H, J = 16.2 Hz, trans CH=CH), 7.86–7.18 (m, 8H, ArH's), 6.85 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 6.09 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.6$  Hz, H-2'), 5.36 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.5$  Hz,  $J_{H-3'-H-2'} =$ 10.1 Hz, H-3'), 4.18 (dd, 1H,  $J_{H-5'-H-4'} = 1.5$  Hz,  $J_{H-5'-H-5'} = 13.5$  Hz, H-5'), 3.94

1817

(d, 1H,  $J_{H-5''-H-5'} = 13.2$  Hz, H-5"), 2.45 (s, 3H, 4-CH<sub>3</sub>-C<sub>6</sub>H<sub>4</sub>CH=N), 2.38, (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH=CH), 2.26, 2.07, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>31</sub>H<sub>32</sub>N<sub>4</sub>O<sub>8</sub>S: C, 59.99; H, 5.2. Found: C, 60.0; H, 5.2.

4-(4-Chlorobenzylideneamino-2-(2,3,4-tri-*O*-acetyl-α-L-arabinopyranosyl)-6-β-(4methylphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one (15d). Using the general procedure, 6e gave 15d (72%); mp. 232°C; IR (KBr) 1743 (C=O acetate), 1697 (C=O amide), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.36 (s, 1H, N=CH), 8.0, 7.13 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.91–7.18 (m, 8H, ArH's), 6.83 (d, 1H,  $J_{H-1'-H-2'} =$ 9.2 Hz, H-1'), 6.08 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.4$  Hz, H-2'), 5.41 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.5$  Hz,  $J_{H-3'-H-2'} = 10$  Hz, H-3'), 4.18 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.3$  Hz, H-5'), 3.93 (d, 1H,  $J_{H-5''-H-5'} =$ 13.4 Hz, H-5''), 2.38 (s, 3H, CH<sub>3</sub>), 2.25, 2.26, 1.99 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>30</sub>H<sub>29</sub>N<sub>4</sub>O<sub>8</sub>SCl: C, 56.2; H, 4.56. Found: C, 56.1; H, 4.6.

4-(4-*N*,*N*-Dimethylaminobenzylideneamino-2-(2,3,4-tri-*O*-acetyl-α-L-arabinopyranosyl)-6-β-(4-methylphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one (15e). Using the general procedure, 6e gave 15e (80%); mp. 233°C; IR (KBr) 1743 (C=O acetate), 1697 (C=O amide), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.13 (s, 1H, N=CH), 8.01, 7.14 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.8–6.71 (m, 8H, ArH's), 6.87 (d, 1H,  $J_{H-1'-H-2'}=9.0$  Hz, H-1'), 6.09 (t, 1H,  $J = (J_{H-2'-H-1'}+J_{H-2'-H-3'})/2 = 9.5$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'}= 3.2$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'}= 3.5$  Hz,  $J_{H-3'-H-2'}= 9.1$  Hz, H-3'), 4.18 (dd, 1H,  $J_{H-5'-H-4'}= 1.6$  Hz,  $J_{H-5'-H-5'}= 13.4$  Hz, H-5'), 3.94 (d, 1H,  $J_{H-5''-H-5'}= 13.3$  Hz, H-5''), 3.1 (s, 6H, N(CH<sub>3</sub>) <sub>2</sub>), 2.37, (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH=CH), 2.25, 2.06, 1.98 (3s, 9H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>32</sub>H<sub>35</sub>N<sub>5</sub>O<sub>8</sub>S: C, 59.16; H, 5.43;N, 10.78. Found: C, 59.2; H, 5.3;N, 10.8.

**4-Benzylidineamino-2-(2,3,4-tri-***O*-acetyl-α-L-arabinopyranosyl)-6-β-(4-methylphyenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazine-5(4*H*)-one (15f). Using the general procedure, 6f gave 15f (75%); mp. 223°C; IR (KBr) 1751 (C=O acetate), 1689 (C=O amide), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.38 (s, 1H, N=CH), 7.99, 7.06 (2d, 2H, J = 16.4 Hz, trans CH=CH), 8.03–6.89 (m, 9H, ArH's), 6.85 (d, 1H,  $J_{H-1'-H-2'} =$ 9.2 Hz, H-1'), 6.09 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.7$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.5$  Hz,  $J_{H-3'-H-2'} = 10.1$  Hz, H-3'), 4.17 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5'} = 13.4$  Hz, H-5'), 3.94 (d, 1H,  $J_{H-5''-H-5'} =$ 13.6 Hz, H-5''), 3.85 (s, 3H, OCH<sub>3</sub>), 2.26, 2.06, 2.0 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>30</sub>H<sub>30</sub>N<sub>4</sub>O<sub>9</sub>S: C, 57.87; H, 4.86; Found: C, 57.9; H,4.9.

4-(4-Methylbenzylidineamino-2-(2,3,4-tri-*O*-acetyl-α-L-arabinopyranosyl)-6β-(4-methoxyphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one (15g). Using the general procedure, 6f gave 15g (69%); mp. 178°C; IR (KBr) 1743 (C=O acetate), 1697 (C=O amide), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.32 (s, 1H, N=CH), 7.98, 7.05 (2d, 2H, J=16.4 Hz, trans CH=CH), 7.85–6.89 (m, 8H, ArH's), 6.85 (d, 1H,  $J_{H-1'-H-2'}=9.2$  Hz, H-1'), 6.09 (t, 1H,  $J=(J_{H-2'-H-1'}+J_{H-2'-H-3'})/2=9.6$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'}=3.4$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'}=3.4$  Hz,  $J_{H-3'-H-2'}=$ 10.2 Hz, H-3'), 4.17 (dd, 1H,  $J_{H-5'-H-4'}=1.6$  Hz,  $J_{H-5'-H-5''}=13.4$  Hz, H-5'), 3.93 (d, 1H, *J*<sub>H-5"-H-5'</sub> = 13.6 Hz, H-5'), 3.84 (s, 3H, 4-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>CH=CH), 2.44 (s, 3H, 4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>CH=N), 2.25, 2.06, 1.99 (3s, 9H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>31</sub>H<sub>32</sub>N<sub>4</sub>O<sub>9</sub>S: C, 58.48; H, 5.06; Found: C, 58.5; H, 5.0.

**4-(4-Chlorobenzylidineamino-2-(2,3,4-tri**-*O*-acetyl-α-L-arabinopyranosyl)-6β-(4methoxyphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one (15h). Using the general procedure, 6f gave 15h (75%); mp. 206°C; IR (KBr) 1743 (C=O acetate), 1697 (C=O amide), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.36 (s, 1H, N=CH), 7.97, 7.04 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.91–6.89 (m, 8H, ArH's), 6.83 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 6.08 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.4$  Hz, H-2'), 5.41 (d, 1H,  $J_{H-4'-H-3'} = 3$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.6$  Hz,  $J_{H-3'-H-2'} =$ 10 Hz, H-3'), 4.17 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5'} = 13.4$  Hz, H-5'), 3.94 (d, 1H,  $J_{H-5''-H-5'} = 13.4$  Hz, H-5''), 3.84 (s, 3H, OCH<sub>3</sub>), 2.25, 2.06, 1.99 (3s, 9H, CH<sub>3</sub>CO). Anal. Calcd. for C<sub>30</sub>H<sub>29</sub>N<sub>4</sub>O<sub>9</sub>SCl: C, 54.84; H, 4.45. Found: C, 55.0; H, 4.4.

**4-(4-***N*,*N*-Dimethylaminobenzylidineamino-2-(2,3,4-tri-*O*-acetyl-α-L-arabinopyranosyl)-**6-β-(4-methoxyphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (15i). Using the general procedure, <b>6f** gave **15i** (75%); mp. 229–230°C; IR (KBr) 1743 (C=O acetate), 1697 (C=O amide), cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.14 (s, 1H, N=CH), 7.99, 7.06 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.82–6.86 (m, 8H, ArH's), 6.73 (d, 1H,  $J_{H-1'-H-2'} = 8.8$  Hz, H-1'), 6.1 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.7$  Hz, H-2'), 5.42 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz, H-4'), 5.28 (dd, 1H,  $J_{H-3'-H-4'} = 3.5$  Hz,  $J_{H-3'-H-2'} = 10.3$  Hz, H-3'), 4.18 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.6$  Hz, H-5'), 3.93 (d, 1H,  $J_{H-5''-H-5'} = 13.6$  Hz, H-5''), 3.85 (s, 3H, OCH<sub>3</sub>), 3.09 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 2.26, 2.06, 1.99 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>32</sub>H<sub>35</sub>N<sub>5</sub>O<sub>9</sub>S: C, 57.73; H, 5.3; N, 10.52. Found: C, 57.7; H, 5.2; N, 10.3.

**4-Benzylidineamino-2-(2,3,4-tri-***O***-acetyl-α-L-arabinopyranosyl)-6-β-(4-chlorophe-nyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (15j). Using the general procedure, <b>6g** gave **15j** (80%); mp. 240°C; IR (KBr) 1747 (C=O acetate), 1697 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.72 (s, 1H, N=CH),7.83, 7.17 (2d, 2H, J= 16.4 Hz, trans CH=CH), 7.99–7.51 (m, 9H, ArH's), 6.86 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 5.91 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.7$  Hz, H-2'), 5.49 (dd, 1H,  $J_{H-3'-H-4'} = 3.6$  Hz,  $J_{H-3'-H-2'} = 10.3$  Hz, H-3'), 5.3 (d, 1H,  $J_{H-4'-H-3'} = 3.8$  Hz, H-4'), 4.17 (dd, 1H,  $J_{H-5'-H-4'} = 1.5$  Hz,  $J_{H-5'-H-5'} = 13.4$  Hz, H-5'), 4.0 (d, 1H,  $J_{H-5''-H-5'} = 13.2$  Hz, H-5''), 2.2, 1.99, 1.98 (3s, 9H, CH<sub>3</sub>CO).

Anal. Calcd. for C<sub>29</sub>H<sub>27</sub>N<sub>4</sub>O<sub>8</sub>SCl: C, 55.55; H, 4.34. Found: C, 55.5; H, 4.4.

**4-(4-Methylbenzylidineamino)-2-(2,3,4-tri**-*O*-acetyl-α-L-arabinopyranosyl)-6-β-(4chlorophenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4*H*)-one (15k). Using the general procedure, **6g** gave 15k (80%); mp. 244°C; IR (KBr) 1747 (C=O acetate), 1697 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.65 (s, 1H, N=CH),7.83, 7.16 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.87–7.41 (m, 8H, ArH's), 6.86 (d, 1H,  $J_{H-1'-H-2'} = 9.2$  Hz, H-1'), 5.9 (t, 1H,  $J = (J_{H-2'-H-1'} + J_{H-2'-H-3'})/2 = 9.8$  Hz, H-2'), 5.48 (dd, 1H,  $J_{H-3'-H-4'} = 3.6$  Hz,  $J_{H-3'-H-2'} = 10$  Hz, H-3'), 5.3 (d, 1H,  $J_{H-4'-H-3'} = 3.4$  Hz,

1819



H-4'), 4.25 (dd, 1H,  $J_{H-5'-H-4'} = 1.6$  Hz,  $J_{H-5'-H-5''} = 13.6$  Hz, H-5''), 4.0 (d, 1H,  $J_{H-5''-H-5'} = 13.6$  Hz, H-5'), 2.43 (s, 3H, CH<sub>3</sub>), 2.19, 1.99, 1.97 (3s, 9H, CH<sub>3</sub>CO). Anal. Calcd. for  $C_{30}H_{29}N_4O_8SCl$ : C, 56.20; H, 4.56. Found: C, 56.2; H, 4.6.

Synthesis of Compound 13b from Compound 15b. A mixture of 15b (1.6 mmol) and benzaldehyde (1 mL) was heated under reflux for 30 minutes. After cooling and washing with cold methanol, the remaining solid was collected by filtration and recrystallized from methanol as yellow crystals of 13b (analytical and spectral data of 13b were mentioned before).

Action of Methanolic Ammonia on 6f and 13a-d. General procedure: A saturated methanolic ammonia solution (40 mol) (prepared by bubbing dry ammonia gas in absolute methanol at  $0^{\circ}$ C) was added to each of 6f, 13a-d (1 mmol). The reaction mixture was then left overnight at room temperature in a stoppered flask (after which time all materials went into solution). The solvent was then removed on rotavap at room temperature. Compounds 16, 17a-d were recrystallized from chloroform/ethanol as yellow crystals.

**4-Amino-2-α-L-arabinopyranosyl-6-β-(4-methoxyphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (16). Using the general procedure, <b>6f** gave 16 (75%); mp. 272°C (decomp.); IR (KBr) 3500–3200 (OH, NH<sub>2</sub>), 1689 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.89, 7.03 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.66, 7.01 (2d, 4H, ArH's), 6.76 (s, 2H, NH2, exchangeable), 6.25 (d, 1H,  $J_{H-1'-H-2'} = 8.8$  Hz, H-1'), 5.09, 5.02, 4.8 (3d, 3H, 3OH, exchangeable), 4.3–3.17 (m, 5H, H-2', H-3', H-4', H-5', H-5''), 3.82 (s, 3H, OCH<sub>3</sub>).

Anal. Calcd. for  $C_{17}H_{20}N_4O_6S$ : C, 49.99; H, 4.93; N, 13.72. Found: C, 50.0; H, 4.9 N, 13.6.

**2-a-L-Arabinopyranosyl-6-styryl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4H)-one (17a).** Using the general procedure, **13a** gave **17a** (71%); mp. 120°C (decomp.); IR (KBr) 3500–3200 (OH, NH), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  12.52 (brs, 1H, NH, exchangeable), 7.8, 7.08 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.67, 7.4 (m, 5H, ArH's), 6.22 (d, 1H,  $J_{\text{H-1'-H-2'}} = 8.8$  Hz, H-1'), 5.35, 5.07, 4.8 (3d, 3H, 3OH, exchangeable), 4.3–3.29 (m, 5H, H-2', H-3', H-4', H-5', H-5'').

Anal. Calcd. for C<sub>16</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>S: C, 52.88; H, 4.71; N, 11.56. Found: C, 52.8; H, 4.8 N, 12.0.

**2-α-L-Arabinopyranosyl-6-β-(4-methylphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (17b). Using the general procedure, 13b gave 17b (72%); mp. 130°C (decomp.); IR (KBr) 3500–3200 (OH, NH), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.27 (s, 1H, NH, exchangeable), 7.58, 7.03 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.51, 7.22 (2d, 4H, ArH's), 6.22 (d, 1H, J\_{H-1'-H-2'} = 9 Hz, H-1'), 5.34, 5.06, 4.8 (3d, 3H, 3OH, exchangeable), 4.3–3.0 (m, 5H, H-2', H-3', H-4', H-5', H-5''). 2.33 (s, 3H, CH<sub>3</sub>).** 

Anal. Calcd. for C<sub>17</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>S: C, 54.1; H, 5.07. Found: C, 54.1; H, 5.1.

**2-α-L-Arabinopyranosyl-6-β-(4-methoxylphenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (17c). Using the general procedure, <b>13c** gave **17c** (75%); mp. 112°C (decomp.); IR (KBr) 3500–3200 (OH, NH), 1697 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.38 (s, 1H, NH, exchangeable), 7.71, 6.99 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.65, 7.03 (2d, 4H, ArH's), 6.24 (d, 1H,  $J_{H-1'-H-2'} = 9.8$  Hz, H-1'), 5.36, 5.12, 4.82 (3d, 3H, 3OH, exchangeable), 4.3–3.3 (m, 5H, H-2', H-3', H-4', H-5', H-5''), 3.86 (s, 3H, OCH<sub>3</sub>).

Anal. Calcd. for C<sub>17</sub>H<sub>19</sub>N<sub>3</sub>O<sub>6</sub>S: C, 51.9; H, 4.87. Found: C, 52.0; H, 4.9.

**2-α-L-Arabinopyranosyl-6-β-(4-chlorophenyl)vinyl-3-thioxo-2,3-dihydro-1,2,4-triazin-5(4***H***)-one (17d). Using the general procedure, 13d gave 17d (73%); mp. 184°C IR (KBr) 3500–3200 (OH, NH), 1701 (C=O amide) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.3 (s, 1H, NH, exchangeable), 7.86, 7.05 (2d, 2H, J = 16.4 Hz, trans CH=CH), 7.67, 7.45 (2d, 4H, ArH's), 6.19 (d, 1H, J\_{H-1'-H-2'} = 8.7 Hz, H-1'), 5.32, 5.02, 4.91 (3d, 3H, 3OH, exchangeable), 4.3–3.32 (m, 5H, H-2', H-3', H-4', H-5', H-5'').** 

Anal. Calcd. for C<sub>16</sub>H<sub>16</sub>N<sub>3</sub>O<sub>5</sub>SCl: C, 48.3; H, 4.05. Found: C, 48.3; H, 4.1.

**Biological Evaluation of Compounds (6g, 15j).** An in vitro model was used as a primary human anticancer screen of compounds **6g, 15j**. A 3-cell line, one-dose assay consisting of MCF 7 (Breast), NCI-H460 (Lung), SF-268 (CNS) was used for the evaluation of the latter compounds. Each cell line was inoculated and preincubated on a microtiter plate. Test agents were then added at a single concentration and the culture were incubated for 48 hours. End point determination was made with sulforhodamine B, a protein-binding dye. Results for each test agent were reported as the percent of growth of the treated cells when compared to the untreated control cells. Compounds which reduce the growth of any one of the cell lines to 32% or less (negative number indicated cell kill) are active (Table 1).

# ACKNOWLEDGMENTS

The authors are very grateful to Dr. V.L. Narayanan and Dr. Edward Sausvile, National Cancer Institute/National Institutes of Health, Bethesda, Maryland, U.S.A, for their valuable anticancer testing.

# REFERENCES

- Mansour, A.K.; Eid, M.M.; Khahil, N.S.A.M. Selective synthesis and reactions of 6-substituted-2-β-galactosyl-1,2,4-triazines of potential anticancer activity. Nucleosides, Nucleotides & Nucleic Acids 2003, 22 (1), 21–44.
- Chang, C.N.; Doong, S.L.; Zhou, J.H.; Beach, J.W.; Jeong, L.S.; Chu, C.K.; Tsai, C.H.; Cheng, Y.C. Deoxycytidine deaminase – resistant stereoisomer is the active form of (±)-2',3'-dideoxy-3'-thiacytidine in the inhibitation of hepatitis B virus replication. J. Biol. Chem. 1992, 267, 13,938–13,942. Chem. Abstr. 1992, 117, 14,2910e.

- Belleau, B.; Dixit, D.; Nguyen-Ba, N.; Kraus, J.L. Purinyl- and pyrimidinyl-1,3-dioxolanes as anti-retrovirals. Eur. Pat. 337, 713, Oct 18, 1989. Chem. Abstr. 1990, 112 198359w.
- Furman, P.A.; Davis, M.; Liotta, D.C.; Paff, M.; Frick, L.W.; Nelson, D.J.; Dornsife, R.E.; Wurster, J.A.; Wilson, L.J.; Fyfe, J.A.; Tuttle, J.V.; Miller, W.H.; Condreay, L.; Avereet, D.R.; Schinazi, R.F.; Painter, G.R. The antihepatitis B virus activities, cytotoxicities, and anabolic profiles of the (-) and (+) enantiomers of cis-5-fluoro-1-[2-(hydroxymethyl)-1,3-oxothiolan-5-yl]cytosine. Antimicrob. Agents Chemother 1992, *36*, 2686–2692. Chem. Abstr. 1993, *118*, 93802f.
- 5. Schinazi, R.F.; Gosselin, G.; Faraj, A.; Korba, B.E.; Liotta, D.C.; Chu, C.K.; Mathe, C.; Imbach, J.; Sommadossi, J. Pure nucleoside enantiomers of  $\beta$ -2',3'-dideoxycytidine analogs are selective inhibitors of hepatitis B virus in vitro. Antimicrob. Agents Chemother **1994**, *38*, 2172–2174. Chem. Abstr. **1994**, *121*, 200749e.
- Coates, J.A.V.; Cammack, N.; Jenkinson, H.J.; Jowett, A.J.; Jowett, M.I.; Pearson, B.A.; Penn, C.R.; Rouse, P.L.; Viner, K.C.; Cameron, J.M. (-)-2'-Deoxy-3'-thiacytidine is a potent, highly selective inhibitor of human immunodeficiency virus type 1 and type 2 replication in vitro. Antimicrob. Agents Chemother 1992, 36, 733–739. Chem. Abstr. 1992, 116, 211044u.
- Doong, S.L.; Tsai, C.H.; Schinazi, R.F.; Liotta, D.C.; Cheng, Y.C. Inhibition of the replication of hepatitis B virus in vitro by 2',3'-dideoxy-3'-thiacytidine and related analog. Proc. Natl. Acad. Sci. USA 1991, 88, 8495–8499. Chem. Abstr. 1991, 115, 269987z.
- Chang, C.N.; Skalski, V.; Zhou, J.H.; Cheng, Y.C. Biochemical pharmacology of the (+)- and (-)-2',3'-dideoxy-3'-thiacytidine as antihepatitis B virus agents. J. Biol. Chem. 1992, 267, 22,414–22,420. Chem. Abstr. 1992, 117, 225781q.
- Grove, K.L.; Guo, X.; Liu, S.H.; Gao, Z.; Chu, C.K.; Cheng, Y.C. Anticancer activity of β-L-dioxolanecytidine, a novel nucleoside analog with the unnatural L configuration. Cancer Res. 1995, 55, 3008–3011. Chem. Abstr. 1995, 123, 132205p.
- Pai, S.B.; Liu, S.H.; Zhu, Y.L.; Chu, C.K.; Cheng, Y.C. Inhibition of hepatitis B virus by a novel L-nucleoside, 2'-fluoro-5-methyl-β-L-arabinofuranosyl uracil. Antimicrob. Agents Chemother 1996, 40, 380–386. Chem. Abstr. 1996, 124, 219475w.
- Pelicano, H.; Pierra, C.; Eriksson, S.; Gosselin, G.; Imbach, J.L.; Maury, G. Enzymatic properties of the unnatural β-L-enantiomers of 2',3'-dideoxyadenosine and 2',3'-didehydro-2',3'dideoxyadenosine. J. Med. Chem. 1997, 40, 3969– 3973. Chem Abstr. 1998, 128, 18411f.
- Mansour, A.K.; Ibrahim, Y.A.; Eid, M.M. Nucleoside derivatives of 1,2,4-triazine. Reaction of some derivatives of 1,2,4-triazine with tetra O-acetyle-α-Dglucopyranosyl bromide. Z. Naturforsch 1976, *31b*, 505–508.
- Mansour, A.K.; Ibrahim, Y.A.; Eid, M.M.; Abdel-Hady, S. A.L.N-Glucosyl derivatives of some 1,2,4-triazines with tetra-O-acetyl-α-D-glucopyranosyl bromide. J. Carbohydrates. Nucleosides. Nucleotides 1981, 8 (2), 81–99.
- 14. Ibrahim, Y.A. Facile approach for the selective glycosidation of cyclic asymmetric amides and thioamides. Carbohydrate Leters **1996**, *1*, 425–432.

### Synthesis of Some New 2-a-L-Arabinopyranosyl-1,2,4-triazines

- Ibrahim, Y.A. Synthesis and structure of 2-deoxyribosyl-6-azauracil derivatives. Carbohydrate Letters 1996, 2, 189–195.
- Mansour, A.J.; Ibrahim, Y.A.; Khalil, N.S.A.M. Selective synthesis and structure of 6-arylvinyl-2-and 4-glucosyl-1,2,4-triazines of expected interesting biological activity. Nucleosides & Nucleotides 1999, 18 (10), 2265–2283.
- 17. Eid, M.M.; Abdelhady, S.A.; Ali, H.A.W. Reaction of some 1,2,4-triazines with acetobromoglucose. Arch. Pharm. **1990**, *323* (4), 243–245.
- Dornow, A.; Menzel, H.; Marx, P. Synthesen sticktoffhaltiger heterocyclen, XXVII. Über 1.2.4-triazine, I. Darstellung einiger neuer s-triazolo [3.2-c]-astriazine. Chem. Ber. 1964, 97, 2173–2178.
- Ahmad, R.; Ajaz, S. Synthesis of 6-(p-Halobenzyl)-1,2,4-triazine-3,5(2H, 4H)dithiones having potential antimicrobial activity. J. Chem. Soc. Pak. 1985, 7 (4), 315–320.
- Badawy, M.A.; Abdel-Hady, S.A.L.; Eid, M.M.; Ibrahim, Y.A. Synthesis and structure of as-triazinoquinolines, II. Chem. Ber. 1984, 117, 1083–188.
- Mansour, A.K.; Eid, M.M.; Hassan, R.A.; Haemers, A.; Pattyn, S.R.; Vanden Berghe, D.A.; Van. Hoof, L. Synthesis, reactions and biological evaluation of some 1,2,3-triazine derivatives. J. Heterocycl. Chem. 1988, 25, 279–283.
- 22. Furniss, B.S.; Hannaford, A.J.; Rogers, V.; Smith, P.W.G.; Tatchell, A.R. *Vogel's Textbook of Practical Organic Chemistry*; Longman group Limited, 1978; Vol. 4, 458pp.

Received January 28, 2003 Accepted April 10, 2003

Copyright @ 2003 by Marcel Dekker, Inc. All rights reserved