



Synthesis and Crystal Structure of 4-Dichloroacetyl-3-phenyl-3,4-dihydro-2H-1,4-benzoxazine

YING FU, ZHAO-ZHENG XU and FEI YE*

College of Science, Northeast Agricultural University, Harbin 150030, HeilongJiang Province, P.R. China

*Corresponding author: Fax: +86 451 55191844; Tel: +86 451 55191854; E-mail: yefei@neau.edu.cn

Received: 26 June 2013;

Accepted: 4 September 2013;

Published online: 10 May 2014;

AJC-15147

A new compound 4-dichloroacetyl-3-phenyl-3,4-dihydro-2H-1,4-benzoxazine with m.f. $C_{16}H_{13}NO_2Cl_2$ was synthesized and characterized. The crystal structure of the compound has been determined by single-crystal X-ray diffraction. The crystal is of monoclinic, space group $P2_1/c$ with $a = 15.105(3)$, $b = 6.2685(13)$, $c = 16.084(3)$ Å, $\alpha = 90^\circ$, $\beta = 103.79(3)^\circ$, $\gamma = 90^\circ$, $V = 1479.1(5)$ Å³, $Z = 4$, $F(000) = 664$, $D_c = 1.447$ g/cm³, $\mu = 0.442$ mm⁻¹, the final $R_1 = 0.0507$ and $wR_2 = 0.1435$ for 2417 observed reflections with $I > 2\sigma(I)$. A total of 13533 reflections were collected, of which 3362 were independent ($R_{int} = 0.0307$). Benzoxazine ring and the benzene plane were almost perpendicular with the dihedral angle of 91.7° with each other.

Keywords: Benzoxazine, Synthesis, X-ray crystallography.

INTRODUCTION

Benzoxazines are promising agents as a starting material for the synthesis of large structures that have properties such as potassium channel activator or antibacterial agents^{1,2}. They were also been employed as the synthon of chiral compounds, pharmaceutical intermediates and polymer³⁻⁵. *N*-Dichloroacetyl benzoxazine is a significant safener for chlorinated amide herbicides and thiocarbamate herbicides. These aspects prompted us to synthesis, characteristics and biological activity of novel *N*-dichloroacetyl benzoxazine derivatives with better safety and efficacy, we introduced different groups substitution on the oxazine^{6,7}. As part of our search for bioactive compounds, here we report the synthesis and crystal structure of 4-dichloroacetyl-3-phenyl-3,4-dihydro-2H-1,4-benzoxazine.


EXPERIMENTAL

The melting point was measured on a Beijing Taike point apparatus (X-4) and uncorrected. The IR of KBr discs were

recorded on FTIR-8400S spectrometer. The ¹H NMR and ¹³C NMR spectrum was recorded on a Bruker AV-300 spectrometer in DMSO-*d*₆, internal standard TMS. The mass spectrum was recorded on a Xevo TQ spectrometer. The elemental analysis was performed on FLASH EA1112 elemental analyzer. All the reagents were analytical reagent grade.

Synthesis and characterization: 4-Dichloroacetyl-3-phenyl-3,4-dihydro-2H-1,4-benzoxazine (**3**) was prepared following **Scheme-I**.

Preparation of compound 2: Pt/C (2g) was added to a solution of 2-(*o*-nitrophenoxy)-phenylethanone (**1**) (0.05 mol, 12.9 g) in toluene (200 mL) and isopropanol (100 mL). The reaction mixture was stirred under H₂ at 60 °C at 1.5 Mpa for 10 h. Then the mixture was filtered. The toluene was removed under vacuum. The 3-phenyl-3,4-dihydro-2H-1,4-benzoxazine (**2**) was obtained as yellow oil with the yield of 93.3 %. Anal. calcd. for $C_{14}H_{13}NO$ (%): C, 79.58; H, 6.21; N, 6.63. Found: C, 79.52; H, 6.28; N, 6.59. IR (KBr, ν_{max} , cm⁻¹): 3363 (NH), 3060 (C-H), 1608, 1427 (C=C). ¹H NMR (300 M, DMSO-*d*₆):



Scheme-I: Synthetic route to the title compound

3.88-3.94, 4.20-4.24 (2H, m, O-CH₂); 4.46-4.48 (1H, t, *J* = 2.7 Hz, NCH); 6.25 (1H, s, NH); 6.50-6.55 (1H, m, H Ar); 6.69-6.71 (3H, m, H Ar); 7.22-7.46 (5H, m, H Ar); ¹³C NMR (75M, DMSO-*d*₆): 53.3, 70.4, 115.4, 116.2, 117.4, 121.7, 127.6, 127.6, 128.1, 128.9, 128.9, 135.4, 140.7, 143.2.

Preparation of compound 3: Dichloroacetyl chloride (17.6 nmol, 2.59 g) was added in a mixture containing 3-phenyl-3,4-dihydro-2*H*-1,4-benzoxazine (**2**) (14 nmol, 2.95 g), K₂CO₃ (15.2 nmol, 2.08 g) and benzene (25 mL). The reaction mixture was stirred at room temperature for 2 h. The organic phase was dried over anhydrous magnesium sulfate and the benzene was removed under vacuum, the 4-dichloroacetyl-3-phenyl-3,4-dihydro-2*H*-1,4-benzoxazine (**3**) was recrystallized from ethanol and light petroleum. The yield was 74.2 % (3.34 g); m.p. 136-138 °C. Anal. calcd. for C₁₆H₁₃NO₂Cl₂(%): C, 59.81; H, 4.08; N, 4.36. Found, %: C, 59.78; H, 4.16; N, 4.32. IR (KBr, ν_{max}, cm⁻¹): 3045 (C-H), 1689 (C=O), 1591, 1492 (C=C). ¹H NMR (300 M, DMSO-*d*₆): 4.47-4.52, 4.99-5.03 (2H, m, O-CH₂-C); 5.82 (1H, s, N-CH-); 6.79-7.75 (9H, m H Ar), 7.24 (1H, s, Cl₂CH); ¹³C NMR (75 M, DMSO-*d*₆): 66.8, 68.2, 117.4, 121.2, 124.0, 124.9, 127.0, 127.9, 127.9, 128.9, 128.9, 129.2, 129.2, 136.4, 147.1, 163.1. Mass spectrum: [M-1]⁺ = 321.

Crystal data and structure determination: The block-shaped single crystal of the title compound was grown by slow cooling from a hot saturated solution in ethanol. The crystal with dimensions of 0.62 × 0.23 × 0.20 mm was mounted on a Bruker AXSII CCD area-detector diffractometer using graphite monochromated Mo Kα radiation (λ = 0.71073 Å) at 293 (2) K. A total of 13533 reflections were collected in the range of 3.48 < θ < 27.48°, of which 3362 were independent and 2417 were observed with I > 2σ(I). The structure was solved by direct methods using SHELXS-97 program⁸ and the semi-empirical from equivalents absorption corrections were applied to all intensity data. All nonhydrogen atoms were refined an isotropically by the full-matrix least square method on F² using SHELXL-97⁹. The hydrogen atoms were determined with theoretical calculations and refined isotropically. A summary of the key crystallographic information is given in Table-1. The final full-matrix least squares refinement gave R = 0.0507 and wR = 0.1435 (w = 1/[σ²(F_o)² + (0.0912P)² + 0.1416 P]) where P = (F_o² + 2F_c²)/3, S = 1.122, (Δ/σ_{max} = 0.000, Δρ_{max} = 0.598 and Δρ_{min} = -0.445 eÅ⁻³). CIF file containing complete information on the studied structure was deposited with CCDC, deposition number 928571 and is freely available upon request from the following web site: www.ccdc.cam.ac.uk/data_request/cif.

RESULTS AND DISCUSSION

The structures of compound **2** and the title compound **3** were confirmed by elemental analysis, IR, ¹H NMR, ¹³C NMR, MS and X-ray diffraction analysis. The IR spectrum of the title compound shows absorption band at 1689 cm⁻¹ originated from the stretching vibration of C=O stretching vibration. The proton magnetic resonance spectra of compounds **2** and **3** have been recorded in DMSO-*d*₆. In the ¹H NMR spectrum the singlet at 3.8-5.8 ppm corresponds to the O-CH₂-CH-N proton indicating the formation of benzoxazine. The signals at 6.25 ppm range are N-H of compound **2**. And it is replaced by a

TABLE-1
CRYSTAL STRUCTURE AND DATA
REFINEMENT PARAMETERS

Empirical formula	C ₁₆ H ₁₃ Cl ₂ NO ₂
Formula weight	322.17
Crystal system, space group	Monoclinic, P2 ₁ /c
a (Å)	15.105(3)
b (Å)	6.2685(13)
c (Å)	16.084(3)
α (°)	90
β (°)	103.79(3)
γ (°)	90
V (Å ³)	1479.1(5)
Z	4
D _c (g/cm ³)	1.447
μ (mm ⁻¹)	0.442
F (000)	664
Crystal size (mm)	0.62 × 0.23 × 0.20
Color/shape	Colorless/block
Temp (K)	293(2)
Theta range for collection	3.48 - 27.48
Reflections collected	13533
Independent reflections	3362
Data/restraints/parameters	3362 / 0 / 190
Goodness-of-fit on F ²	1.122
Final R indices [I > 2σ(I)]	R ₁ = 0.0507, wR ₂ = 0.1435
R indices (all data)	R ₁ = 0.0691, wR ₂ = 0.1604
Largest difference peak/hole (eÅ ⁻³)	0.598 and -0.445

single sharp peak at 7.24 ppm for the Cl₂CH of the title compound. With regard to the mass spectra, the title compound show M-1 signals. The measured data of elemental analytical results is in accord with the theoretical values.

Crystal structure: The molecular structure with atom-numbering scheme and the packing diagram of the title compound are shown in Figs. 1 and 2, respectively. The selected bond lengths and bond angles are shown in Table 2.

According to X-ray analysis of the structure, the bond lengths of C(8)-N(1), C(1)-N(1) and C(7)-O(1) were close to the typical C-N and C-O bond length. The C(15)-O(2) (1.206 Å) indicated a double bond C=O (1.19-1.23 Å). However, compared with the normal aliphatic bond value, bond length C(15)-N(1) (1.359 Å) and C(6)-O(1) (1.370 Å) were shorter than normal.




Fig. 1. Molecular structure of 4-dichloroacetyl-3-phenyl-3,4-dihydro-2*H*-1,4-benzoxazine, showing displacement ellipsoids drawn at the 30 % probability level




Fig. 2. Packing of the title compound

TABLE-2
SELECTED BOND LENGTHS (Å) AND BOND ANGLES (°)

Atom	Distance	Atom	Angle
C(1)-C(6)	1.387(3)	C(6)-C(1)-C(2)	119.53(18)
C(1)-C(2)	1.394(3)	C(6)-C(1)-N(1)	118.30(17)
C(1)-N(1)	1.430(2)	C(2)-C(1)-N(1)	121.86(18)
C(2)-C(3)	1.372(3)	O(1)-C(6)-C(1)	122.98(18)
C(3)-C(4)	1.379(4)	O(1)-C(6)-C(5)	117.2(2)
C(4)-C(5)	1.383(4)	C(1)-C(6)-C(5)	119.8(2)
C(5)-C(6)	1.390(3)	O(1)-C(7)-C(8)	112.73(17)
C(6)-O(1)	1.370(3)	N(1)-C(8)-C(9)	112.41(16)
C(7)-O(1)	1.433(3)	N(1)-C(8)-C(7)	105.24(17)
C(7)-C(8)	1.523(3)	C(9)-C(8)-C(7)	115.24(17)
C(8)-N(1)	1.477(2)	C(14)-C(9)-C(10)	118.6(2)
C(8)-C(9)	1.518(3)	C(14)-C(9)-C(8)	121.58(19)
C(9)-C(14)	1.388(3)	C(10)-C(9)-C(8)	119.7(2)
C(9)-C(10)	1.393(3)	O(2)-C(15)-N(1)	123.31(19)
C(10)-C(11)	1.382(4)	O(2)-C(15)-C(16)	120.01(18)
C(11)-C(12)	1.357(5)	N(1)-C(15)-C(16)	116.53(17)
C(12)-C(13)	1.385(5)	C(15)-C(16)-Cl(1)	110.06(15)
C(13)-C(14)	1.388(3)	C(15)-C(16)-Cl(2)	106.64(14)
C(15)-O(2)	1.206(2)	Cl(1)-C(16)-Cl(2)	110.83(12)
C(15)-N(1)	1.359(3)	C(15)-N(1)-C(1)	127.04(16)
C(15)-C(16)	1.542(3)	C(15)-N(1)-C(8)	118.52(16)
C(16)-Cl(1)	1.756(2)	C(1)-N(1)-C(8)	113.17(16)
C(16)-Cl(2)	1.769(2)	C(6)-O(1)-C(7)	117.11(17)

It was obvious that there was $p\pi$ - $p\pi$ large conjunctive effect between O(1), benzene ring [C(1), C(3), C(4), C(5), C(6) and C(2)], N(1), C(15)-O(2) [C=O], which resulting in shorter bond length than the typical C-N bond length and C-O bond length. The title compound contained two benzene rings and an oxazine ring. The torsion angle of O(1)-C(7)-C(8)-N(1) was $-59.0(2)^\circ$, forming 7,8-half-chair conformation. Benzene ring plate I[C(1), C(3), C(4), C(5), C(6) and C(2)] made a dihedral angle of 2.6° with oxazine plane II[O(1), C(6), C(1), N(1), C(8), C(7)]. It means that they were almost coplanar because of the conjunctive effect. Benzoxazine ring and plan III[C(9), C(10), C(11), C(12), C(13) and C(14)] were almost perpendicular with the dihedral angle being 91.7° .

ACKNOWLEDGEMENTS

This work was supported by the National Nature Science Foundation of China (31101473), the Natural Science Foundation of Heilongjiang Province (B201212), the Science and Technology Research Project of Heilongjiang Education Department (12521015) and the Research Science Foundation in Technology Innovation of Harbin (2012RFQXN015).

REFERENCES

1. G. Caliendo, E. Perissutti, V. Santagada, F. Fiorino, B. Severino, D. Cirillo, R. d'Emmanuele di Villa Bianca, L. Lippolis, A. Pinto and R. Sorrentino, *Eur. J. Med. Chem.*, **39**, 815 (2004).
2. V.M. Ismailov, I.A. Mamedov and N.N. Yusubov, *Russ. J. Org. Chem.*, **40**, 284 (2004).
3. T. Matola and I. Jablonkai, *Crop Prot.*, **26**, 278 (2007).
4. S. Silva, S. Tardy, S. Routier, F. Suzenet, A. Tatibouët, A.P. Rauter and P. Rollin, *Tetrahedron Lett.*, **49**, 5583 (2008).
5. G.P. Cao, W.J. Chen and X.B. Liu, *Polym. Degrad. Stab.*, **93**, 739 (2008).
6. Y. Fu, L.H. Qu, S.S. Zhang, F. Ye, L.X. Zhao, S. Gao and Z.Y. Xing, *Heterocycl. Commun.*, **18**, 143 (2012).
7. Y. Fu, S.S. Zhang and F. Ye, *Indian J. Heterocycl. Chem.*, **20**, 405 (2011).
8. G.M. Sheldrick, SHELXS-97, Program for X-Ray Crystal Structure Solution, University of Göttingen, Göttingen, Germany (1997).
9. G.M. Sheldrick, SHELXS-97, Program for X-Ray Crystal Structure Refinement, University of Göttingen, Göttingen, Germany (1997).