# 6-Methyl-2,4-Disubstituted Pyridazin-3(2H)-ones: A Novel Class of Small-Molecule Agonists for Formyl Peptide Receptors

Agostino Cilibrizzi,<sup>†</sup> Mark T. Quinn,<sup>‡</sup> Liliya N. Kirpotina,<sup>‡</sup> Igor A. Schepetkin,<sup>‡</sup> Jeff Holderness,<sup>‡</sup> Richard D. Ye,<sup>§</sup> Marie-Josephe Rabiet,<sup>||</sup> Claudio Biancalani,<sup>†</sup> Nicoletta Cesari,<sup>†</sup> Alessia Graziano,<sup>†</sup> Claudia Vergelli,<sup>†</sup> Stefano Pieretti,<sup>⊥</sup> Vittorio Dal Piaz,<sup>†</sup> and Maria Paola Giovannoni<sup>\*,†</sup>

<sup>†</sup>Dipartimento di Scienze Farmaceutiche, Università degli Studi di Firenze, Via Ugo Schiff 6, Sesto Fiorentino 50019 Firenze, Italy, <sup>‡</sup>Department of Veterinary Molecular Biology, Montana State University, Bozeman, Montana 59717, <sup>§</sup>Department of Pharmacology, University of Illinois, Chicago, Illinois 60612, <sup>II</sup>CEA, DSV, iRTSV, Laboratoire Biochimie et Biophysique des Systèmes Intégrés, 17 rue des Martyrs, Grenoble F-38054, France, and <sup>L</sup>Department of Therapeutic Research and Medicine Evaluation, Viale Regina Elena 299, 00161 Roma, Italy

Received May 7, 2009

Following a ligand-based drug design approach, a potent mixed formyl peptide receptor 1 (FPR1) and formyl peptide receptor-like 1 (FPRL1) agonist (14a) and a potent and specific FPRL1 agonist (14x) were identified. These compounds belong to a large series of pyridazin-3(2*H*)-one derivatives substituted with a methyl group at position 6 and a methoxy benzyl at position 4. At position 2, an acetamide side chain is essential for activity. Likewise, the presence of lipophilic and/or electronegative substituents in the position para to the aryl group at the end of the chain plays a critical role for activity. Affinity for FPR1 receptors was evaluated by measuring intracellular calcium flux in HL-60 cells transfected with FPR1, FPRL1, and FPRL2. Agonists were able to activate intracellular calcium mobilization and chemotaxis in human neutrophils. The most potent chemotactic agent (EC<sub>50</sub> =  $0.6 \,\mu$ M) was the mixed FPR/FPRL1 agonist 14h.

## Introduction

Innate immunity is a very important mechanism in defending humans against infectious microbes.<sup>1</sup> Thus, immune disregulation due to viral infections (i.e., AIDS) and immunosuppression treatments following transplantation expose patients to life-threatening risks. Moreover, in the context of increasing aging of the population in more developed countries, it was observed that older people exhibit a natural immune function disregulation, which may be exacerbated in chronic stress conditions.<sup>2</sup> In addition, the ongoing emergence of bacterial and viral strains resistant to multiple classes of chemotherapeutics increases morbidity, mortality, and costs associated with nosocomial infections.<sup>3</sup> Thus, identification and development of bioactive molecules that selectively stimulate the innate immune response is an important challenge both for biologists and for chemists.<sup>4</sup>

The human formyl peptide receptor (FPR) family belongs to a group of G-protein coupled receptors that are predominantly expressed on neutrophils and monocytes.<sup>5</sup> Three FPR subtypes (FPR1<sup>*a*</sup>, FPRL1, and FPRL2) have been identified in humans, while eight FPR-related receptors have been characterized in mice.<sup>6</sup> Evidence for an important role of FPRs both in infective and inflammatory processes, has been well documented in the literature. The finding that bacteria are the major natural source of chemotactic formyl peptides (i.e., *N*-formylmethionine-leucine-phenylalanine, fMLF), which are high affinity agonists for FPR1, strongly supports the hypothesis that these receptors work as antibacterial receptors.<sup>7</sup> According to this idea, Fpr1-knockout mice show reduced resistance to infections provoked by some types of bacteria.<sup>8</sup> Moreover, a key role of FPRL1 has been proposed in host response to HIV-1 infection, and synthetic peptide domains of HIV-1 envelope proteins are able to activate FPRL1 leading to attenuation of cell response to chemokines through cross-desensitization and down-regulation of the monocyte receptors CCR5 and CXCR4.<sup>9</sup> The potent in vitro anti-HIV-1 activity of the synthetic gp41 peptide T2 (DP178), which is an agonist of FPR, strongly supports this view.<sup>10</sup>

Evidence for a pro-inflammatory role of FPRL1 in various neurodegenerative diseases was also found. In fact, the 42 amino acid form of amyloid  $\beta$  (AB42), which is believed to play a key role in Alzheimer disease-induced neuronal damage by recruitment and activation of mononuclear phagocytes, is a chemotactic agonist of FPRL1.<sup>11</sup> Likewise, the neurotoxic protein fragment PrP106-126, which behaves as the pathologic isoform of prion protein, was chemotactic and independently induced cytokine secretion in human monocytes through interaction with FPRL1.<sup>12</sup> On the other hand, antiinflammatory properties are displayed by different types of FPRL1 ligands. For example, lipoxin A4 (LXA4) is an endogenous arachidonic acid metabolite that has been shown to promote resolution of inflammation through FPRL1 agonism.<sup>13</sup> Likewise, annexin A1, a glucocorticoid-regulated protein that is particularly abundant in neutrophils, is able to inhibit leucocytes migration/activation through hFPRL1 agonism.<sup>14</sup> Taken together, these data clearly support a key role of FPRL1 in a variety of acute inflammatory and infective pathologies. Thus, this receptor has recently become

<sup>\*</sup>To whom correspondence should be addressed. Phone: +30-055-4573682. Fax: +39 55 4573671. E-mail mariapaola.giovannoni@ unifi.it.

<sup>&</sup>lt;sup>*a*</sup> Abbreviations: FPRs, formyl peptide receptors; FPR1, formyl peptide receptor 1; FPRL1, formyl peptide receptor-like 1; FPRL2, formyl peptide receptor-like 2; fMLF, *N*-formylmethionine-leucine-phenylalanine; LXA4, lipoxin A4; WT, wild-type; HBD, hydrogen bond donor; HBA, hydrogen bond acceptor.



Figure 1. Structures of FPRL1 agonists.

a prospective target for therapeutic intervention.<sup>15</sup> In addition, the identification of novel agonists and antagonists may be a useful tool to clarify the FPR-mediated (patho)physiological effects.

In addition to natural peptides like the bacterial fMLF,<sup>16</sup> synthetic peptides<sup>17</sup> and small molecule nonpeptide ligands of FPRL1 have been recently reported. Quin-C1<sup>18,19</sup> (compound 1, Figure 1) was the first small-molecule FPRL1-selective agonist shown to induce chemotaxis in peripheral blood neutrophils, with a relative potency 1000-fold lower than the synthetic peptide agonist WKYMVm. More recently, researchers from Amgen reported a series of pyrazolone derivatives displaying nanomolar selective affinity for FPRL1.<sup>20</sup> These types of agonists, of which **2** is an example, are characterized by the presence of an arylurea substructure linked to the heterocyclic backbone and served us as starting point to develop novel chemotypes of FPR ligands. Thus we report here the synthesis and evaluation of two series of functionalized pyridazinones of general structure **3** and **4**.

## Chemistry

The synthetic pathways employed to prepare the final compounds are depicted in Schemes 1-5.

Compound  $5^{21}$  in anhydrous toluene was refluxed with the appropriate aryl isocyanate to afford the urea derivatives 6a-c (Scheme 1). The 4-butoxy analogue 7 was synthesized with an alternative procedure starting from the same precursor and using triphosgene in anhydrous THF, followed by treatment with the appropriate aniline. The aromatic amide 8 was obtained from 5 by treatment with the opportune aroyl chloride in toluene at reflux. Finally, coupling of the amine 5 with 4-butoxyphenylboronic acid in CH<sub>2</sub>Cl<sub>2</sub> in presence of Cu(Ac)<sub>2</sub> gave the aromatic amine 9.

In Scheme 2 is depicted the synthesis of compounds 14a-x (Table 2). The dihydropyridazinone  $10^{22}$  was converted into the previously described 4-benzyl derivatives  $11b,c^{23,24}$  and

the new 11a by condensation with the appropriate aromatic aldehyde in the presence of KOH. Compounds 11, in turn, were alkylated with ethyl bromoacetate to give the esters 12ac, whose 12b,c were previously reported.<sup>24,25</sup> Alkaline hydrolysis of compounds 12 gave the known  $13c^{23}$  and the new carboxylic acids 13a,b. These compounds were treated with ethyl chloroformate in THF in presence of triethylamine, affording the intermediate mixed anhydrides, which, in turn, were transformed into the final amides 14a-s and 14u-x by treatment with the appropriate aryl or cycloalkyl amine. To obtain the ester 14t, the mixed anhydride was treated with 4-bromophenol.

Scheme 3 outlines the synthetic procedure for compounds 15–19: the precursor 11a was alkylated with the appropriate bromo ester to give compounds 15a,b, which, in turn, were converted into the corresponding acids 16a,b. The final step was the transformation of these compounds to the final amides 17a,b using the same procedure described for compounds 14. Compounds 18a,b were obtained by alkylation of 11a with the appropriate halide in standard conditions.

Moreover the precursor 11a was converted into the final 19a,b through a Mannich reaction  $(CH_2O + NH_3)$ . The intermediate amine was not isolated and was converted into the urea 19a by treatment with 4-bromophenyl isocyanate and into the amide 19b with 4-bromobenzoyl chloride.

The final compounds 21, 23, and 25 were synthesized as shown in Scheme 4. The intermediate 12a was reduced with sodium borohydride in THF/MeOH to generate the primary alcohol 20. This compound was the starting material for the synthesis both of the ether 21, through a coupling reaction with the 4-bromophenylboronic acid in presence of  $Cu(Ac)_2$ , and of compound 23, through the mesylate 22, which was converted into the final compound 23 by nucleophilic replacement with 4-bromo aniline. Treatment of 22 with ammonia gave the intermediate 24 from which the amide 25b and the urea 25a were obtained using 4-bromobenzoyl chloride or 4-bromophenyl isocyanate, respectively, using the same

#### Scheme 1<sup>*a*</sup>



<sup>*a*</sup> Reagents and conditions: (a) anhydrous toluene, aryl isocyanate, reflux, 3-7 h; (b) THF, anhydrous sodium acetate, triphosgene 70 °C, 2 h, 4-butoxyaniline, 12 h, rt; (c) Et<sub>3</sub>N, RCOCl, toluene, 110 °C, 4 h; (d) Cu(Ac)<sub>2</sub>, 4-butoxyphenylboronic acid, Et<sub>3</sub>N, rt, 12 h.

reaction conditions described for the analogues 7 and 8 in Scheme 1.

In Scheme 5 is depicted the synthesis of compounds **29** and **30**. Treatment of the commercially available 6-methylpyridazinone **26** with *m*-methoxybenzyl chloride in acetonitrile resulted in compound **27**, which, in turn, gave the corresponding 4-amino derivative **28** by heating with  $N_2H_4$  under hard conditions. The amide **29** and the urea **30** were obtained from **28** by treatment with the proper aroyl chloride or the aryl isocyanate by following the same conditions reported for **25a** and **25b**.

#### **Results and Discussion**

All synthesized compounds were evaluated for their ability to induce intracellular Ca<sup>2+</sup> flux in HL-60 cells transfected with FPR1, FPRL1, or FPRL2, and results were reported in Tables 1 and 2. All compounds were also evaluated in WT (wild-type nontransfected HL-60 cells) and were inactive. Moreover, both  $EC_{50}$  values and relative efficacy compared to the peptide agonists, fMLF and WKYMVm, were determined.

In the early phase of our project, we synthesized pyridazin-3(2*H*)-ones bearing a phenyl group at position 2 and a functionalized side chain at position 4 (compounds 6a-c, 7, 8, and 9), which may be regarded as enlarged analogues of 2 (Figure 1). Because these compounds failed to show activity (see Table 1), we turned our attention to analogues with a substituted benzyl at position 4 and a similar functionalized chain at N-2 (compounds 4, Figure 1). In this series, we were able to identify compound 14a, a potent mixed FPR1 and FPRL1 agonist.

Thus 14a was selected as lead and extensive structureactivity relationship (SAR) studies on this prototype were performed by modifying the nature and the length of the functionalized side chain and by replacing Br with a variety of substituents. Elongation of carbon chain from one to two methylene groups gave compound 17a, which resulted less potent at FPR1 and FPRL1 of 14a (Table 1). Branching the carbon chain resulted in compound 17b, which did not activate FPRL1 but retained activity for FPR1 similar to that of 14a. Thus, 17b is an FPR1-specific agonist. Modifications more substantial of the functionalized chain were obtained with compounds 18a,b and were completely detrimental. Further inactive compounds were the urea derivative 19a, the inverse amide 19b, and their superior homologues 25a and 25b. Replacement of CONH with an ether group and a secondary amine gaves compounds 21 and 23, respectively, which resulted also inactive. Moreover, compounds 29 and **30**, in which the substituents at position 2 and 4 were interchanged, were inactive.

Other modifications of the functionalized chain were performed with compounds 14r-u (Table 2). When the NH was converted to NCH<sub>3</sub> (compound 14r), the activity disappeared. Compound 14s, where the CONH is in the middle of the spacer, was also inactive. Replacement of CONH with COO (14t) or inclusion of amidic nitrogen in piperazine nucleus (14u) gave the same effect.

The nature and the position of the substituent on the phenyl group at the end of the chain also played a crucial role in ligand activity. Moving Br to positions meta (14b) and ortho (14c) resulted in a complete loss of FPR1/FPRL1 activity. Among halo-derivatives, the 4-chloro analogue 14d exhibited

## Scheme 2<sup>*a*</sup>



<sup>*a*</sup> Reagents and conditions: (a) aromatic aldehyde, 5% KOH, EtOH, reflux, 2-5 h; (b) ethyl bromoacetate, K<sub>2</sub>CO<sub>3</sub>, anhydrous CH<sub>3</sub>CN, reflux, 1-6 h; (c) NaOH, 60–80 °C, 1-4 h; (d) ethyl chloroformate, anhydrous THF, Et<sub>3</sub>N, appropriate substituted aryl(cycloalkyl)amine (or 4-bromophenol for compound **14t**), 12 h, rt.

the same profile as 14a. The corresponding 4-iodo derivative 14e was 2 times less potent at FPRL1. For this compound, a weak effect at FPRL2 was also observed. The 4-fluoro analogue 14f was less potent compared to 14a, but specificity for FPR1 was evident. The elimination of Br (compound 14g) was associated with loss of activity. Replacement of Br in 14a with substituents having similar steric properties, such as t-But (14i), OCF<sub>3</sub> (14o), and CN (14q), led to loss of activity at both FPR1 and FPRL1. The 4-trifluoromethyl and the 4-nitro analogues (14n and 14p, respectively) as well as compound 14h, bearing a methyl group at position 4, had relatively low activity. Introduction of alkoxy groups gave interesting results: the 4-methoxy derivative 14j and the 3,4dimethoxy derivative 14l had low activity at both receptors, whereas the 3,4-methylendioxy derivative 14m was specific for FPR1. It is worth noting that compounds 14v, 14w, and 14k, where Br in the para position was substituted with the typical buthoxy group characteristic of Quin-C1, were found to be completely devoid of activity.

Only a few substitutions were performed at the level of the aromatic system in the benzyl group. Moving OCH<sub>3</sub> from the meta to the para position (compound **14x**) resulted in high activity (EC<sub>50</sub> = 2.4  $\mu$ M) and selectivity for FPRL1. This exciting result prompted us to design several additional analogues, which are currently under investigation. Taken together, these data suggest that regarding the aromatic system at the end of the functionalized chain in position 2, the presence of a lipophilic and/or electronegative substituent, such as F, Br, or CH<sub>3</sub>, in position para is an essential requirement for potency and/or selectivity. Likewise, the

presence of an acetamide spacer at pyridazine N-2 also plays a crucial role in specificity and potency. The role of both CO and NH in the side chain seems to indicate that a hydrogen bond donor (HBD) neighboring an acceptor (HBA) system is also an essential requirement for binding at FPRs, according to the Quin-C1 and pyrazolone structures **2**, where the presence of an urea is a common structural element. Moreover, this HBD\HBA system must be placed at an appropriate distance from both the aromatic and the heterocyclic scaffold.

For compound 14a were also calculated dose–response curves (Figure 2). Note that no response was observed in control, undifferentiated HL-60 cells, which do not express FPRs. Furthermore, the effect of selected agonists on Ca<sup>2+</sup> flux in human neutrophils was also determined to verify the HL-60 results in primary phagocytes (Table 3). We found that both selective and nonselective agonists identified in HL-60 cell assays also induced Ca<sup>2+</sup> flux in human neutrophils, with EC<sub>50</sub> values in the range  $0.8-21.7 \,\mu$ M.

The most potent chemotactic compound was **14h** (EC<sub>50</sub> =  $0.6 \,\mu$ M), followed by **14j** (EC<sub>50</sub> =  $0.9 \,\mu$ M), which were both nonselective agonists. The FPRL1-selective agonist (**14x**) showed lower potency (EC<sub>50</sub> =  $13.1 \,\mu$ M) as a chemotactic agent.

In conclusion, we have identified a novel chemotype of interesting and selective FPR1 or FPRL1 agonists and mixed FPR/FPRL1 agonists active in the low micromolar range in human neutrophils. Moreover, some of these compounds proved to be able to stimulate chemotaxis at submicromolar concentrations.

### Scheme 3<sup>*a*</sup>



<sup>*a*</sup> Reagents and conditions: (a) appropriate alkyl halide,  $K_2CO_3$ , anhydrous CH<sub>3</sub>CN, reflux, 3 h; (b) NaOH, 60 °C, rt; (c) ethyl chloroformate, anhydrous THF, Et<sub>3</sub>N, 4-bromoaniline, 12 h, rt; (d) appropriate alkyl halide,  $K_2CO_3$ , anhydrous CH<sub>3</sub>CN, reflux, 2–4 h; (e) 40% CH<sub>2</sub>O, 33% NH<sub>3</sub>, dioxane, CH<sub>2</sub>Cl<sub>2</sub>, 4-bromophenyl isocyanate (for compound **19a**), or 4-bromobenzoyl chloride (for compound **19b**), 0 °C to rt, 6–12 h.

## **Experimental Section**

Chemistry. All melting points were determined on a Buchi apparatus and are uncorrected. IR spectra were measured as Nujol mulls with a PerkinElmer spectrometer (FT-IR, Spectrum 1000). <sup>1</sup>H NMR spectra were recorded with Avance 400 instruments (Bruker Biospin Version 002 with SGU). Chemical shifts are reported in ppm, using the solvent as internal standard. Mass spectra (m/z) were recorded on a ESI-TOF mass spectrometer (Bruker Micro TOF). Extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, and the solvents were removed under reduced pressure. Merck F-254 commercial plates were used for analytical TLC to follow the course of reaction. Silica gel 60 (Merck 70-230 mesh) was used for column chromatography. Microanalyses were performed with a Perkin-Elmer 260 elemental analyzer for C, H, N, and the results were within  $\pm 0.4\%$  of the theoretical values unless otherwise stated. Reagents and starting materials were commercially available.

**General Procedure for 6a–c.** To a stirred solution of compound  $5^{21}(0.35 \text{ mmol})$  in anhydrous toluene (2 mL), the proper aryl isocyanate (0.40 mmol) was added. The mixture was refluxed for 4–7 h. After cooling, the precipitate was collected by suction and purified by crystallization from toluene (6b and

**6c**). In the case of **6a**, a second batch of product was obtained from evaporation of the filtrate. The residue, after treatment with cold water, was extracted with  $CH_2Cl_2$  (3 × 20 mL). Removal of the solvent gave a residue that was purified by column chromatography using  $CH_2Cl_2/CH_3OH$  9.9:0.1 as eluent.

**1-(4-Iodophenyl)-3-(6-methyl-3-oxo-2-phenyl-2,3-dihydropyridazin-4-yl)urea** (6a). Yield = 45%; mp = 264–265 °C (EtOH/toluene). IR (cm<sup>-1</sup>) 3270 (NH), 3256 (NH), 1708 (CO), 1627 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.44 (s, 3H, CH<sub>3</sub>), 6.50 (d, 2H, Ar, J = 8.7 Hz), 7.21 (t, 1H, Ar, J = 7.5 Hz), 7.36–7.41 (m, 4H, Ar), 7.68 (d, 2H, Ar, J = 7.7 Hz), 8.15 (s, 1H, Ar), 8.41 (exch br s, 1H, NH), 9.38 (exch br s, 1H, NH). MS (ESI) m/z 447.03 [M + H]<sup>+</sup>. Anal. (C<sub>18</sub>H<sub>15</sub>IN<sub>4</sub>O<sub>2</sub>) C, H, N.

**1-(4-Bromophenyl)-3-(6-methyl-3-oxo-2-phenyl-2,3-dihydropyridazin-4-yl)urea (6b).** Yield = 50%; mp = 263–265 °C (toluene). IR (cm<sup>-1</sup>) 3271 (NH), 3250 (NH), 1704 (CO), 1625 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.43 (s, 3H, CH<sub>3</sub>), 6.60 (d, 2H, Ar, *J* = 8.6 Hz), 7.19–7.22 (m, 3H, Ar), 7.38 (t, 2H, Ar, *J* = 7.8 Hz), 7.69 (d, 2H, Ar, *J* = 7.9 Hz), 8.16 (s, 1H, Ar), 8.44 (exch, br, s, 1H, NH), 9.40 (exch br s, 1H, NH). MS (ESI) *m*/*z* 399.05 [M + H]<sup>+</sup>. Anal. (C<sub>18</sub>H<sub>15</sub>BrN<sub>4</sub>O<sub>2</sub>) C, H, N.

#### Scheme 4<sup>*a*</sup>



<sup>*a*</sup> Reagents and conditions: (a) anhydrous THF, NaBH<sub>4</sub>, CH<sub>3</sub>OH, 60 °C, 1 h; (b) Cu(Ac)<sub>2</sub>, 4-bromophenylboronic acid, anhydrous CH<sub>2</sub>Cl<sub>2</sub>, Et<sub>3</sub>N, rt, 20 h; (c) pyridine, CH<sub>2</sub>Cl<sub>2</sub>, methanesulfonyl chloride, 0 °C to rt, 4 h; (d) 2-propanol, 4-bromoaniline, 60 °C, 6 h; (e) 33% NH<sub>3</sub>, isopropanol, 60 °C, 3 h; (f) anhydrous toluene, 4-bromophenyl isocyanate, 110 °C, 5 h (for compound **25a**); anhydrous CH<sub>2</sub>Cl<sub>2</sub>, Et<sub>3</sub>N, 4-bromobenzoyl chloride, 0 °C, 6 h (for compound **25b**).

**1-(4-Chlorophenyl)-3-(6-methyl-3-oxo-2-phenyl-2,3-dihydropyridazin-4-yl)urea (6c).** Yield = 85%; mp = 274–276 °C (toluene). IR (cm<sup>-1</sup>) 3269 (NH), 3255 (NH), 1705 (CO), 1626 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.44 (s, 3H, CH<sub>3</sub>), 6.65 (d, 2H, Ar, J = 8.8 Hz), 7.07 (d, 2H, Ar, J = 8.8 Hz), 7.21 (t, 1H, Ar, J = 7.5 Hz), 7.38 (t, 2H, Ar, J = 7.9 Hz), 7.69 (d, 2H, Ar, J = 7.6 Hz), 8.16 (s, 1H, Ar), 8.43 (exch br s, 1H, NH), 9.40 (exch br s, 1H, NH). MS (ESI) m/z 355.10 [M + H]<sup>+</sup>. Anal. (C<sub>18</sub>H<sub>15</sub>ClN<sub>4</sub>O<sub>2</sub>) C, H, N.

1-(4-Butoxyphenyl)-3-(6-methyl-3-oxo-2-phenyl-2,3-dihydropyridazin-4-yl)urea (7). To a cooled and stirred suspension of 5 (0.5 mmol) and anhydrous sodium acetate (1.19 mmol) in anhydrous THF (5 mL), triphosgene (1.18 mmol) was added. The mixture was stirred for an additional 10 min at 0 °C and refluxed for 2 h. The solvent was removed in vacuo, and the residue was dissolved in anhydrous THF (2 mL). 4-Butoxyaniline (1.25 mmol) was added, and the mixture was stirred at room temperature for 12 h. After dilution with cold water, the suspension was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 20 mL). After removal of the solvent, the residue was purified by column chromatography using absolute EtOH/CH<sub>2</sub>Cl<sub>2</sub>/petroleum ether/toluene/33% ammonia 3.3:19.7:6.5:70:0.5 as eluent. Yield = 31%; mp = 189–191 °C (EtOH). IR (cm<sup>-1</sup>) 3270 (NH), 3255 (NH), 1705 (CO), 1625 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.01 (t, 3H, CH<sub>2</sub>CH<sub>3</sub>, J = 7.4 Hz), 1.52 (sext, 2H, CH<sub>2</sub>CH<sub>3</sub>, J = 7.4 Hz), 1.79 (quint, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, J = 7.0 Hz), 2.40 (s, 3H, 6-CH<sub>3</sub>), 3.95 (t, 2H, OCH<sub>2</sub>, J = 6.6 Hz), 6.70 (s, 4H, Ar), 7.17 (t, 1H, Ar, J = 7.5 Hz), 7.32 (t, 2H, Ar, J = 7.8 Hz), 7.61 (d, 2H, Ar, J = 7.9 Hz), 8.11 (exch br s, 1H, NH), 8.14 (s, 1H, Ar), 9.33 (exch br s, 1H, NH). MS (ESI) m/z 393.19 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>24</sub>N<sub>4</sub>O<sub>3</sub>) C, H, N.

**4-Butoxy-***N***-(6-methyl-3-oxo-2-phenyl-2,3-dihydropyridazin-4-yl)benzamide (8).** 4-Butoxybenzoic acid (0.36 mmol) was converted to the corresponding chloride as follows: SOCl<sub>2</sub> (20.7 mmol) was added and the stirred suspension was cooled to 0 °C and treated with Et<sub>3</sub>N (0.99 mmol). After 4 h of reflux, the mixture was evaporated in vacuo and the residue was washed Scheme 5<sup>a</sup>



<sup>*a*</sup> Reagents and conditions: (a) 3-methoxybenzyl chloride, anhydrous CH<sub>3</sub>CN, anhydrous K<sub>2</sub>CO<sub>3</sub>, 80 °C, 2 h; (b) hydrazine hydrate, 180 °C, 11 h; (c) 4-bromobenzoyl chloride, Et<sub>3</sub>N, 0 °C, 4 h; (d) 4-bromophenyl isocyanate, anhydrous toluene, reflux, 7 h.

with cyclohexane  $(3 \times 5 \text{ mL})$ . The residue was dissolved in anhydrous toluene (5 mL), and the solution was treated with a solution of compound 5 (0.40 mmol) in toluene (1 mL) and Et<sub>3</sub>N (1.48 mmol). After 4 h of reflux, the solvent was removed in vacuo and the residue was treated with cold water and extracted with  $CH_2Cl_2$  (3 × 20 mL). After removal of the solvent, the residue was purified by column chromatography (eluent: cyclohexane/ethyl acetate 3:1). Yield =37%; mp =95-97 °C (EtOH). IR (cm<sup>-1</sup>) 3269 (NH), 1682 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.01 (t, 3H, CH<sub>2</sub>CH<sub>3</sub>, J = 7.4 Hz), 1.53 (sext, 2H, CH<sub>2</sub>CH<sub>3</sub>, J = 7.5 Hz), 1.82 (quint, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, J = 7.0 Hz), 2.45 (s, 3H, 6-CH<sub>3</sub>), 4.06 (t, 2H, CH<sub>2</sub>O, J = 6.5 Hz), 6.99 (d, 2H, Ar, J = 8.8 Hz), 7.43 (t, 1H, Ar, J = 7.4 Hz), 7.52 (t, 2H, Ar, J = 7.8 Hz), 7.65 (d, 2H, Ar, J = 7.7 Hz), 7.91 (d, 2H, Ar, J = 8.8 Hz), 8.24 (s, 1H, Ar), 9.46 (exch br s, 1H, NH). MS (ESI) m/z 378.18 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>) C, H, N.

4-(4-Butoxyphenylamino)-6-methyl-2-phenylpyridazin-3(2H)one (9). To a suspension of 5 (0.55 mmol), copper acetate (0.82 mmol) and 4-butoxyphenylboronic acid (1.08 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL), Et<sub>3</sub>N (1.08 mmol) was added and the mixture was stirred at room temperature for 12 h. The suspension was extracted with 15% aqueous ammonia (10 mL), and the organic layer was washed with 10 mL of water and dried over sodium sulfate. After removal of the solvent, the residue was purified by column chromatography using as eluent cyclohexane/ethyl acetate 3:1. The analytical sample was obtained from a further purification performed on a silica gel preparative column (eluent: cyclohexane/ethyl acetate 3:1). Yield = 16%; mp = 137-139 °C (EtOH). IR (cm<sup>-1</sup>) 3275 (NH), 1633 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.02 (t, 3H,  $CH_2CH_3$ , J = 7.4 Hz), 1.54 (sext, 2H,  $CH_2CH_3$ , J = 7.0 Hz), 1.81 (quint, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.31 (s, 3H, 6-CH<sub>3</sub>), 4.01 (t, 2H,  $CH_2O, J = 6.5 Hz$ ), 6.40 (s, 1H, Ar), 6.97 (d, 2H, Ar, J = 8.9 Hz), 7.20 (d, 2H, Ar, J = 8.8 Hz), 7.41 (t, 1H, Ar, J = 7.4 Hz), 7.51

(t, 3H, Ar, J = 7.7 Hz), 7.64 (d, 2H, Ar, J = 8.0 Hz). MS (ESI) m/z 350.19 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub>) C, H, N.

**4-(3-Methoxybenzyl)-6-methylpyridazin-3(2***H***)-one (11a). To a solution of <b>10** (1.79 mmol) in 7 mL of a solution of KOH 5% (w/v) in absolute EtOH, 3-methoxybenzaldehyde (1.79 mmol) was added and the mixture was refluxed under stirring for 3 h. After cooling, the sample was concentrated in vacuo, diluted with cold water (10–15 mL), and acidified with 2 N HCl. The suspension was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL). Removal of the solvent afforded compound **11a**, which was purified by flash chromatography using cyclohexane/ethyl acetate 1:1 as eluent. Yield = 61%; mp = 133–134 °C (EtOH). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$ 2.26 (s, 3H, 6-CH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 6.73 (s, 1H, Ar), 6.82–6.86 (m, 3H, Ar), 7.276–7.28 (m, 1H, Ar).

[5-(3-Methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]acetic acid ethyl ester (12a). A mixture of 11a (1.13 mmol),  $K_2CO_3$  (2.26 mmol), and ethyl bromoacetate (3.40 mmol) in CH<sub>3</sub>CN (3 mL) was refluxed under stirring for 6 h. The mixture was then concentrated in vacuo, diluted with cold water, and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL). The solvent was evaporated in vacuo, and compound 12a was purified by column chromatography using cyclohexane/ethyl acetate 1:1 as eluent. Yield = 98%; oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.29t, 3H, CH<sub>2</sub>CH<sub>3</sub>, J = 7.1 Hz), 2.22 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 2H, CH<sub>2</sub>-Ar), 4.22-4.25 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 4.85 (s, 2H, NCH<sub>2</sub>CO), 6.67 (s, 1H, Ar), 6.78-6.82 (m, 3H, Ar), 7.26 (t, 1H, Ar, J = 7.8 Hz).

General Procedure for 13a,b. A suspension of the appropriate derivatives 12a or 12b,<sup>24</sup> respectively (0.5-1.11 mmol), and 6 N NaOH (6–8 mL) in ethanol (2 mL) was stirred at rt to 80 °C for 1–4 h. The mixtures were then concentrated in vacuo, diluted with cold water, and acidified with 6 N HCl, and the final products were filtered off with suction and recrystallized from ethanol.

Table 1. Activity of the Compounds in Human HL-60 Cells Expressing Human FPR1, FPRL1, or FPRL2





		Y	R <sub>1</sub>	$R_2$	$\operatorname{Ca}^{2+}$ mobilization $\operatorname{EC}_{50}(\mu M)$ and efficacy $(\%)^a$		
compd	Х				FPR1	FPRL1	FPRL2
6a	NHCONH		Ι		NA	NA	NA
6b	NHCONH		Br		NA	NA	NA
6c	NHCONH		Cl		NA	NA	NA
7	NHCONH		$OC_4H_9$		NA	NA	NA
8	NHCO		$OC_4H_9$		NA	NA	NA
9	NH		$OC_4H_9$		NA	NA	NA
14a	$CH_2$	CH <sub>2</sub> CONH	3-OCH <sub>3</sub>	4-Br	$3.4 \pm 1.6$ (75)	$3.8 \pm 1.5$ (70)	NA
17a	$CH_2$	(CH <sub>2</sub> ) <sub>2</sub> CONH	3-OCH <sub>3</sub>	4-Br	$9.7 \pm 2.7 (30)$	$5.4 \pm 1.2$ (25)	NA
17b	$CH_2$	CH(CH <sub>3</sub> )CONH	3-OCH <sub>3</sub>	4-Br	$3.2 \pm 1.5$ (90)	NA	NA
18a	CH <sub>2</sub>	CH <sub>2</sub>	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
18b	CH <sub>2</sub>	CH <sub>2</sub> CO	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
19a	$CH_2$	CH <sub>2</sub> NHCONH	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
19b	$CH_2$	CH <sub>2</sub> NHCO	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
21	$CH_2$	$(CH_2)_2O$	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
23	$CH_2$	$(CH_2)_2NH$	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
25a	$CH_2$	(CH <sub>2</sub> ) <sub>2</sub> NHCONH	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
25b	$CH_2$	(CH <sub>2</sub> ) <sub>2</sub> NHCO	3-OCH <sub>3</sub>	4-Br	NA	NA	NA
29	NHCO	$CH_2$	4-Br	3-OCH <sub>3</sub>	NA	NA	NA
30	NHCONH	$CH_2$	4-Br	3-OCH <sub>3</sub>	NA	NA	NA
fMLF					0.01	20.4	1.9
WKYMVm					0.5	0.001	0.01

<sup>*a*</sup>NA, no activity was observed (no response was observed during first 2 min after addition of compounds under investigation). The EC<sub>50</sub> values are presented as the mean  $\pm$  SD of three independent experiments, in which median effective concentration values (EC<sub>50</sub>) were determined by nonlinear regression analysis of the dose–response curves (5–6 points) generated using GraphPad Prism 5 with 95% confidential interval (p < 0.05). Efficacy (in bracket) is expressed as percent of the response induced by 5 nM fMLF (FPR1) or 5 nM WKYMVm (FPRL1 and FPRL2).

[5-(3-Methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]acetic Acid (13a). Yield = 97%; mp = 194–195 °C (EtOH). <sup>1</sup>H NMR (DMSO)  $\delta$  2.21 (s, 3H, 3-CH<sub>3</sub>), 3.72 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 2H, CH<sub>2</sub>-Ar), 4.69 (s, 2H, NCH<sub>2</sub>CO), 6.79–6.85 (m, 3H, Ar), 7.07 (s, 1H, Ar), 7.22 (t, 1H, Ar, J = 7.8 Hz).

[5-(4-Methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]acetic Acid (13b). Yield = 96%; mp = 144–145 °C (EtOH). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.26 (s, 3H, 3-CH<sub>3</sub>), 3.84 (s, 3H, OCH<sub>3</sub>), 3.87 (s, 2H, CH<sub>2</sub>-Ar), 4.93 (s, 2H, NCH<sub>2</sub>CO), 6.69 (s, 1H, Ar), 6.91 (d, 2H, Ar, *J* = 8.5 Hz), 7.16 (d, 2H, Ar, *J* = 8.5 Hz).

General Procedure for 14a-x. To a cooled (-5 °C) and stirred solution of compound 13a (0.35 mmol) in anhydrous tetrahydrofuran (3–5 mL), Et<sub>3</sub>N (1.22 mmol) was added. After 30 min, the mixture was allowed to warm up to 0 °C, and ethyl chloroformate (0.38 mmol) was added. After 1 h, the appropriate substituted aryl(cycloalkyl)amine (for compounds 14a-s and 14u-x) or 4-bromophenol (for compound 14t), commercially available (0.7 mmol), was added. The reactions were carried out at room temperature for 12 h. The mixtures were then concentrated in vacuo, diluted with cold water (20-30 mL), and extracted with  $CH_2Cl_2$  (3 × 15 mL). The solvent was evaporated to afford final compounds 14a-x, which were purified by column chromatography using cyclohexane/ethyl acetate 1:1 as eluent for compounds 14a, 14d, 14f-m, 14q, 14w; toluene/NH<sub>4</sub>OH/EtOH/CH<sub>2</sub>Cl<sub>2</sub>/ petroleum ether 7:0.05:0.30:2:0.65 for compounds 14b,c; 14x; cyclohexane/ethyl acetate 1:2 for compounds 14e, 14n-p, 14s,

14v; cyclohexane/ethyl acetate 2:1 for compounds 14r,t and CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 9.5:0.5 for compound 14u.

*N*-(4-Bromophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14a). Yield = 86%; mp = 171− 172 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1708 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.20 (s, 3H, 3-CH<sub>3</sub>), 3.70 (s, 3H, OCH<sub>3</sub>), 3.73 (s, 2H, CH<sub>2</sub>-Ar), 4.80 (s, 2H, NCH<sub>2</sub>CO), 6.77−6.82 (m, 3H, Ar), 7.13 (s, 1H, Ar), 7.20 (t, 1H, Ar, *J* = 8.0 Hz), 7.47 (s, 4H, Ar). MS (ESI) *m*/*z* 442.08 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N. *N*-(3-Bromophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-

*N*-(3-Bromophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14b). Yield = 85%; mp = 73-75 °C (EtOH). IR (cm<sup>-1</sup>) 3295 (NH), 1708 (CO), 1642 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.30 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.92 (s, 2H, CH<sub>2</sub>-Ar), 4.95 (s, 2H, NCH<sub>2</sub>CO), 6.80-6.86 (m, 4H, Ar), 7.14 (t, 1H, Ar, *J* = 8.0 Hz), 7.22 (m, 1H, Ar, *J* = 8.2 Hz), 7.27-7.31 (m, 1H, Ar), 7.39 (d, 1H, Ar, *J* = 8.0 Hz), 7.77 (s, 1H, Ar), 9.05 (exch br s, 1H, NH). MS (ESI) *m*/*z* 271.11 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

*N*-(2-Bromophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14c). Yield = 80%; oil. IR (cm<sup>-1</sup>) 3300 (NH), 1708 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.30 (s, 3H, 3-CH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.92 (s, 2H, CH<sub>2</sub>-Ar), 5.00 (s, 2H, NCH<sub>2</sub>CO), 6.76 (s, 1H, Ar), 6.80–6.86 (m, 3H, Ar), 7.00 (t, 1H, Ar, *J* = 7.7 Hz), 7.27–7.34 (m, 2H, Ar), 7.52 (d, 1H, Ar, *J* = 8.0 Hz), 8.36 (d, 1H, Ar, *J* = 8.2 Hz), 8.54 (exch br s, 1H, NH). MS (ESI) *m*/z 442.08 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.





		Y	R <sub>1</sub>	$Ca^{2+}$ mobilization $EC_{50}$ ( $\mu$ M) and efficacy (%) <sup><i>a</i></sup>		
compd	Х			FPR1	FPRL1	FPRL2
14a	OCH <sub>3</sub>	Н	$NH-C_6H_4$ -Br (p)	3.4 ± 1.6 (75)	3.8 ± 1.5 (70)	NA
14b	$OCH_3$	Η	$NH-C_6H_4$ -Br (m)	NA	NA	NA
14c	OCH <sub>3</sub>	Н	$NH-C_6H_4$ -Br (o)	NA	NA	NA
14d	OCH <sub>3</sub>	Н	$NH-C_6H_4-Cl(p)$	$2.6 \pm 0.3$ (110)	$4.0 \pm 1.6$ (35)	NA
14e	OCH <sub>3</sub>	Н	$NH-C_6H_4-I(p)$	$2.8 \pm 0.2$ (90)	$6.8 \pm 2.2$ (40)	$13.0 \pm 3.1 (30)$
14f	OCH <sub>3</sub>	Н	$NH-C_6H_4-F(p)$	$7.6 \pm 0.2$ (40)	NA	NA
14g	OCH <sub>3</sub>	Н	NH-C <sub>6</sub> H <sub>5</sub>	NA	NA	NA
14h	OCH <sub>3</sub>	Н	$NH-C_6H_4-CH_3(p)$	$7.2 \pm 2.2 (120)$	$10.9 \pm 3.4 (50)$	NA
14i	OCH <sub>3</sub>	Н	$NH-C_{6}H_{4}-tC_{4}H_{9}(p)$	NA	NA	NA
14j	OCH <sub>3</sub>	Н	$NH-C_6H_4-OCH_3(p)$	$7.7 \pm 2.5$ (65)	$14.4 \pm 2.0$ (35)	NA
14k	OCH <sub>3</sub>	Н	$NH-C_6H_4-OC_4H_9(p)$	NA	NA	NA
14 L	OCH <sub>3</sub>	Н	$NH-C_{6}H_{4}-(OCH_{3})_{2}$ (m, p)	$15.5 \pm 2.9$ (25)	$16.8 \pm 3.2$ (25)	NA
14m	OCH <sub>3</sub>	Н	$\mathbf{A}^b$	$2.3 \pm 1.1$ (50)	NA	NA
14n	OCH <sub>3</sub>	Н	$NH-C_6H_4-CF_3(p)$	$5.7 \pm 1.8 (50)$	$8.8 \pm 2.3$ (95)	NA
140	OCH <sub>3</sub>	Н	$NH-C_6H_4-OCF_3(p)$	NA	NA	NA
14p	OCH <sub>3</sub>	Н	$NH-C_6H_4-NO_2(p)$	$10.5 \pm 2.9$ (60)	$12.3 \pm 2.5$ (55)	NA
14q	OCH <sub>3</sub>	Н	$NH-C_6H_4-CN(p)$	NA	NA	NA
14r	OCH <sub>3</sub>	Н	$N(CH_3)-C_6H_4-Br(p)$	NA	NA	NA
14s	OCH <sub>3</sub>	Н	$NHCH_2$ - $C_6H_4$ -Br (p)	NA	NA	NA
14t	OCH <sub>3</sub>	Н	$O-C_6H_4$ -Br (p)	NA	NA	NA
14u	OCH <sub>3</sub>	Н	$\mathbf{B}^{b}$	NA	NA	NA
14v	Н	OCH <sub>3</sub>	$NH-C_6H_4-OC_4H_9(p)$	NA	NA	NA
14w	Н	Cl	$NH-C_6H_4-OC_4H_9(p)$	NA	NA	NA
14x	Н	OCH <sub>3</sub>	$NH-C_6H_4$ -Br (p)	NA	$2.4 \pm 0.9$ (70)	NA
fMLF			· · ·	0.01	20.4	1.9
WKYMVm				0.5	0.001	0.01

<sup>*a*</sup>NA, no activity was observed (no response was observed during first 2 min after addition of compounds under investigation). The EC<sub>50</sub> values are presented as the mean  $\pm$  SD of three independent experiments, in which median effective concentration values (EC<sub>50</sub>) were determined by nonlinear regression analysis of the dose–response curves (5–6 points) generated using GraphPad Prism 5 with 95% confidential interval (p < 0.05). Efficacy (in bracket) is expressed as percent of the response induced by 5 nM fMLF (FPR1) or 5 nM WKYMVm (FPRL1 and FPRL2). <sup>*b*</sup>



*N*-(4-Chlorophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14d). Yield = 86%; mp = 170– 171 °C (EtOH). IR (cm<sup>-1</sup>) 3296 (NH), 1705 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.30 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 4.95 (s, 2H, NCH<sub>2</sub>CO), 6.79 (s, 1H, Ar), 6.80– 6.85 (m, 3H, Ar), 7.16 (d, 2H, Ar, *J* = 8.8 Hz), 7.25–7.29 (m, 1H, Ar), 7.38 (d, 2H, Ar, *J* = 8.8 Hz), 9.28 (exch br s, 1H, NH). MS (ESI) *m*/*z* 398.13 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>ClN<sub>3</sub>O<sub>3</sub>) C, H, N.

*N*-(4-Iodophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14e). Yield = 99%; mp = 179– 180 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1707 (CO), 1645 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.30 (s, 3H, 3-CH<sub>3</sub>), 3.79 (s, 3H, OCH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 4.95 (s, 2H, NCH<sub>2</sub>CO), 6.78–6.84 (m, 4H, Ar), 7.17 (d, 2H, Ar, *J* = 8.7 Hz), 7.24–7.28 (m, 1H, Ar), 7.45 (d, 2H, Ar, *J* = 8.7 Hz), 9.41 (exch br s, 1H, NH). MS (ESI) *m/z* 490.06 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>IN<sub>3</sub>O<sub>3</sub>) C, H, N.

*N*-(4-Fluorophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14f). Yield = 97%; mp = 145-147 °C

(EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1707 (CO), 1643 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.30 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 4.95 (s, 2H, NCH<sub>2</sub>CO), 6.80–6.84 (m, 4H, Ar), 6.90 (t, 2H, Ar, J = 8.7 Hz), 7.25–7.29 (m, 1H, Ar), 7.39–7.42 (m, 2H, Ar), 9.21 (exch br s, 1H, NH). MS (ESI) *m*/*z* 382.16 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>FN<sub>3</sub>O<sub>3</sub>) C, H, N.

**2-[5-(3-Methoxybenzyl)-3-methyl-6-oxo-***6H***-pyridazin-1-yl]***-N***-phenyl-acetamide (14g).** Yield = 97%; oil. IR (cm<sup>-1</sup>) 3295 (NH), 1708 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.28 (s, 3H, 3-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.91 (s, 2H, CH<sub>2</sub>-Ar), 4.96 (s, 2H, NCH<sub>2</sub>CO), 6.78–6.86 (m, 4H, Ar), 7.08 (t, 1H, Ar, *J* = 7.4 Hz), 7.26–7.30 (m, 3H, Ar), 7.50 (d, 2H, Ar, *J* = 8.0 Hz), 8.93 (exch br s, 1H, NH). MS (ESI) *m*/*z* 364.16 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub>) C, H, N.

**2-[5-(3-Methoxybenzyl)-3-methyl-6-oxo-6H-pyridazin-1-yl]**-*N*-**4-tolyl-acetamide (14h).** Yield = 78%; mp = 142–144 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1708 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.28 (s, 3H, 3-CH<sub>3</sub>), 2.30 (s, 3H, CH<sub>3</sub>-Ar), 3.81 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 4.94 (s, 2H, NCH<sub>2</sub>CO), 6.76



**Figure 2.** Analysis of  $Ca^{2+}$  mobilization in phagocytes treated with compound **14a**. HL-60-FPR1 and HL-60-FPRL1 cells (A) or human neutrophils (B) were loaded with FLIPR calcium 3 dye, and  $Ca^{2+}$  flux was analyzed, as described. Responses were normalized to the response induced by 5 nM fMLF for HL-60-FPR1 cells and neutrophils, or 5 nM WKYMVm for HL-60-FPRL1 cells, which were assigned a value of 100%. (C) Representative kinetics of  $Ca^{2+}$  mobilization after treatment with compound **14a** or fMLF. Human neutrophils were treated with the compound **14a** (1.5 and 3  $\mu$ M), 5 nM fMLF (positive control), or 1% DMSO (negative control), and  $Ca^{2+}$  flux was monitored for the indicated times. The data are from one experiment that is representative of three independent experiments.

**Table 3.**  $Ca^{2+}$  Mobilization and Chemotactic Activity in HumanNneutrophils Treated with Selected FPR1/FPRL1 Agonists<sup>a</sup>

	EC <sub>50</sub> (µM)				
compd	Ca <sup>2+</sup> mobilization	chemotaxis			
14a	$2.6 \pm 0.3$	$2.1 \pm 0.8$			
14d	$6.7 \pm 1.1$	$1.6 \pm 0.2$			
14e	$3.2 \pm 1.2$	$1.8 \pm 0.3$			
14f	$3.9 \pm 0.6$	$8.2 \pm 1.4$			
14h	$3.2 \pm 0.3$	$0.6 \pm 0.3$			
14j	$1.6 \pm 0.8$	$0.9 \pm 0.2$			
141	$1.1 \pm 0.6$	$1.1 \pm 0.6$			
14m	$3.6 \pm 1.0$	$1.2 \pm 0.6$			
14n	$3.6 \pm 0.8$	$4.5 \pm 2.5$			
14p	$21.7 \pm 4.2$	$1.9 \pm 0.6$			
14x	$4.3 \pm 1.1$	$13.1 \pm 2.3$			
17a	$11.3 \pm 2.8$	$11.8 \pm 2.6$			
17b	$0.8 \pm 0.2$	$0.6\pm0.4$			

<sup>*a*</sup> The data are presented as the mean  $\pm$  SD of three independent experiments with cells from different donors, in which median effective concentration values (EC<sub>50</sub>) were determined by nonlinear regression analysis of the dose–response curves (5–6 points) generated using GraphPad Prism 5 with 95% confidential interval (p < 0.05).

(s, 1H, Ar), 6.80–6.86 (m, 3H, Ar), 7.08 (d, 2H, Ar, J = 8.3 Hz), 7.26–7.30 (m, 1H, Ar), 7.38 (d, 2H, Ar, J = 8.4 Hz), 8.82 (exch br s, 1H, NH). MS (ESI) m/z 378.18 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>) C, H, N.

*N*-(4-*tert*-Butylphenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14i). Yield =95%; mp =54-56 °C (EtOH). IR (cm<sup>-1</sup>) 3296 (NH), 1705 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.30 (s, 9H, C-(CH<sub>3</sub>)<sub>3</sub>), 2.27 (s, 3H, 3-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 4.96 (s, 2H, NCH<sub>2</sub>CO), 6.76 (s, 1H, Ar), 6.80 (s, 1H, Ar), 6.81–6.86 (m, 2H, Ar), 7.26–7.30 (m, 3H, Ar), 7.43 (d, 2H, Ar, J = 8.6 Hz), 8.88 (exch br s, 1H, NH). MS (ESI) m/z 420.22 [M + H]<sup>+</sup>. Anal. (C<sub>25</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub>) C, H, N.

**2-**[5-(3-Methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-*N*-(4-methoxyphenyl)-acetamide (14j). Yield = 75%; mp = 65– 67 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1705 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.28 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.91 (s, 2H, CH<sub>2</sub>-Ar), 4.94 (s, 2H, NCH<sub>2</sub>CO), 6.77 (s, 1H, Ar), 6.80 (s, 1H, Ar), 6.81–6.86 (m, 4H, Ar), 7.27–7.30 (m, 1H, Ar), 7.42 (d, 2H, Ar, *J* = 8.9 Hz), 8.71 (exch br s, 1H, NH). MS (ESI) *m*/*z* 394.18 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>23</sub>N<sub>3</sub>O<sub>4</sub>) C, H, N.

*N*-(4-Butoxyphenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14k). Yield = 73%; oil. IR (cm<sup>-1</sup>) 3300 (NH), 1708 (CO), 1645 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.99 (t, 3H, O(CH<sub>2</sub>)<sub>3</sub>*CH*<sub>3</sub>), 1.50 (sext, 2H, OCH<sub>2</sub>CH<sub>2</sub>*CH*<sub>2</sub>, *J* = 7.5 Hz), 1.76 (quint, 2H, OCH<sub>2</sub>*CH*<sub>2</sub>CH<sub>2</sub>, *J* = 7.0 Hz), 2.28 (s, 3H, 3-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 3.93 (t, 2H, OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, *J* = 6.5 Hz), 4.94 (s, 2H, NCH<sub>2</sub>CO), 6.76 (s, 1H, Ar), 6.80–6.86 (m, 5H, Ar), 7.26–7.30 (m, 1H, Ar), 7.40 (d, 2H, Ar, *J* = 8.9 Hz), 8.76 (exch br s, 1H, NH). MS (ESI) *m*/*z* 436.23 [M + H]<sup>+</sup>. Anal. (C<sub>25</sub>H<sub>29</sub>N<sub>3</sub>O<sub>4</sub>) C, H, N.

*N*-(3,4-Dimethoxyphenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14l). Yield = 74%; mp =  $57-59 \,^{\circ}\text{C}$  (EtOH). IR (cm<sup>-1</sup>) 3298 (NH), 1708 (CO), 1640 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.29 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 3.84 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 4.95 (s, 2H, NCH<sub>2</sub>CO), 6.72 (d, 1H, Ar, J = 8.6 Hz), 6.78 (s, 2H, Ar), 6.82–6.88 (m, 3H, Ar), 7.25–7.29 (m, 1H, Ar), 7.32 (d, 1H, Ar, J = 2.2 Hz), 8.93 (exch br s, 1H, NH). MS (ESI) m/z 424.19 [M + H]<sup>+</sup>. Anal. (C<sub>23</sub>H<sub>25</sub>N<sub>3</sub>O<sub>5</sub>) C, H, N.

*N*-Benzo[1,3]dioxol-5-yl-2-[5-(3-methoxybenzyl)-3-methyl-6oxo-6*H*-pyridazin-1-yl]-acetamide (14m). Yield = 98%; oil. IR (cm<sup>-1</sup>) 3300 (NH), 1707 (CO), 1643 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.28 (s, 3H, 3-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.89 (s, 2H, CH<sub>2</sub>-Ar), 4.93 (s, 2H, NCH<sub>2</sub>CO), 5.92 (s, 2H, O-CH<sub>2</sub>-O), 6.65 (d, 1H, Ar, *J* = 8.3 Hz), 6.77–6.85 (m, 5H, Ar), 7.21 (d, 1H, Ar, *J* = 2.0 Hz), 7.25–7.29 (m, 1H, Ar), 9.04 (exch br s, 1H, NH). MS (ESI) *m*/*z* 408.16 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>21</sub>N<sub>3</sub>O<sub>5</sub>) C, H, N.

**2-**[**5-**(**3-**Methoxybenzyl)-**3-**methyl-**6-**oxo-**6***H*-pyridazin-1-yl]-*N*-(**4-**trifluoromethylphenyl)-acetamide (14n). Yield = 80%; mp = 175–176 °C (EtOH). IR (cm<sup>-1</sup>) 3297(NH), 1708 (CO), 1646 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.33 (s, 3H, 3-CH<sub>3</sub>), 3.79 (s, 3H, OCH<sub>3</sub>), 3.91 (s, 2H, CH<sub>2</sub>-Ar), 5.00 (s, 2H, NCH<sub>2</sub>CO), 6.81–6.85 (m, 3H, Ar), 6.89 (s, 1H, Ar), 7.25–7.29 (m, 1H, Ar), 7.38 (d, 2H, Ar, *J* = 8.7 Hz), 7.47 (d, 2H, Ar, *J* = 8.7 Hz), 9.62 (exch br s, 1H, NH). MS (ESI) *m*/*z* 432.16 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>20</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub>) C, H, N.

**2-[5-(3-Methoxybenzyl)-3-methyl-6-oxo-6***H*-pyridazin-1-yl]-*N*-(4-trifluoromethoxyphenyl)-acetamide (140). Yield = 87%; mp = 168–169 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1708 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.31 (s, 3H, 3-CH<sub>3</sub>), 3.78 (s, 3H, OCH<sub>3</sub>), 3.89 (s, 2H, CH<sub>2</sub>-Ar), 5.00 (s, 2H, NCH<sub>2</sub>CO), 6.79–6.83 (m, 3H, Ar), 6.87 (s, 1H, Ar), 7.00 (d, 2H, Ar, J = 8.6 Hz), 7.24– 7.28 (m, 1H, Ar), 7.41 (d, 2H, Ar, J = 9.0 Hz), 9.54 (exch br s, 1H, NH). MS (ESI) *m*/*z* 448.15 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>20</sub>F<sub>3</sub>N<sub>3</sub>O<sub>4</sub>) C, H, N.

**2-[5-(3-Methoxybenzyl)-3-methyl-6-oxo-***6H***-pyridazin-1-yl]**-*N*-(**4-nitrophenyl)-acetamide** (**14p**). Yield = 49%; mp = 165–166 °C (EtOH). IR (cm<sup>-1</sup>) 3298 (NH), 1708 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.34 (s, 3H, 3-CH<sub>3</sub>), 3.79 (s, 3H, OCH<sub>3</sub>), 3.92 (s, 2H, CH<sub>2</sub>-Ar), 5.01 (s, 2H, NCH<sub>2</sub>CO), 6.80–6.84 (m, 3H, Ar), 6.94 (s, 1H, Ar), 7.25–7.28 (m, 1H, Ar), 7.49 (d, 2H, Ar, *J* = 9.2 Hz), 7.99 (d, 2H, Ar, *J* = 9.2 Hz), 9.92 (exch br s, 1H, NH). MS (ESI) *m/z* 409.15 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>N<sub>4</sub>O<sub>5</sub>) C, H, N.

*N*-(4-Cyanophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14q). Yield = 55%; mp = 156– 158 °C (EtOH). IR (cm<sup>-1</sup>) 3285 (NH), 2221 (CN), 1716 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.32 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.91 (s, 2H, CH<sub>2</sub>-Ar), 4.98 (s, 2H, NCH<sub>2</sub>CO), 6.80–6.88 (m, 4H, Ar), 7.26–7.30 (m, 1H, Ar), 7.50–7.57 (m, 4H, Ar), 9.60 (exch br s, 1H, NH). MS (ESI) *m*/*z* 389.16 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>20</sub>N<sub>4</sub>O<sub>3</sub>) C, H, N.

**2-**[**5-**(**3-**Methoxybenzyl)-**3-**methyl-**6-**oxo-**6***H*-pyridazin-1-yl]-*N*-(**4-**methoxyphenyl)-*N*-methyl-acetamide (14r). Yield = 45%; mp = 125-127 °C (EtOH). IR (cm<sup>-1</sup>) 1709 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.21 (s, 3H, 3-CH<sub>3</sub>), 3.32 (s, 3H, CH<sub>3</sub>N), 3.82 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 2H, CH<sub>2</sub>-Ar), 4.64 (s, 2H, NCH<sub>2</sub>CO), 6.62 (s, 1H, Ar), 6.77 (s, 1H, Ar), 6.80-6.85 (m, 2H, Ar), 7.26-7.30 (m, 3H, Ar), 7.60 (d, 2H, Ar, *J* = 8.4 Hz). MS (ESI) *m*/*z* 456.09 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>22</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

*N*-(4-Bromobenzyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14s). Yield = 97%; mp = 184– 185 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1708 (CO), 1644 (CO). <sup>1</sup>H NMR (DMSO- $d_6$ )  $\delta$  2.21 (s, 3H, 3-CH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 2H, CH<sub>2</sub>-Ar), 4.27 (d, 2H, CH<sub>2</sub>Ar), 4.68 (s, 2H, NCH<sub>2</sub>CO), 6.80–6.86 (m, 3H, Ar), 7.05 (s, 1H, Ar), 7.23– 7.26 (m, 3H, Ar), 7.51 (d, 2H, Ar, J = 8.3 Hz), 8.63 (exch br t, 1H, NH, J = 5.8 Hz). MS (ESI) m/z 456.09 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>22</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

[5-(3-Methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]acetic Acid 4-Bromo-phenyl Ester (14t). Yield = 97%; mp = 111-112 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1745 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.27 (s, 3H, 3-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 5.10 (s, 2H, NCH<sub>2</sub>COO), 6.73 (s, 1H, Ar), 6.80 (s, 1H, Ar), 6.83-6.86 (m, 2H, Ar), 7.05 (d, 2H, Ar, J = 8.7 Hz), 7.26–7.30 (m, 1H, Ar), 7.50 (d, 2H, Ar, J = 8.7 Hz). MS (ESI) m/z 443.06 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>19</sub>BrN<sub>2</sub>O<sub>4</sub>) C, H, N.

**4-(3-Methoxybenzyl)-6-methyl-2-[2-(4-methylpiperazin-1-yl)-2-oxo-ethyl]-pyridazin-3(2***H***)-one (14u). Yield = 62%; oil. IR (cm<sup>-1</sup>) 1673 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>) \delta 2.25 (s, 3H, 3-CH<sub>3</sub>), 2.55 (s, 3H, CH<sub>3</sub>N), 2.74–2.81 (m, 4H, Ar), 3.77–3.82 (m, 7H (4H, Ar; 3H, OCH<sub>3</sub>)), 3.87 (s, 2H, CH<sub>2</sub>-Ar), 4.96 (s, 2H, NCH<sub>2</sub>CO), 6.69 (s, 1H, Ar), 6.80 (s, 1H, Ar), 6.82–6.85 (m, 2H, Ar), 7.26–7.30 (m, 1H, Ar). MS (ESI)** *m***/***z* **371.21 [M + H]<sup>+</sup>. Anal. (C<sub>20</sub>H<sub>26</sub>N<sub>4</sub>O<sub>3</sub>) C, H, N.** 

*N*-(4-Butoxyphenyl)-2-[5-(4-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14v). Yield = 60%; mp = 160– 161 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1707 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.98 (t, 3H, O(CH<sub>2</sub>)<sub>3</sub>*CH<sub>3</sub>*, *J* = 7.4 Hz), 1.45–1.54 (m, 2H, OCH<sub>2</sub>CH<sub>2</sub>*CH*<sub>2</sub>CH<sub>3</sub>), 1.73–1.80 (m, 2H, OCH<sub>2</sub>*CH*<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 2.28 (s, 3H, 3-CH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 3.94 (t, 2H, O*CH*<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, *J* = 6.5 Hz), 4.93 (s, 2H, NCH<sub>2</sub>CO), 6.74 (s, 1H, Ar), 6.82 (d, 2H, Ar, *J* = 9.0 Hz), 6.90 (d, 2H, Ar, *J* = 8.6 Hz), 7.17 (d, 2H, Ar, *J* = 8.5 Hz), 7.39 (d, 2H, Ar, *J* = 9.0 Hz), 8.67 (exch br s, 1H, NH). MS (ESI) *m*/*z* 436.23 [M + H]<sup>+</sup>. Anal. (C<sub>25</sub>H<sub>29</sub>N<sub>3</sub>O<sub>4</sub>) C, H, N.

*N*-(**4-Butoxyphenyl**)-**2**-[**5**-(**4-chlorobenzyl**)-**3-methyl-6-oxo-6H-pyridazin-1-yl]-acetamide** (14w). Yield = 70% mp = 146– 147 °C (EtOH). IR (cm<sup>-1</sup>) 3298 (NH), 1708 (CO), 1642 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.99 (t, 3H, O(CH<sub>2</sub>)<sub>3</sub>*CH*<sub>3</sub>, *J* = 7.4 Hz), 1.49 (sext, 2H, OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, *J* = 7.5 Hz), 1.76 (quint, 2H, OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, *J* = 7.0 Hz), 2.29 (s, 3H, 3-CH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 3.93 (t, 2H, CH<sub>2</sub>-O, *J* = 6.5 Hz), 4.94 (s, 2H, NCH<sub>2</sub>CO), 6.76 (s, 1H, Ar), 6.81 (d, 2H, Ar, *J* = 9.0 Hz), 7.18 (d, 2H, Ar, *J* = 8.4 Hz), 7.32 (d, 2H, Ar, *J* = 8.4 Hz), 7.38 (d, 2H, Ar, *J* = 9.0 Hz), 8.69 (exch br s, 1H, NH). MS (ESI) *m/z* 440.17 [M + H]<sup>+</sup>. Anal. (C<sub>24</sub>H<sub>26</sub>ClN<sub>3</sub>O<sub>3</sub>) C, H, N.

*N*-(4-Bromophenyl)-2-[5-(4-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-acetamide (14x). Yield = 73%; mp = 139– 141 °C (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1709 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.29 (s, 3H, 3-CH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 4.94 (s, 2H, NCH<sub>2</sub>CO), 6.78 (s, 1H, Ar), 6.90 (d, 2H, Ar, *J* = 8.6 Hz), 7.17 (d, 2H, Ar, *J* = 8.6 Hz), 7.40 (s, 4H, Ar), 9.01 (exch br s, 1H, NH). MS (ESI) *m*/*z* 442.08 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

General Procedure for 15a,b. Compounds 15a,b were obtained starting from 11a following the general procedure described for 12a, using the appropriate alkyl halide in the place of ethyl bromoacetate.

**3-[5-(3-Methoxybenzyl)-3-methyl-6-oxo-***6H***-pyridazin-1-yl]-propionic Acid Ethyl Ester (15a).** Yield = 86%; oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.26 (t, 3H,OCH<sub>2</sub>*CH*<sub>3</sub>, *J* = 7.1 Hz), 2.22 (s, 3H, 3-CH<sub>3</sub>), 2.84 (t, 2H, NCH<sub>2</sub>*CH*<sub>2</sub>COO, *J* = 7.2 Hz), 3.83 (s, 3H, OCH<sub>3</sub>), 3.87 (s, 2H, CH<sub>2</sub>-Ar), 4.17 (q, 2H, OCH<sub>2</sub>CH<sub>3</sub>, *J* = 7.1 Hz), 4.45 (t, 2H, N*CH*<sub>2</sub>CH<sub>2</sub>COO, *J* = 7.3 Hz), 6.64 (s, 1H, Ar), 6.80 (s, 1H, Ar), 6.83 (d, 2H, Ar, *J* = 7.9 Hz), 7.26–7.30 (m, 1H, Ar).

**2-**[**5-**(**3-**Methoxybenzyl)-**3-**methyl-**6-**oxo-**6***H*-pyridazin-**1-***y*]propionic Acid Ethyl Ester (15b). Yield = 85%; oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.22 (t, 3H,OCH<sub>2</sub>*CH*<sub>3</sub>, *J* = 7.1 Hz), 1.65 (d, 3H, *CH*<sub>3</sub>CHN, *J* = 7.2 Hz), 2.21 (s, 3H, 3-CH<sub>3</sub>), 3.78 (s, 3H, OCH<sub>3</sub>), 3.84 (s, 2H, CH<sub>2</sub>-Ar), 4.19 (q, 2H, O*CH*<sub>2</sub>CH<sub>3</sub>, *J* = 6.7 Hz), 5.51 (q, 1H, CH<sub>3</sub>*CH*N, *J* = 7.2 Hz), 6.64 (s, 1H, Ar), 6.77–6.81 (m, 3H, Ar), 7.22–7.26 (m, 1H, Ar).

General Procedure for 16a,b. Compounds 16a,b were obtained starting from 15a,b following the same general procedure described for 13a,d. After dilution with cold water and acidification with 6N HCl, the mixtures were extracted with  $CH_2Cl_2$  (3 × 15 mL) and the solvent evaporated in vacuo.

**3-[3-(3-Methoxybenzyl)-5-methyl-2-oxo-2***H***-pyridin-1-yl]-propionic Acid (16a).** Yield =96%; mp = 86–88 °C (EtOH). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.22 (s, 3H, 3-CH<sub>3</sub>), 2.89 (t, 2H, NCH<sub>2</sub>-*CH*<sub>2</sub>COO, *J* = 7.2 Hz), 3.81 (s, 3H, OCH<sub>3</sub>), 3.87 (s, 2H, CH<sub>2</sub>-Ar), 4.46 (t, 2H, NCH<sub>2</sub>COO, *J* = 7.2 Hz), 6.66 (s, 1H, Ar),

6.78 (s, 1H, Ar), 6.81–6.85 (m, 2H, Ar), 7.25–7.29 (m, 1H, Ar), 9.99 (exch br s, 1H, OH).

**2-**[**5-**(**3-**Methoxybenzyl)-**3-**methyl-**6-**oxo-**6***H*-pyridazin-**1-**yl]propionic Acid (16b). Yield = 85%; oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.71 (d, 3H, *CH*<sub>3</sub>CHN, *J* = 7.2 Hz), 2.25 (s, 3H, 3-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 5.54 (q, 1H, CH<sub>3</sub>*CH*N, *J* = 7.2 Hz), 6.67 (s, 1H, Ar), 6.79–6.85 (m, 3H, Ar), 7.26–7.29 (m, 1H, Ar).

General Procedure for 17a,b. Compounds 17a,b were obtained starting from 16a,b following the general procedure described for 14a-x. Purification of the final compounds was performed by column chromatography using cyclohexane/ethyl acetate 1:2 as eluent for compound 17a, cyclohexane/ethyl acetate 2:1 for compound 17b.

*N*-(4-Bromophenyl)-3-[3-(3-methoxybenzyl)-5-methyl-2-oxo-2*H*-pyridazin-1-yl]-propionamide (17a). Yield = 82%; mp = 123-125 °C (EtOH). IR (cm<sup>-1</sup>) 3297 (NH), 1710 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.27 (s, 3H, 3-CH<sub>3</sub>), 3.00 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>COO, J = 6.3 Hz), 3.80 (s, 3H, OCH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 4.54 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>COO, J = 6.3 Hz), 6.76-6.84 (m, 4H, Ar), 7.25 (t, 1H, Ar, J = 7.9 Hz), 7.40 (d, 2H, Ar, J = 8.8Hz), 7.50 (d, 2H, Ar, J = 8.8 Hz), 9.29 (exch br s, 1H, NH). MS (ESI) m/z 456.09 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>22</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

*N*-(4-Bromophenyl)-2-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6*H*-pyridazin-1-yl]-propionamide (17b). Yield = 53%; oil; IR (cm<sup>-1</sup>) 3300 (NH), 1709 (CO), 1643 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.71 (d, 3H, *CH*<sub>3</sub>CHN, *J* = 7.1 Hz), 2.31 (s, 3H, 3-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 5.71 (q, 1H, CH<sub>3</sub>*CHN*, *J* = 7.0 Hz), 6.79-6.85 (m, 4H, Ar), 7.25-7.29 (m, 1H, Ar), 7.35-7.36 (m, 4H, Ar), 9.18 (exch br s, 1H, NH). MS (ESI) *m*/*z* 456.09 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>22</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

General Procedure for 18a,b. Compounds 18a,b were obtained starting from 11 following the procedure described for 12a. The final compounds were purified by column chromatography using cyclohexane/ethyl acetate 2:1 as eluent for compound 18a and cyclohexane/ethyl acetate 1:1 for compound 18b.

**2-(4-Bromobenzyl)-4-(3-methoxybenzyl)-6-methyl-pyridazin-3**(*2H*)-one (18a). Yield = 68%; mp = 112–114 °C (EtOH). IR (cm<sup>-1</sup>) 1638 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.23 (s, 3H, 6-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 2H, CH<sub>2</sub>-Ar), 5.25 (s, 2H, NCH<sub>2</sub>Ar), 6.64 (s, 1H, Ar), 6.78–6.85 (m, 3H, Ar), 7.26–7.30 (m, 1H, Ar), 7.35 (d, 2H, Ar, J = 8.4 Hz), 7.45–7.47 (m, 2H, Ar). MS (ESI) *m*/*z* 399.07 [M + H]<sup>+</sup>. Anal. (C<sub>20</sub>H<sub>19</sub>BrN<sub>2</sub>O<sub>2</sub>) C, H, N.

**2-[2-(4-Bromophenyl)-2-oxo-ethyl]-4-(3-methoxybenzyl)-6methyl-pyridazin-3**(*2H*)-**one** (18b). Yield = 95%; oil. IR (cm<sup>-1</sup>) 1715 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.25 (s, 3H, 6-CH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 3.89 (s, 2H, CH<sub>2</sub>-Ar), 5.52 (s, 2H, NCH<sub>2</sub>CO), 6.72 (s, 1H, Ar), 6.80 (s, 1H, Ar), 6.84 (d, 2H, Ar, *J* = 7.9 Hz), 7.27–7.31 (m, 1H, Ar), 7.66 (d, 2H, Ar, *J* = 8.6 Hz), 7.88 (d, 2H, Ar, *J* = 8.5 Hz). MS (ESI) *m*/*z* 427.07 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>19</sub>BrN<sub>2</sub>O<sub>3</sub>) C, H, N.

General Procedure for 19a,b. A mixture of 11a (0.32 mmol), 40% formaldehyde (3 mL), and 33% NH<sub>3</sub> (1.5 mL) in dioxane (1.5-2 mL) was heated at 50 °C for 1 h. The solvent was then evaporated in vacuo, and the residue was extracted with  $CH_2Cl_2$ (3 × 15 mL). The organic layer was dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated to afford an oil.

For compound **19a**, the residual oil was dissolved in 2 mL of anhydrous  $CH_2Cl_2$  and 4-bromophenyl isocyanate (0.35 mmol) was added. The mixture was stirred at room temperature for 12 h and then the solid residue was filtered off and the solution was evaporated in vacuo to afford compound **19a**, which was purified by flash chromatography using cyclohexane/ethyl acetate 3:1 as eluent.

For compound **19b**, the residual oil was dissolved in 3 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub> and, after cooling (0 °C), 4-bromobenzoyl chloride (0.44 mmol) was added and the mixture was stirred at 0 °C for 6 h. Finally, the residue was washed with cold 0.5 N NaOH (3 × 10 mL) and with cold water (2 × 10 mL). Evaporation of the organic layer afforded compound **19b**, which was purified by flash chromatography using cyclohexane/ethyl acetate 3:1 as eluent.

**1-(4-Bromophenyl)-3-[5-(3-methoxybenzyl)-3-methyl-6-oxo-6H-pyridazin-1-ylmethyl]-urea (19a).** Yield = 27%; mp = 112– 114 °C (EtOH). IR (cm<sup>-1</sup>) 3270 (NH), 3265 (NH), 1708 (CO), 1630 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.25 (s, 3H, 3-CH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 6.13 (s, 2H, NCH<sub>2</sub>N), 6.68 (s, 1H, Ar), 6.78–6.81 (m, 1H, Ar), 6.83–6.86 (m, 2H, Ar), 6.99 (exch br s, 1H, NH), 7.27–7.31 (m, 3H, Ar), 7.41 (m, 2H, Ar, *J* = 8.8 Hz). MS (ESI) *m*/*z* 458.07 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>21</sub>BrN<sub>4</sub>O<sub>3</sub>) C, H, N.

**4-Bromo-***N*-[**5**-(**3-methoxybenzyl**)-**3-methyl**-**6-oxo**-*6H*-**pyridazin-1-ylmethyl**]-**benzamide** (**19b**). Yield = 30%; oil. IR (cm<sup>-1</sup>) 3290 (NH), 1708 (CO), 1640 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.26 (s, 3H, 3-CH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 3.90 (s, 2H, CH<sub>2</sub>-Ar), 6.29 (s, 2H, NCH<sub>2</sub>N), 6.69 (s, 1H, Ar), 6.81–6.87 (m, 3H, Ar), 7.28–7.32 (m, 1H, Ar), 7.58 (d, 2H, Ar, *J* = 8.6 Hz), 7.94 (d, 2H, Ar, *J* = 8.6 Hz). MS (ESI) *m*/*z* 443.06 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>20</sub>-BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

**2-(2-Hydroxyethyl)-4-(3-methoxybenzyl)-6-methyl-pyridazin-3(2***H***)-one (20). To a refluxed mixture of compound 12a (0.47 mmol) and NaBH<sub>4</sub> (2.64 mmol) in anhydrous THF (6 mL), CH<sub>3</sub>OH (1.45 mL) was slowly added. After stirring for 1 h at 60 °C, the mixture was concentrated in vacuo, diluted with cold water (10–15 mL), and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL). Evaporation of the solvent afforded the final compound (20). Yield = 95%; oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) \delta 2.23 (s, 3H, 6-CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 2H, CH<sub>2</sub>-Ar), 4.00 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>OH,** *J* **= 5.1 Hz), 4.35 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>OH,** *J* **= 5.0 Hz), 4.66 (exch br s, 1H, CH<sub>2</sub>OH), 6.68 (s, 1H, Ar), 6.78–6.84 (m, 3H, Ar), 7.25–7.29 (m, 1H, Ar).** 

**2-[2-(4-Bromophenoxy)-ethyl]-4-(3-methoxybenzyl)-6-methyl-pyridazin-3**(*2H*)-one (21). Compound 21 was obtained starting from 20 following the general procedure described for 9. Compound 21 was purified by flash chromatography using toluene/ethyl acetate 8:2 as eluent. Yield = 26%; mp = 82– 83 °C (EtOH). IR (cm<sup>-1</sup>) 1643 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.24 (s, 3H, 6-CH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.87 (s, 2H, CH<sub>2</sub>-Ar), 4.36 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>O, *J* = 5.9 Hz), 4.54 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>O, *J* = 5.8 Hz), 6.66 (s, 1H, Ar), 6.79–6.86 (m, 5H, Ar), 7.27–7.35 (m, 1H, Ar), 7.35–7.37 (m, 2H, Ar). MS (ESI) *m*/*z* 431.08 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>21</sub>BrN<sub>2</sub>O<sub>3</sub>) C, H, N.

Methanesulfonic Acid 2-[5-(3-Methoxybenzyl)-3-methyl-6oxo-6*H*-pyridazin-1-yl]-ethyl Ester (22). To a cooled (0 °C) and stirred solution of 20 (0.47 mmol) and pyridine (0.5 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (2 mL), methanesulfonyl chloride (0.61 mmol) was added dropwise, and the mixture was stirred at room temperature for 4 h. Then ice cold water was added and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL): evaporation of the solvent afforded the desired compound. Yield =85%; oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.22 (s, 3H, 3-CH<sub>3</sub>), 2.98 (s, 3H, CH<sub>3</sub>SO<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 2H, CH<sub>2</sub>-Ar), 3.87–3.90 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>O), 4.45 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>O, *J* = 6.6 Hz), 6.66 (s, 1H, Ar), 6.78 (s, 1H, Ar), 6.82 (d, 2H, Ar, *J* = 8.1 Hz), 7.24–7.28 (m, 1H, Ar).

**2-[2-(4-Bromophenylamino)-ethyl]-4-(3-methoxybenzyl)-6methyl-pyridazin-3(***2H***)-one (23).** A solution of **22** (0.4 mmol) and 4-bromoaniline (0.8 mmol) in 2-propanol (2 mL) was heated under stirring for 6 h at 60 °C. After the mixture was concentrated in vacuo and cold water (30 mL) was added, the suspension was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL). Evaporation of the solvent afforded the final compound **23**, which was purified by column chromatography using toluene/ethyl acetate 8:2 as eluent. Yield = 58%; mp = 86–88 °C (EtOH). IR (cm<sup>-1</sup>) 3350 (NH), 1643 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.24 (s, 3H, 6-CH<sub>3</sub>), 3.58 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>NHAr, J = 5.7 Hz), 3.82 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 2H, CH<sub>2</sub>-Ar), 4.46 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>NHAr, J = 5.8 Hz), 6.65–6.68 (m, 3H, Ar), 6.79–6.86 (m, 3H, Ar), 7.27–7.31 (m, 3H, Ar). MS (ESI) *m*/*z* 428.10 [M + H]<sup>+</sup>. Anal. (C<sub>21</sub>H<sub>22</sub>BrN<sub>3</sub>O<sub>2</sub>) C, H, N. **2-(2-Aminoethyl)-4-(3-methoxybenzyl)-6-methyl-pyridazin-3**(*2H*)**-one (24).** A mixture of **22** (0.43 mmol) and 33% NH<sub>3</sub> (3 mL) in isopropanol (2 mL) was stirred at 60 °C for 3 h. After concentration of the solvent and dilution with cold water (20 mL), the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL). Evaporation of the solvent afforded desired compound **24**. Yield = 68%; oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.25 (s, 3H, 6-CH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.88 (s, 2H, CH<sub>2</sub>-Ar), 4.02–4.04 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>), 4.37 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, *J* = 4.9 Hz), 5.38 (exch br s, 2H, NH<sub>2</sub>), 6.69 (s, 1H, Ar), 6.79–6.85 (m, 3H, Ar), 7.26–7.30 (m, 1H, Ar).

General Procedure for 25a,b. Compounds 25a,b were obtained by starting from 24. For compound 25a, the general procedure reported for 6a-c was followed, and the final compound was purified by flash chromatography using cyclohexane/ethyl acetate 1:1 as eluent. For compound 25b, Et<sub>3</sub>N (1.8 mmol) and 4-bromobenzoyl chloride (1.43 mmol) were added to a cooled (0 °C) and stirred solution of 24 (0.73 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub>(2 mL), and the mixture was stirred at 0 °C for 6 h. The solid residue was filtered off, and the solution was washed with 6 N NaOH (3 × 10 mL) and then with cold water (2 × 10 mL). The organic layer was dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated in vacuo to afford compound 25b, which was purified by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH 99:1 as eluent.

**1-(4-Bromophenyl)-3-{2-[5-(3-methoxybenzyl)-3-methyl-6oxo-6***H***-pyridazin-1-yl]-ethyl}-urea (25a). Yield = 15%; mp = 114–115 °C (EtOH). IR (cm<sup>-1</sup>) 3270 (NH), 3265 (NH), 1705 (CO), 1630 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>) \delta 2.23 (s, 3H, 3-CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 3.87 (s, 2H, CH<sub>2</sub>-Ar), 4.47 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>NH, J = 5.2 Hz), 4.57 (t, 2H, NCH<sub>2</sub>CH<sub>2</sub>NH, J = 5.1 Hz), 6.69 (s, 1H, Ar), 6.78–6.84 (m, 3H, Ar), 7.04 (exch br s, 1H, NH), 7.25–7.28 (m, 3H, Ar), 7.41 (d, 2H, Ar, J = 8.8 Hz). MS (ESI) m/z 472.08 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>23</sub>BrN<sub>4</sub>O<sub>3</sub>) C, H, N.** 

**4-Bromo-***N*-{**2-**[**5-**(**3-methoxybenzyl**)-**3-methyl-6-oxo-**6*H*-**pyridazin-1-yl**]-**ethyl**}-**benzamide** (**25b**). Yield = 13%; oil. IR (cm<sup>-1</sup>) 3300 (NH), 1707 (CO), 1643 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.18 (s, 3H, 3-CH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.87 (s, 2H, CH<sub>2</sub>-Ar), 4.55 (t, 2H, NCH<sub>2</sub>*CH*<sub>2</sub>NH, *J* = 5.4 Hz), 4.70 (t, 2H, N*CH*<sub>2</sub>*CH*<sub>2</sub>NH, *J* = 5.4 Hz), 6.67 (s, 1H, Ar), 6.68–6.85 (m, 3H, Ar), 7.26 (d, 1H, Ar, *J* = 7.8 Hz), 7.56 (d, 2H, Ar, *J* = 8.5 Hz), 7.86 (d, 2H, Ar, *J* = 8.5 Hz). MS (ESI) *m*/*z* 457.08 [M + H]<sup>+</sup>. Anal. (C<sub>22</sub>H<sub>22</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

**2-(3-Methoxybenzyl)-6-methylpyridazin-3(2***H***)-one (27). Compound <b>27** was obtained starting from **26** following the general procedure described for **12a**. Compound **27** was purified by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH 9.9:0.1 as eluent. Yield = 86%; mp = 53-55 °C (cyclohexane). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.32 (s, 3H, CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 5.26 (s, 2H, CH<sub>2</sub>), 6.81-6.84 (m, 1H, Ar), 6.87 (d,1H, Ar, *J* = 9.4 Hz), 6.97-7.05 (m, 2H, Ar), 7.07 (d, 1H, Ar, *J* = 9.4 Hz), 7.22-7.26 (m, 1H, Ar).

**4-Amino-2-(3-methoxybenzyl)-6-methylpyridazin-3**(*2H*)-one (**28**). A suspension of **27** (0.78 mmol) and hydrazine hydrate (322 mmol) was stirred in a sealed tube at 180 °C for 12 h. After cooling, ice-cold water was added, the suspension was kept at 0 °C for 2 h, and the precipitate was filtered off. The solution was saturated with NH<sub>4</sub>Cl and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 25 mL). Removal of the solvent afforded a second batch of crude product. Yield = 89%; mp = 96–98 °C (EtOH). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.23 (s, 3H, CH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 4.85 (exch br s, 2H, NH<sub>2</sub>), 5.27 (s, 2H, CH<sub>2</sub>), 6.16 (s, 1H, Ar), 6.81–6.85 (m, 1H, Ar), 6.97–7.02 (m, 2H, Ar), 7.23–7.28 (m, 1H, Ar).

**4-Bromo-***N*-[**2**-(**3-methoxybenzyl**)-**6-methyl-3-oxo-2,3-dihydropyridazin-4-yl]benzamide (29).** Compound **29** was obtained from compound **28** following the procedure described for **25b**. The final compound was purified by flash chromatography using cyclohexane/ethyl acetate 3:1 as eluent. Yield = 28%; mp =  $162-164 \degree C$  (EtOH). IR (cm<sup>-1</sup>) 3300 (NH), 1709 (CO), 1644 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.40 (s, 3H, CH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 5.32 (s, 2H, CH<sub>2</sub>), 6.84–6.87 (m, 1H, Ar), 6.98–7.02 (m, 2H, Ar), 7.26–7.30 (m, 1H, Ar), 7.66 (d, 1H, Ar, J = 8.7 Hz), 7.80 (d, 2H, Ar, J = 8.6 Hz), 8.14 (s, 1H, Ar), 9.37 (exch br s, 1H, NH). MS (ESI) m/z 428.06 [M + H]<sup>+</sup>. Anal. (C<sub>20</sub>H<sub>18</sub>BrN<sub>3</sub>O<sub>3</sub>) C, H, N.

1-(4-Bromophenyl)-3-[2-(3-methoxybenzyl)-6-methyl-3-oxo-2,3-dihydropyridazin-4-yl]urea (30). Compound 30 was obtained from compound 28 following the general procedure reported for 6a-c. After dilution with cold water, the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL), and the solvent was evaporated in vacuo. Compound 30 was purified by column chromatography using first CH<sub>2</sub>Cl<sub>2</sub> to remove the 4-bromophenyl urea, followed by cyclohexane/ethyl acetate 2:1 as eluent. Yield = 56%; mp = 207-209 °C (EtOH). IR (cm<sup>-1</sup>) 3270 (NH), 3265 (NH), 1705 (CO), 1630 (CO). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.12 (s, 3H, CH<sub>3</sub>), 3.57 (s, 3H, OCH<sub>3</sub>), 5.08 (s, 2H, CH<sub>2</sub>), 6.51-6.58 (m, 3H, Ar), 6.97-7.04 (m, 3H, Ar), 7.23 (d, 2H, Ar, J = 8.7 Hz), 7.86 (s, 1H, Ar), 8.70 (exch br s, 1H, NH), 8,94 (exch, br,s, 1H, NH). MS (ESI) *m*/ *z* 443.07 [M + H]<sup>+</sup>. Anal. (C<sub>20</sub>H<sub>19</sub>BrN<sub>4</sub>O<sub>3</sub>) C, H, N.

**Biological Assays. Cell Culture.** Human HL-60 cells stably transfected with FPR1 (HL-60-FPR1), FPRL1 (HL-60-FPRL1), or FPRL2 (HL-60-FPRL2) were as previously described.<sup>25</sup> The transfected cells were cultured in RPMI supplemented with 10% heat inactivated fetal calf serum, 10 mM HEPES, 100  $\mu$ g/mL streptomycin, 100 U/mL penicillin, and G418 (1 mg/mL), as described previously.<sup>26</sup> Parent (wild-type) HL-60 cells were cultured under the same conditions but without G418.

**Isolation of Human Neutrophils.** Blood was collected from healthy donors in accordance with a protocol approved by the Institutional Review Board at Montana State University. Neutrophils were purified from the blood using dextran sedimentation, followed by Histopaque 1077 gradient separation and hypotonic lysis of red blood cells, as described previously.<sup>27</sup> Isolated neutrophils were washed twice and resuspended in HBSS without Ca<sup>2+</sup> and Mg<sup>2+</sup> (HBSS<sup>-</sup>). Neutrophil preparations were routinely >95% pure, as determined by light microscopy, and >98% viable, as determined by trypan blue exclusion.

Ca<sup>2+</sup> Mobilization Assay. Changes in intracellular Ca<sup>2+</sup> were measured with a FlexStation II scanning fluorometer using a FLIPR 3 calcium assay kit (Molecular Devices, Sunnyvale, CA) for human neutrophils and HL-60 cells. All active compounds were evaluated in parent (wild-type) HL-60 cells for supporting that the agonists are inactive in nontransfected cells. Human neutrophils or HL-60 cells, suspended in HBSS<sup>-</sup> containing 10 mM HEPES, were loaded with Fluo-4 AM dye (Invitrogen)  $(1.25 \,\mu g/mL \text{ final concentration})$  and incubated for 30 min in the dark at 37 °C. After dye loading, the cells were washed with HBSS<sup>-</sup> containing 10 mM HEPES, resuspended in HBSS containing 10 mM HEPES and Ca<sup>2+</sup> and Mg<sup>2+</sup> (HBSS<sup>+</sup>), and aliquotted into the wells of a flat-bottomed, half-area-well black microtiter plates ( $2 \times 10^{5}$  cells/well). The compound source plate contained dilutions of test compounds in HBSS<sup>+</sup>. Changes in fluorescence were monitored ( $\lambda_{ex} = 485 \text{ nm}, \lambda_{em} = 538 \text{ nm}$ ) every 5 s for 240 s at room temperature after automated addition of compounds. Maximum change in fluorescence, expressed in arbitrary units over baseline, was used to determine agonist response. Responses were normalized to the response induced by 5 nM fMLF (Sigma Chemical Co., St. Louis, MO) for HL-60-FPR1 and neutrophils, or 5 nM WKYMVm (Calbiochem, San Diego, CA) for HL-60-FPRL1, which were assigned a value of 100%. Curve fitting (5-6 points) and calculation of median effective concentration values (EC<sub>50</sub>) were performed by nonlinear regression analysis of the dose-response curves generated using Prism 5 (GraphPad Software, Inc., San Diego, CA).

**Chemotaxis Assay.** Neutrophils were suspended in HBSS<sup>+</sup> containing 2% (v/v) fetal bovine serum (FBS) ( $2 \times 10^6$  cells/mL), and chemotaxis was analyzed in 96-well ChemoTx chemotaxis chambers (Neuroprobe, Gaithersburg, MD), as described previously.<sup>27</sup> In brief, lower wells were loaded with

30  $\mu$ L of HBSS<sup>+</sup> containing 2% (v/v) FBS and the indicated concentrations of test compound, DMSO (negative control), and 1 nM fMLF as a positive control. The number of migrated cells was determined by measuring ATP in lysates of transmigrated cells using a luminescence-based assay (CellTiter-Glo; Promega, Madison, WI), and luminescence measurements were converted to absolute cell numbers by comparison of the values with standard curves obtained with known numbers of neutrophils. The results are expressed as percentage of negative control and were calculated as follows: (number of cells migrating in response to control medium) × 100. EC<sub>50</sub> values were determined by nonlinear regression analysis of the dose–response curves generated using Prism 5 software.

Acknowledgment. We thank Dr. Francois Boulay (Laboratoire de Biochimie et Biophysique des Systèmes Intégrés, CEA-Grenoble, France) for help in obtaining HL-60 cells transfected with FPR subtypes. This work was supported in part by NIH contract HHSN266200400009C and P20 RR020185. We also thank Dr. Brian Bothner and Navid Mohaved (Department of Chemistry & Biochemistry, Montana State University, Bozeman, MT) for their help in mass spectrometry analysis.

**Supporting Information Available:** Elemental analysis results for all target compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

#### References

- Lee, H. Y.; Bae, Y. S. The Anti-infective Peptide, Innate Defense-Regulator Peptide, Stimulates Neutrophil Chemotaxis Via a Formyl Peptide Receptor. *Biochem. Biophys. Res. Commun.* 2008, *369*, 573–578.
- (2) Gouin, J. P.; Hantsoo, L.; Kiekolt-Glaser, J. K. Immune Disregulation and Chronic Stress Among Older Adults: A Review. *Neuroimmunomodulation* **2008**, *15*, 251–259.
- (3) Jones, R. N. Resistance Patterns Among Nosocomial Phatogens: Trends Over the Past Few Years. *Chest* 2001, 119, 397S–4045S.
- (4) Zhang, L.; Falla, T. J. Host Defense Peptides for Use as Potential Therapeutics. *Curr. Opin. Invest. Drugs* 2009, 10, 164–171.
- (5) Selvatici, R.; Falzarano, S.; Mollica, A.; Spisani, S. Signal Transduction Pathways Triggered by Selective Formylpeptide Analogues in Human Neutrophils. *Eur. J. Pharmacol.* 2006, 534, 1–11.
- (6) Migeotte, I.; Communi, D.; Parmentier, M. Formylpeptide Receptors: A Promiscuous Subfamily of G Protein-Coupled Receptors Controlling Immune Responses. *Cytokine Growth Factor Rev.* 2006, 17, 501–519.
- (7) Schiffmann, E.; Showell, H. V.; Corcoran, B. A.; Ward, P. A.; Smith, E.; Becker, E. L. The Isolation and Partial Characterization of Neutrophil Chemotatic Factors from *Escherichia coli.*. J. Immunol. 1975, 114, 1831–1837.
- (8) Gao, J. L.; Lee, E. J.; Murphy, P. M. Impaired Antibacterial Host Defence in Mice Lacking the *N*-Formylpeptide Receptor. *J. Exp. Med.* **1999**, *189*, 657–662.
- (9) Le, Y.; Oppenheim, J. J.; Wang, J. M. Pleiotropic Roles of Formyl Peptide Receptors. *Cytokine Growth Factor Rev.* 2001, 12, 91–105.
- (10) Kilby, J. M.; Hopkins, S.; Venetta, T. M.; DiMassimo, B.; Cloud, G. A.; Lee, J. Y.; Alldredge, L.; Hunter, E.; Lambert, D.; Bolognesi, D.; Matthews, T.; Johnson, M. R.; Nowak, M. A.; Shaw, G. M.; Saag, M. S. Potent Suppression of HIV-1 Replication in Human by T-20 a Peptide Inhibitor of gp41-mediated Virus Entry. *Nat. Med.* **1998**, *4*, 1302–1307.

- (11) Le, Y.; Gong, W.; Tiffany, H. L.; Tumanov, A.; Nedospasov, S.; Shen, W.; Dunlop, N. M.; Gao, J. L.; Murphy, P. M.; Oppenheim, J. J.; Wang, J. M. Amyloid (beta)42 Activates a G-protein-coupled Chemoattractant Receptor, FPR-like-1. *J. Neurosci.* 2001, *21*, RC123.
- (12) Le, Y.; Yazawa, H.; Gong, W.; Yu, Z.; Ferrans, V. J.; Murphy, M. P.; Wang, J. M. The Neurotoxic Prion Peptide Fragment PrP(106–126) is a Chemotactic Agonist for the G-Protein-Coupled Receptor Formyl Peptide Receptor-like 1. J. Immunol. 2001, 66, 1448–1451.
- (13) Chiang, N.; Serhan, C. N.; Dahlén, S. E.; Drazen, J. M.; Hay, D. W.; Rovati, G. E.; Shimizu, T.; Yokomizo, T.; Brink, C. The Lipoxin Receptor ALX: Potent Ligand-Specific and Stereoselective Actions in Vivo. *Pharmacol Rev.* 2006, *58*, 463–487.
- (14) Perretti, M.; Chiang, N.; La, M.; Fierro, I. M.; Marullo, S.; Getting, S. J.; Solito, E.; Serhan, C. N. Endogenous Lipid- and Peptide-Derived Anti-inflammatory Pathways Generated with Glucocorticoid and Aspirin Treatment Activate the Lipoxin A4 Receptor. *Nat. Med.* 2002, 8, 1296–1302.
- Receptor. Nat. Med. 2002, 8, 1296–1302.
  (15) Edwards, B. S.; Bologa, C.; Young, S. M.; Balakin, V.; Prossnitz, E. R.; Savchuck, N. P.; Skalar, L. A.; Oprea, T. I. Integration of Virtual Screening with High-Throughput Flow Cytometry to Identify Novel Small Molecule Formyl Peptide Receptor Antagonists. Mol. Pharmacol. 2005, 68, 1301–1310.
- (16) Bae, Y. S.; Song, J. Y.; Kim, Y.; He, R.; Ye, R. D.; Kwak, J. Y.; Suh, P. G.; Ryu, S. H. Differential Activation of Formyl Peptide Receptor Signalling by Peptide Ligands. *Mol. Pharmacol.* 2003, 64, 841–847.
- (17) Le, Y.; Gong, W.; Li, B.; Dunlop, N. M.; Shen, W.; Su, S. B.; Ye, R. D.; Wang, J. M. Utilization of Two Seven-Transmembrane, G Protein-Coupled Receptors, Formyl Peptide Receptor-like 1 and Formyl Peptide Receptor, by the Synthetic Hexapeptide WKYMVm for Human Phagocyte Activation. J. Immunol. 1999, 163, 6777–6784.
- (18) Nanamori, M.; Cheng, X.; Mei, J.; Sang, H.; Xuan, Y.; Zhou, C.; Wang, M. W.; Ye, R. D. A novel nonpeptide ligand for formyl peptide receptor-like 1. *Mol. Pharmacol.* 2004, *66*, 1213–1222.
- (19) Frohn, M.; Xu, H.; Zou, X.; Chang, C.; McElvaine, M.; Plant, M. H.; Wong, M.; Tagari, P.; Hungate, R.; Burli, R. W. New "Chemical Probes" to Examine the Role of the hFPRL1 (or ALXR) Receptor in Inflammation. *Bioorg. Med. Chem. Lett.* 2007, 17, 6633–6637.
- (20) Burli, R. W.; Xu, H.; Zou, X.; Muller, K.; Golden, J.; Frohn, M.; Adlam, M.; Plant, M. H.; Wong, M.; McElvain, M.; Regal, K.; Viswanadhan, V. N.; Tagari, P.; Hungate, R. Potent hFPRL1 (ALXR) Agonists as Potential Anti-inflammatory Agents. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3713–3718.
- (21) Overend, W. G.; Wiggins, L. F. Conversion of Sucrose into Pyridazine Derivatives. II. 4-Amino-2-phenyl-6-methyl-3-pyridazinone, 4-Amino-2-(p-nitrophenyl)-6-methyl-3-pyridazone, and Their Sulfalinamide Derivatives. J. Chem. Soc. 1947, 549–554.
- (22) Meng, Q.; Hesse, M. Nitrogen–Nitrogen Bond Cleavage: a Route to Macrocyclic Dilactams. *Synlett* **1990**, *3*, 148–150.
- (23) Ismail, M. F.; El Khamry, A. A.; Shams, N. A.; El Sawi, O. M. Base-Catalyzed Condensation of Aromatic Aldehydes with 4,5-Dihydrogen-6-methylpyridazin-3(2H)-one. *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.* 1980, 19, 203–206.
- (24) Ismail, M. F.; Shams, N. A.; El Sawi, O. M. Synthesis of 3-Mercaptopyridazine Derivatives. *Synthesis* 1980, 410–412.
- (25) Ismail, M. F.; Shams, N. A.; El Sawi, O. M. Some Reaction with 4-(Arylmethyl)-6-Methylpyridazin-3(2H)-ones. *Egyptian J. Chem.* 1982, 24, 223–226.
- (26) Christophe, T.; Karlsson, A.; Rabiet, M. J.; Boulay, F.; Dahlgren, C. Phagocyte Activation by Trp-Lys-Tyr-Met-Val-Met, Acting Through FPRL1/LXA<sub>4</sub>R, Is Not Affected by Lipoxin A<sub>4</sub>. J. Immunol. 2002, 56, 470–476.
- (27) Schepetkin, I. A.; Kirpotina, L. N.; Khlebnikov, A. I.; Quinn, M. T. High-throughput Screening for Small-Molecule Activators of Neutrophils: Indentification of Novel *N*-Formyl Peptide Receptor Aagonists. *Mol. Pharmacol.* 2007, *71*, 1061–1074.