

Interaction of Methoxychlor and Related Compounds with Estrogen Receptor α and β , and Androgen Receptor: Structure-Activity Studies

KEVIN W. GAIDO, SUSAN C. MANESS, DONALD P. MCDONNELL, SHANGARA S. DEHAL, DAVID KUPFER, and STEPHEN SAFE

Chemical Industry Institute of Toxicology, Research Triangle Park, North Carolina (K.W.G., S.C.M.); Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, North Carolina (D.P.M.); Department of Pharmacology and Molecular Toxicology, University of Massachusetts Medical Center, Worcester, Massachusetts (S.S.D., D.K.); and Department of Veterinary Physiology and Pharmacology, Texas A&M University, College Station, Texas (S.S.)

Received March 10, 2000; accepted June 15, 2000

This paper is available online at <http://www.molpharm.org>

ABSTRACT

We previously demonstrated differential interactions of the methoxychlor metabolite 2,2-bis(*p*-hydroxyphenyl)-1,1,1-trichloroethane (HPTE) with estrogen receptor α (ER α), ER β , and the androgen receptor (AR). In this study, we characterize the ER α , ER β , and AR activity of structurally related methoxychlor metabolites. Human hepatoma cells (HepG2) were transiently transfected with human ER α , ER β , and AR plus an appropriate steroid-responsive luciferase reporter vector. After transfection, cells were treated with various concentrations of HPTE or structurally related compounds in the presence (for detecting antagonism) and absence (for detecting agonism) of 17 β -estradiol and dihydrotestosterone. The monohydroxy analog of methoxychlor, as well as monohydroxy and dihydroxy analogs

of 2,2-bis(*p*-hydroxyphenyl)-1,1-dichloroethylene, had ER α agonist activity and ER β and AR antagonist activity similar to HPTE. The trihydroxy metabolite of methoxychlor displayed only weak ER α agonist activity and did not alter ER β or AR activities. Replacement of the trichloroethane or dichloroethylene group with a methyl group resulted in a compound with ER α and ER β agonist activity that retained antiandrogenic activities. This study identifies some of the structural requirements for ER α and ER β activity and demonstrates the complexity involved in determining the mechanism of action of endocrine-active chemicals that simultaneously act as agonists or antagonists through one or more hormone receptors.

Methoxychlor [1,1,1-trichloro-2,2-bis(4-methoxyphenyl)ethane] is a chlorinated hydrocarbon pesticide structurally similar to dichlorodiphenyltrichloroethane [DDT; 1,1,1-trichloro-2,2-bis(chlorophenyl)ethane]. Like *o,p'*-DDT, methoxychlor is estrogenic in vivo (Bulger et al., 1978; Gray et al., 1989; Alm et al., 1996; Chapin et al., 1997; Cummings, 1997; Hall et al., 1997). However, methoxychlor has low affinity for the estrogen receptor (ER) and the in vivo estrogenic activity is caused by metabolism to phenolic estrogenic metabolites. The primary estrogenic metabolite of methoxychlor is 2,2-bis(*p*-hydroxyphenyl)-1,1,1-trichloroethane (HPTE). HPTE competes with estradiol for binding to ER, induces ornithine decarboxylase and uterotrophic activity in ovariectomized rats, and is approximately 100-fold more active than methoxychlor (Bulger et al., 1978; Ousterhout et al., 1981; Shelby et al., 1996).

The financial assistance of the National Institutes of Health (ES000834, ES09106, and ES04917) and the Texas Agricultural Experiment Station is gratefully acknowledged.

Estrogenic responses are mediated through two separate estrogen receptors, ER α and ER β . These two receptors have homologous DNA and ligand binding regions (Kuiper and Gustafsson, 1997; Tremblay et al., 1997; Ogawa et al., 1998), and most compounds have similar binding affinities and transcriptional activities with ER α and ER β (Kuiper et al., 1996, 1998; Mosselman et al., 1996; Tremblay et al., 1997).

We previously demonstrated that HPTE is an ER α agonist and an ER β antagonist in HepG2 human hepatoma cells transfected with estrogen-responsive reporter constructs (Gaido et al., 1999). This unique activity of HPTE makes it an ideal compound with which to evaluate the in vitro and in vivo differences in ER-subtype dependent responses. We have also shown that HPTE is an androgen receptor (AR) antagonist in vitro (Maness et al., 1998). The differential activity of HPTE with ER α , ER β , and AR may explain why some of the responses induced by methoxychlor in vivo differ from those induced by estradiol. For example, the ability of

ABBREVIATIONS: DDT, dichlorodiphenyltrichloroethane; ER, estrogen receptor; AR, androgen receptor; TLC, thin-layer chromatography; E2, 17 β -estradiol; GC-MS, gas chromatography-mass spectrometry; *p,p'*-DDE, 2,2-bis(*p*-hydroxyphenyl)-1,1-dichloroethylene; C3, complement 3; Luc, luciferase; CMV, cytomegalovirus; CPRG, chlorophenol red- β -D-galactopyranoside; HPTE, 2,2-bis(*p*-hydroxyphenyl)-1,1,1-trichloroethane; DHT, dihydrotestosterone; THC, tetrahydrochrysene.

methoxychlor to act as an ER antagonist in the ovary (Hall et al., 1997) may be caused by the high level of ER β expression relative to ER α in this tissue (Saunders et al., 1997).

The physiological consequences of a chemical that is an ER α agonist, an ER β antagonist, and an AR antagonist are unknown, and HPTE can serve as a model for investigating the effects of an agent that modulates multiple endocrine pathways. Additional studies with HPTE and structural analogs may lead to further insights on ligand specificity for ER α , ER β , and AR. Therefore we compared the ER α , ER β , and AR activity of HPTE and structural analogs and show that some chemicals similar in structure to HPTE also demonstrate unique ER α , ER β , and AR activity.

Materials and Methods

Chemicals. HPTE was synthesized by dissolving 1 g of methoxychlor (Aldrich Chemical Co., Milwaukee, WI) in 100 ml of methylene chloride and then treating with excess boron tribromide in methylene chloride (Aldrich) for 24 h. Water (5 ml) was carefully added, and crude HPTE was isolated in methylene chloride. The residue (0.8 g) was purified by preparative thin-layer chromatography (TLC). The resulting HPTE was >97% pure as determined by gas-liquid chromatography.

Monohydroxymethoxychlor was synthesized by dissolving 1.0 g of methoxychlor in methylene chloride. Approximately 1.5 mol equivalents of boron dibromide in methylene chloride was slowly added over a period of 1 to 2 h. The progress of demethylation was monitored by TLC. The monohydroxymethoxychlor metabolite was isolated by preparative TLC using hexane/acetone (92:8) as solvent. Yields of 250 to 300 mg were obtained and the products were greater than 98% pure as determined by gas chromatography-mass spectrometry (GC-MS).

Trihydroxymethoxychlor and the corresponding trimethoxymethoxychlor were synthesized by ChemSyn Labs (Lenexa, KS). Dimethoxy-DDE was synthesized by dissolving 1.0 g of methoxychlor in dimethyl sulfoxide. Anhydrous sodium bicarbonate (3.0 g) was added and the mixture was heated at 140°C for 1 h. The mixture was diluted with water and the dimethoxy-DDE product was isolated by extraction with chloroform. The crystalline residue from the chloroform extract (0.75 g) was greater than 98% pure as determined by GC-MS. Dihydroxy-DDE was prepared from 2,2-bis(*p*-hydroxyphenyl)-1,1-dichloroethylene (*p,p'*-DDE) following the same procedure as described above for HPTE. Dihydroxy-DDE was greater than 98% pure as determined by GC-MS. Monohydroxy-DDE was prepared from *p,p'*-DDE following the same procedure as described above for monohydroxymethoxychlor. Monohydroxy-DDE was greater than 98% pure as determined by GC-MS. All other chemicals were obtained from Sigma Chemical Co. (St. Louis, MO) and were \geq 97% pure.

Plating and Transfection. Transfection experiments were performed as described previously (Maness et al., 1998; Gaido et al., 1999). HepG2 human hepatoma cells (ATCC, Rockville, MD) were plated in triplicate in 24-well plates (Falcon Plastics, Oxnard, CA) at a density of 10^5 cells/well in complete medium consisting of phenol red-free Eagle's minimal essential medium (GIBCO/BRL, Grand Island, NY) supplemented with 10% resin-stripped fetal bovine serum (Hyclone, Logan, UT), 2% L-glutamine, and 0.1% sodium pyruvate. Cells were incubated overnight at 37°C in a humidified atmosphere of 5% CO $_2$ /air and then transfected after the Superfect procedure (Qiagen, Valencia, CA) with three plasmids. For detection of ER α activity, cells were transfected with human ER α expression plasmid, plus an estrogen-responsive complement 3-luciferase (C3-Luc) reporter plasmid, and a constitutively active cytomegalovirus (CMV)- β -galactosidase reporter plasmid (transfection and toxicity control) (Tzukerman et al., 1994; Gaido et al., 1999). For detection of ER β

activity, cells were transfected with a human ER β expression plasmid, a C3-Luc reporter plasmid, and CMV- β -galactosidase reporter plasmid (Gaido et al., 1999; Hall and McDonnell, 1999). For detection of AR activity, cells were transfected with a human AR expression plasmid, plus an androgen-responsive MMTV-Luc reporter plasmid, and CMV- β -galactosidase reporter plasmid (Maness et al., 1998). Transfected cells were rinsed with PBS and dosed with various concentrations of test chemical and dimethyl sulfoxide (vehicle control; Sigma) in complete medium. After a 24 h incubation, cells were rinsed with PBS and lysed with 65 μ l of lysing buffer (25 mM Tris-phosphate, pH 7.8, 2 mM 1,2-diaminocyclohexane-*N,N,N',N'*-tetraacetic acid, 10% glycerol, 0.5% Triton X-100, 2 mM dithiothreitol). Lysate was divided into two 96-well plates for luciferase and β -galactosidase determination.

Luciferase activity was determined by adding 100 μ l of Luc assay reagent (Promega, Madison, WI) to the first 96-well plate containing 20 μ l of lysate. Luminescence was determined immediately using a ML3000 microtiter plate luminometer (Dynatech Laboratories, Chantilly, VA).

β -Galactosidase activity was determined by adding 20 μ l of β -galactosidase assay reagent to 30 μ l of lysate in the second 96-well plate. β -Galactosidase assay reagent consisted of a 4 mg/ml solution of chlorophenol red- β -D-galactopyranoside (CPRG; Sigma) in 150 μ l of CPRG buffer (60 mM Na $_2$ HPO $_4$, 40 mM NaH $_2$ PO $_4$, 10 mM KCl, 1 mM MgSO $_4$, 50 mM β -mercaptoethanol, pH 7.8). Absorbance at 570 nm was determined over a 30 min period using a V_{max} kinetic microplate reader (Molecular Devices, Menlo Park, CA).

HepG2 cells lack detectable levels of endogenous ER α , ER β , and AR and in the absence of transfected receptor, Luc activity remains below the level of detection (data not shown). Background activity after receptor transfection averaged 150 ± 56 normalized Luc units with ER α , 31 ± 6 normalized Luc units with ER β , and 5 ± 1 normalized Luc units with AR. We have previously confirmed by Western analysis that ER α and ER β are expressed at equal concentrations under the conditions of our assay (Hall and McDonnell, 1999).

Statistical Analysis. Unless otherwise noted, values presented in this study represent the means \pm S.E. resulting from at least three separate experiments with triplicate wells for each treatment dose. Dose-response data were analyzed using the sigmoidal dose-response function of the graphical and statistical program Prism (GraphPad, San Diego, CA).

Results

We compared the activity of HPTE and structural analogs in HepG2 cells transfected with expression vectors for human ER α , ER β , and AR along with the appropriate reporter plasmid (Fig. 1). HepG2 cells were dosed with set concentrations of chemical alone and in combination with an inducing dose of either 17 β -estradiol (E2) or dihydrotestosterone (DHT) for determining antagonistic activity with ER α/β and AR, respectively. HPTE (Fig. 1C) exhibited ER α agonist, and ER β and AR antagonist activity as described previously (Maness et al., 1998; Gaido et al., 1999). HPTE does display some partial ER β agonist activity of approximately 13% of that obtained with a maximally inducing dose of estradiol (Maness et al., 1998). The monohydroxy metabolite of methoxychlor, as well as the mono- and dihydroxy analogs of *p,p'*-DDE (Fig. 1, B, H, and I), also had ER α agonist and ER β and AR antagonist activity. Bisphenol A exhibited ER α and ER β agonist activity but did not have antiandrogenic activity (Fig. 1K). Replacement of the trichloromethyl of HPTE or dichloromethylene group of dihydroxy-DDE results in a conversion from ER β antagonist activity to full ER β agonist activity but retains ER α agonist and AR antagonist activity (Fig. 1, C and




Fig. 1. ER α , ER β , and AR activity of methoxychlor analogs. HepG2 cells were transiently transfected with expression plasmids for human ER α , ER β , and AR plus C3-Luc and a constitutively active β -galactosidase expression plasmid (transfection and toxicity control). Cells were treated with increasing concentrations of methoxychlor analogs alone for detecting agonist activity (hatched bars) and with an inducing dose of either estradiol or DHT for detecting antagonist activity (shaded bars). Luciferase activity was normalized to β -galactosidase activity. Values represent the means \pm S.E. of three separate experiments and are presented as percentage response, with 100% activity defined as the activity achieved with 10^{-7} M estradiol for ER α and ER β , and 10^{-7} M DHT for AR. The abscissa represents log molar concentration. Agonists cause an increase in percentage response (rising hatched bars) with increasing concentration, whereas antagonists cause a decrease in percentage response (declining shaded bars) with increasing concentration.

I versus L). Trimethoxy and trihydroxy ring substituted compounds (Fig. 1, D and E) exhibited minimal ER α agonist activity and did not affect ER β or AR-dependent responses.

Concentration-response curves for selected ER α and ER β agonists are presented in Fig. 2, A and B. EC₅₀ values for ER α and ER β agonist activity are presented in Table 1. HPTE and dihydroxy-DDE were most potent as ER α agonists and were approximately 17- and 25-fold less potent, respectively, than estradiol. HPTE and dihydroxy-DDE were followed in ER α agonist potency by monohydroxy methoxychlor, bisphenol A, monohydroxy-DDE, bishydroxyphenylmethane, and bishydroxyphenylethane. Bishydroxyphenylmethane and bishydroxyphenylethane were equally potent as ER β agonists and were approximately 285-fold less potent than estradiol.

We characterized the ER β antagonist activity of selected compounds in HepG2 cells by determining the effect of various concentrations across a complete estradiol dose-response range (Fig. 3, A-C). Each of the tested compounds caused parallel shifts in the estradiol dose-response curve, indicating competitive antagonism. Schild regression analyses were performed and equilibrium dissociation (K_B) values determined (Table 1). HPTE, monohydroxy methoxychlor, monohydroxy-DDE, and dihydroxy-DDE demonstrated relatively similar antagonist potencies.

Similar experiments were performed to characterize AR antagonist activity (Fig. 4, Table 1). HPTE, dihydroxy-DDE, and *p,p'*-DDE demonstrated similar AR antagonist poten-

cies. Monohydroxy-DDE, bishydroxyphenylmethane, and bishydroxyphenylethane were approximately 3- to 5-fold less potent than HPTE, dihydroxy-DDE, and *p,p'*-DDE.

Discussion

ER α and ER β bind structurally diverse classes of chemicals, and it is difficult to compare structure-dependent receptor binding or transactivation between chemical classes. In contrast, structure-activity correlations within a single structural class such as the triphenylethanes can be used to design ER agonists and antagonists (e.g., tamoxifen) for clinical applications as selective estrogen receptor modulators. For a series of mono-, di-, and trihydroxyphenyl ethane/ethylene analogs structurally related to methoxychlor, the order of potency as ER α agonists was dihydroxyphenyl > monohydroxyphenyl > trihydroxyphenyl, suggesting that the optimal structure for ER α agonist activity contained two *p*-hydroxyphenyl groups substituted at a single ethane or ethylene carbon atom (i.e., bis-substitution). In contrast, the fully methylated metabolites (e.g., Fig. 1, A versus C; Fig. 1, G versus D) were significantly less active as ER α agonists; this is consistent with previous studies with methoxychlor and HPTE (Bulger et al., 1978; Ousterhout et al., 1981; Shelby et al., 1996). A similar pattern was observed for the substituted diphenylmethane analogs even though only a limited number of these compounds were tested (Fig. 1, J and N versus M).

With few exceptions (Fig. 1, K and L), the bishydroxy/methoxyphenyl ethane or ethylene analogs were not significant ER β agonists, and only two bishydroxyphenylmethanes (Fig. 1, J and N) induced measurable ER β -dependent reporter gene activity. Thus, our results demonstrate that this series of bishydroxyphenyl-substituted ethanes, ethylenes, and methanes are preferential ER α agonists and exhibit weak to nondetectable ER β agonist activity. These results are unique because previous studies for ER subtype-dependent ligand binding and transactivation report similar ER α and ER β activity for various structural classes of estrogenic compounds (Kuiper et al., 1996, 1998; Mosselman et al., 1996; Tremblay et al., 1997).

The results of our studies also demonstrate that methoxychlor and structurally related analogs exhibit minimal ER α antagonist activity but that three compounds (Fig. 1, B, H,




Fig. 2. Dose response of selected methoxychlor analogs with ER α (A) and ER β (B). Experiments were performed as described in Fig. 1. Luciferase activity was normalized to β -galactosidase activity. Values represent the means \pm S.E. of three separate experiments. ■, E2; ●, monohydroxy-methoxychlor; △, monohydroxy-DDE; ◇, dihydroxy-DDE; ▽, bishydroxyphenyl methane; ▲, bishydroxyphenyl ethane.

TABLE 1

A comparison of the agonist and antagonist potencies of methoxychlor analogs with ER α , ER β , and AR

Chemicals	ER α		ER β		AR	
	EC ₅₀	K _B	EC ₅₀	K _B	EC ₅₀	K _B
$\times 10^{-8}$ M						
Estradiol	0.3		0.7			
DHT					1.0	
HPTE	5.1 ^a		3.0 ^a			31.6
Monohydroxy-methoxychlor	19.8		7.2			
Monohydroxy-DDE	67.0		8.5			100.0
Dihydroxy-DDE	7.4		5.1			29.4
Bishydroxyphenylmethane	150.0		200.0			142.0
Bishydroxyphenylethane	160.0		210.0			136.0
<i>p,p'</i> -DDE						36.9 ^b
Bisphenol A	64.0 ^c		89.0 ^c			

^a From Gaido et al. (1999).

^b From Maness et al. (1998).

^c From Gould et al. (1998).

and I) in addition to HPTE (Fig. 1C) are highly effective ER β antagonists. Structural features required for this response include bis(4-hydroxyphenyl) or bis(4-hydroxyphenyl),(4-methoxyphenyl) groups attached to chlorine-substituted ethane/ethylene moieties. Additional compounds are required to more accurately define structural requirements for ER β antagonist activities; however, our results clearly demonstrate remarkable structure-dependent differences among these compounds for activity as ER β antagonists.

Several compounds investigated in this study exhibited antiandrogenic activity. Four analogs that were ER β antagonists (Fig. 1, B, C, H, and I) were among the most active

antiandrogens and exhibited activity similar to that observed for *p,p'*-DDE. Interestingly, both *p,p'*-DDE (Fig. 1F) and dihydroxy-DDE (Fig. 1I) exhibited similar antiandrogenic activities, and the interchange of two *p*-chloro and two *p*-hydroxyl substituents had minimal effects on this AR response. In contrast, the two *p*-hydroxyl groups (but not *p*-chloro substituents) conferred both ER α agonist and ER β antagonist activity on dihydroxy-DDE, demonstrating that subtle substituent changes can affect some but not all ligand-activated hormone receptor action.

Methoxy-DDE and monohydroxy-DDE are impurities in technical grade methoxychlor (Bulger et al., 1985), and dihydroxy-DDE is formed during the metabolism of methoxychlor in mice (Kapoor et al., 1970). HPTE, monohydroxy-methoxychlor, monohydroxy-DDE, and dihydroxy-DDE have previously been shown to compete with estradiol for ER binding in vitro and demonstrate uterotrophic activity in vivo (Bulger et al., 1978, 1985; Ousterhout et al., 1981). Thus, exposure to methoxychlor results in a complex interaction of multiple metabolites with different activities at ER α , ER β , and AR.

The molecular mechanism by which a ligand can act as an ER α agonist and an ER β antagonist is of both toxicological and pharmacological interest. The overall structure of the ER β ligand-binding domain is very similar to that of ER α (Pike et al., 1999), and most compounds demonstrate similar binding affinities and transcriptional activities with ER α and ER β (Kuiper et al., 1996, 1997; Mosselman et al., 1996; Tremblay et al., 1997). The helix 12 region present on both receptors plays an important role in the mechanism of ER action (Darimont et al., 1998). This region folds over the ligand binding pocket and exposes a region on both receptors involved in coactivator binding. ER α and ER β antagonists such as raloxifene and hydroxytamoxifen contain bulky constituents that reposition helix 12 and block receptor interaction with coactivators (Brzozowski et al., 1997; Pike et al., 1999). HPTE analogs used in this study do not have substituents of the size and character of raloxifene and consequently less likely to physically reposition helix 12. However, X-ray crystallography and sequence analysis comparison of the ligand-binding domains of ER α and ER β suggest that the agonist orientation of helix 12 in ER β may be unstable and thus easier to antagonize than ER α (Pike et al., 1999). The ER α agonist/ER β antagonists identified in this study may be able to stabilize helix 12 in the agonist orientation for ER α but not for ER β . X-ray crystallographic studies are needed to confirm this hypothesis.

The *R,R*-enantiomer of tetrahydrochrysene (*R,R*-THC) has also recently been shown to have differential ER α and ER β activity (Meyers et al., 1999; Sun et al., 1999). Like HPTE, *R,R*-THC behaves as an ER α agonist and an ER β antagonist. In contrast, the *S,S*-enantiomer (*S,S*-THC) is an agonist with both ER α and ER β . The equilibrium dissociation value (K_D) for *R,R*-THC has not been determined, and whether *R,R*-THC is an ER β competitive antagonist remains to be demonstrated. THC compounds differ considerably in structure from the methoxychlor analogs presented in this study, and this class of compounds will provide additional information regarding the ligand specificity of ER α and ER β binding and transactivation.

Less is known about the mechanism of AR antagonism by AR ligands. AR antagonists are generally thought to prevent




Fig. 3. Effect of various concentrations of methoxychlor analogs on an estradiol dose-response curve with ER β . Experiments were performed as described in Fig. 1 with 10^{-10} to 10^{-5} M E2 either alone or in combination with the indicated concentrations of monohydroxy-methoxychlor (A), monohydroxy-DDE (B), and dihydroxy-DDE (C). Luciferase activity was normalized to β -galactosidase activity. Values represent the means of three separate experiments.




Fig. 4. Effect of various concentrations of methoxychlor analogs on a DHT dose-response curve with AR. Experiments were performed as described in Fig. 1 with 10^{-9} to 10^{-5} M DHT either alone or in combination with the indicated concentrations of HPTE (A), bishydroxyphenylmethane (B), bishydroxyphenylethane (C), monohydroxy-DDE (D), and dihydroxy-DDE (E). Luciferase activity was normalized to β -galactosidase activity. Values represent the means of three separate experiments.

or reduce binding of AR to DNA (Kelce et al., 1995, 1998). However, the specific mechanisms responsible for this inhibition of AR-DNA binding remain unknown.

Much still remains to be determined about the precise roles of ER α , ER β , and AR in reproductive development and endocrine function, especially in humans; the physiological consequences of exposure to chemicals that are ER α agonists, ER β antagonists, and AR antagonists are unknown. HPTE and its structural analogs give us further insights into the ligand specificity of ER α , ER β , and AR and serve as model chemicals for investigating ER α , ER β , and AR steroid hormone receptor interactions.

Acknowledgments

We thank Dr. Paul Foster, Dr. Chris Corton, and Dr. Katrina Waters for critical review of the manuscript and Dr. Barbara Kuyper for editorial assistance.

References

- Alm H, Tiemann U and Torner H (1996) Influence of organochlorine pesticides on development of mouse embryos in vitro. *Reprod Toxicol* **10**:321–326.
- Brzozowski AM, Pike A, Dauter Z, Hubbard RE, Bonn T, Engstrom O, Ohman L, Greene GL, Gustafsson JA and Carlquist M (1997) Molecular basis of agonism and antagonism in the oestrogen receptor. *Nature (Lond)* **389**:753–758.
- Bulger WH, Feil VJ and Kupfer D (1985) Role of hepatic monooxygenases in generating estrogenic metabolites from methoxychlor and from its identified contaminants. *Mol Pharmacol* **27**:115–124.
- Bulger WH, Muccitelli RM and Kupfer D (1978) Studies on the in vivo and in vitro

- estrogenic activities of methoxychlor and its metabolites. Role of hepatic monooxygenase in methoxychlor activation. *Biochem Pharmacol* **27**:2417–2423.
- Chapin RE, Harris MW, Davis BJ, Ward SM, Wilson RE, Mauney MA, Lockhart AC, Smialowicz RJ, Moser VC, Burka LT, Collins BJ, Haskins EA, Allen JD, Judd L, Purdie WA, Harris HL, Lee CA and Corniffe GM (1997) The effects of perinatal/juvenile methoxychlor exposure on adult rat nervous, immune, and reproductive system function. *Fundam Appl Toxicol* **40**:138–157.
- Cummings AM (1997) Methoxychlor as a model for environmental estrogens. *Crit Rev Toxicol* **27**:367–379.
- Darimont BD, Wagner RL, Apriletti JW, Stallcup MR, Kushner PJ, Baxter JD, Fletterick RJ and Yamamoto KR (1998) Structure and specificity of nuclear receptor-coactivator interactions. *Genes Dev* **12**:3343–3356.
- Gaido KW, Leonard LS, Maness SC, Hall JM, McDonnell DP, Saville B and Safe S (1999) Differential interaction of the methoxychlor metabolite 2,2-bis-(p-hydroxyphenyl)-1,1,1-trichloroethane with estrogen receptors alpha and beta. *Endocrinology* **140**:5746–5753.
- Gould JC, Leonard LS, Maness SC, Wagner BL, Conner K, Zacharewski T, Safe S, McDonnell DP and Gaido KW (1998) Bisphenol A interacts with the estrogen receptor alpha in a distinct manner from estradiol. *Mol Cell Endocrinol* **142**:203–214.
- Gray LE Jr, Ostby J, Ferrell J, Rehnberg G, Linder R, Cooper R, Goldman J, Slott V and Laskey J (1989) A dose-response analysis of methoxychlor-induced alterations of reproductive development and function in the rat. *Fundam Appl Toxicol* **12**:92–108.
- Hall JM and McDonnell DP (1999) The estrogen receptor beta-isoform (ERbeta) of the human estrogen receptor modulates ERalpha transcriptional activity and is a key regulator of the cellular response to estrogens and antiestrogens. *Endocrinology* **140**:5566–5578.
- Hall DL, Payne LA, Putnam JM and Huet-Hudson YM (1997) Effect of methoxychlor on implantation and embryo development in the mouse. *Reprod Toxicol* **11**:703–708.
- Kapoor IP, Metcalf RL, Nystrom RF and Sangha GK (1970) Comparative metabolism of methoxychlor, methiochlor, and DDT in mouse, insects, and in a model ecosystem. *J Agric Food Chem* **18**:1145–1152.
- Kelce WR, Gray LE and Wilson EM (1998) Antiandrogens as environmental endocrine disruptors. *Reprod Fertil Dev* **10**:105–111.
- Kelce WR, Stone CR, Laws SC, Gray LE, Kemppainen JA and Wilson EM (1995) Persistent DDT metabolite p,p'-DDE is a potent androgen receptor antagonist. *Nature (Lond)* **375**:581–585.
- Kuiper GGJM, Carlsson B, Grandien K, Enmark E, Haggblad J, Nilsson S and Gustafsson JA (1997) Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* **138**:863–870.
- Kuiper G, Enmark E, Peltouhikko M, Nilsson S and Gustafsson JA (1996) Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci USA* **93**:5925–5930.
- Kuiper G and Gustafsson JA (1997) The novel estrogen receptor-beta subtype: Potential role in the cell- and promoter-specific actions of estrogens and antiestrogens. *FEBS Lett* **410**:87–90.
- Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, van der Burg B and Gustafsson JA (1998) Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* **139**:4252–4263.
- Maness SC, McDonnell DP and Gaido KW (1998) Inhibition of androgen receptor-dependent transcriptional activity by DDT isomers and methoxychlor in HepG2 human hepatoma cells. *Toxicol Appl Pharmacol* **151**:135–142.
- Meyers MJ, Sun J, Carlson KE, Katzenellenbogen BS and Katzenellenbogen JA (1999) Estrogen receptor subtype-selective ligands: asymmetric synthesis and biological evaluation of cis- and trans-5,11-dialkyl-5,6,11,12-tetrahydrochrysenes. *J Med Chem* **42**:2456–2468.
- Mosselman S, Polman J and Dijkema R (1996) ER beta: identification and characterization of a novel human estrogen receptor. *FEBS Lett* **392**:49–53.
- Ogawa S, Inoue S, Watanabe T, Hiroi H, Orimo A, Hosoi T, Ouchi Y and Muramatsu M (1998) The complete primary structure of human estrogen receptor beta (hER beta) and its heterodimerization with ER alpha in vivo and in vitro. *Biochem Biophys Res Commun* **243**:122–126.
- Ousterhout J, Struck RF and Nelson JA (1981) Estrogenic activities of methoxychlor metabolites. *Biochem Pharmacol* **30**:2869–2871.
- Pike ACW, Brzozowski AM, Hubbard RE, Bonn T, Thorsell AG, Engstrom O, Ljunggren J, Gustafsson JK and Carlquist M (1999) Structure of the ligand-binding domain of oestrogen receptor beta in the presence of a partial agonist and a full antagonist. *EMBO J* **18**:4608–4618.
- Saunders P, Maguire SM, Gaughan J and Millar MR (1997) Expression of oestrogen receptor beta (ER beta) in multiple rat tissues visualised by immunohistochemistry. *Endocrinology* **154**:R13–16.
- Shelby MD, Newbold RR, Tully DB, Chae K and Davis VL (1996) Assessing environmental chemicals for estrogenicity using a combination of in vitro and in vivo assays. *Environ Health Perspect* **104**:1296–1300.
- Sun J, Meyers M, Fink B, Rajendran R, Katzenellenbogen J and Katzenellenbogen B (1999) Novel ligands that function as selective estrogens or antiestrogens for estrogen receptor alpha or estrogen receptor beta. *Endocrinology* **140**:800–804.
- Tremblay GB, Tremblay A, Copeland NG, Gilbert DJ, Jenkins NA, Labrie F and Giguere V (1997) Cloning, chromosomal localization, and functional analysis of the murine estrogen receptor beta. *Mol Endocrinol* **11**:353–365.
- Tzukerman MT, Esty A, Santiso-Mere D, Danielian P, Parker MG, Stein RB, Pike JW and McDonnell DP (1994) Human estrogen receptor transactivational capacity is determined by both cellular and promoter context and mediated by two functionally distinct intramolecular regions. *Mol Endocrinol* **8**:21–30.

Send reprint requests to: Dr. Kevin W. Gaido, CIIT, P.O. Box 12137, Research Triangle Park, NC 27709. E-mail: gaido@ciit.org
