

Alkenes from Epoxides by Reductive Elimination with Magnesium Bromide-Magnesium Amalgam

By F. BERTINI, P. GRASSELLI, and G. ZUBIANI
(Istituto di Chimica del Politecnico, Milano)

and G. CAINELLI*

(Istituto di Chimica Organica, Università, Via Amendola, 173, 70126 Bari, Italy)

Summary Olefins may be obtained by treating epoxides in tetrahydrofuran solution with magnesium amalgam and magnesium bromide.

Olefins are produced by this process from a variety of epoxides (Table). The yields are generally of the order of

TABLE*


				% Yield ^b
1,2-Epoxy-1-phenylethane	32
1,2-Epoxy-1,1-diphenylethane	38
1,2-Epoxy-1-n-undecylethane	22
1,2-Epoxycyclohexane	50
3,3-Epoxymethylene-5 α -cholestane ^c	80

* All products completely characterised.

^b Yield of isolated, purified compounds.

^c A mixture of isomers was employed.


β -SUBSTITUTED alkyl halides readily undergo a reductive elimination which can be represented as:



X = halogen; Y = OH, OR, OCOR.

The best reducing agents for such systems are metals, *e.g.* sodium,¹ zinc,² and magnesium.³

On the other hand, epoxides react under appropriate conditions with metal halides, especially magnesium bromide in ether, to give, in many cases, the corresponding halohydrin salts:⁴



We have observed that epoxides readily undergo reductive elimination on being stirred overnight with an equimolar amount of magnesium bromide in tetrahydrofuran or 1,2-dimethoxyethane in the presence of slightly more than 1 g.-atom of a dilute magnesium amalgam (0.5–3% Mg) at room temperature or, if necessary, at reflux, under nitrogen or argon.

30–40%. However, no effort has been made to increase the yields, and in many cases a considerable amount of starting material was recovered.

The magnesium bromide etherate-magnesium amalgam mixture can easily be prepared by dissolving (under argon) in the appropriate quantity of mercury, 2.1 g.-atoms of magnesium, covering the resulting amalgam with solvent, and then treating, if necessary with cooling, with a mole of bromine.

The magnesium salt of a bromohydrin, resulting from the interaction between epoxide and magnesium bromide etherate, is most probably an intermediate in this reaction. This salt subsequently undergoes a magnesium-induced reductive elimination to the corresponding olefins.

(Received, November 10th, 1969; Com. 1712.)

¹ H. House and R. Ro, *J. Amer. Chem. Soc.*, 1958, **80**, 182.

² L. Fieser and R. Ettore, *J. Amer. Chem. Soc.*, 1953, **75**, 1700; L. Crombie and S. Harper, *J. Chem. Soc.*, 1950, 1705, 1715; 1956, 136; S. Cristol and L. Rademacher, *J. Amer. Chem. Soc.*, 1959, **81**, 1600.

³ R. Fuson, "Advanced Organic Chemistry," Wiley, New York, 1950, p. 144.

⁴ Cf. for instance, E. T. McBee, C. E. Hathaway, and C. W. Roberts, *J. Amer. Chem. Soc.*, 1956, **78**, 3851; Houben-Weyl "Methoden der organischen Chemie," Thieme Verlag, Stuttgart, vol. 6/3, p. 437; C. A. Stewart and C. A. Vanderwerf, *J. Amer. Chem. Soc.*, 1954, **76**, 1259.