


This article was downloaded by: [Istanbul Universitesi Kutuphane ve Dok]

On: 22 December 2014, At: 12:18

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/lcar20>

A CONVENIENT MULTIGRAM PREPARATION OF FUNCTIONALIZED 2-AZIDO-2-DEOXY-D-MANNOSE AS A USEFUL ORTHOGONALLY PROTECTED BUILDING BLOCK FOR OLIGOSACCHARIDE SYNTHESIS

Veronica Draghetti^a, Laura Poletti^a, Davide Prospero^a & Luigi Lay^a

^a Università degli Studi di Milano, Via Venezian, Milano, 21-20131, Italy

Published online: 19 Aug 2006.

To cite this article: Veronica Draghetti, Laura Poletti, Davide Prospero & Luigi Lay (2006) A CONVENIENT MULTIGRAM PREPARATION OF FUNCTIONALIZED 2-AZIDO-2-DEOXY-D-MANNOSE AS A USEFUL ORTHOGONALLY PROTECTED BUILDING BLOCK FOR OLIGOSACCHARIDE SYNTHESIS, *Journal of Carbohydrate Chemistry*, 20:9, 813-819

To link to this article: <http://dx.doi.org/10.1081/CAR-100108658>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

**A CONVENIENT MULTIGRAM PREPARATION
OF FUNCTIONALIZED 2-AZIDO-2-DEOXY-D-
MANNOSE AS A USEFUL ORTHOGONALLY
PROTECTED BUILDING BLOCK FOR
OLIGOSACCHARIDE SYNTHESIS**

**Veronica Draghetti, Laura Poletti, Davide Prospero,
and Luigi Lay***

Università degli Studi di Milano, Dip. di Chimica Organica e
Industriale, Via Venezian, 21–20131 Milano, Italy

ABSTRACT

Multigram preparation of strategically protected 2-azido-2-deoxy mannose **5** from pentaacetyl glucose **1** is described. This manno-derivative was obtained in a straightforward manner and in high overall yield and represents a flexible building-block for complex oligosaccharide synthesis.

INTRODUCTION

Mannosamine is a 2-deoxy-2-aminosugar playing different biological functions. It is an intermediate in the biosynthesis of *N*-acetyl D-neuraminic acid,¹ and it acts as inhibitor of GPI biosynthesis² and platelet aggregation factors in mammalian cells.³ It is also, *inter alia*, a component of cell walls and of the capsular polysaccharides of different bacteria.⁴ Due to its role in recognition phenomena, it represents a crucial building block in the preparation of natural and modified carbohydrate-based therapeutics. Complex carbohydrate synthesis often needs large amounts of strategically protected monosaccharides as intermediates. It is there-


*Corresponding author. Fax +39 02 2664952; E-mail: luigi.lay@unimi.it

fore highly desirable to have a quick, simple and straightforward procedure to provide a properly protected mannosamine precursor in multi-gram quantities, and suitable for utilisation in the synthesis of *N*-acetyl mannosamine-containing oligosaccharides.

In a project aimed at synthesising analogues of bacterial capsular polysaccharides, we required a synthetic way of providing multi-gram amounts of protected mannosamine precursor **5** (Scheme 1), to be used as a building block in further steps of the synthesis. To date, only two methods have been applied to the preparation of a large amount of functionalised 2-azido-2-deoxy-D-mannosamine. Paulsen *et al.*^{5b} adopted a poorly stereoselective azido nitration⁵ on D-glucal, yielding a 2:1 manno:gluco mixture. Augè *et al.*⁶ obtained 2-azido-2-deoxy-D-mannose in three steps from glucose, using *N,N'*-sulfuryl diimidazole as a key reagent. Nevertheless, this preparation furnishes very low overall yields due to the abundant formation of by-products in the early steps. Other procedures leading to mannosamine have also been reported, such as azido phenylselenylation⁷ of D-glucal and electrophilic azidation of aldonolactones.⁸ Unfortunately, they show some disadvantages such as low yields or poor stereoselectivities, use of undesired toxic reagents, or reaction conditions which make it difficult to scale-up for a multigram preparation. Finally, Ishido *et al.*⁹ reported the nucleophilic displacement of a 2-*O*-triflate of D-glucose as a method to obtain 2-azido-2-deoxy-D-manno derivatives with high selectivity and yields. Therefore, we envisaged this reaction as the key step to prepare our target molecule on a large scale.


RESULTS

In this endeavour, we developed a very simple and efficient procedure leading in eight steps to protected mannosamine precursor **5** in 39% overall yield from 1,2,3,4,6-penta-*O*-acetyl- α -D-glucopyranose **1** (Scheme 1). In compound **5** the



Scheme 1. a) 33% HBr in AcOH, CH₂Cl₂, 0°C to rt, 2 h; b) *sym*-collidine, 5 eq MeOH dry, 0.5 eq TBAB, 20 h, (92% o.y.); c) 0.2 eq MeONa 1M, MeOH, 14 min.; d) 9 eq BnBr, 4.5 eq NaH, DMF dry, 24 h (77% from **2**); e) Allylic alcohol, 0.22 eq TMSOTf, 0°C, 15'; f) 0.2 eq MeONa, MeOH, 2 h (85% from **3**); g) 4 eq Py, 3 eq Tf₂O, CH₂Cl₂ dry, 0°C, 2 h; h) 3 eq Bu₄N⁺N₃⁻, dry toluene, 55–60°C, 24 h (65% from **4**).





Scheme 2. a) HOAc/H₂O = 20:1 v/v, 2.4 eq NaOAc, 1.1 eq PdCl₂, 76%; b) Ac₂O/HOAc 2:1 v/v, 3 eq ZnCl₂, 84%.

amino group is masked by an azido group, widely used in the precursors of 2-deoxy-2-aminosugars for its poor steric hindrance and easy reversion to an amino group. Our procedure utilises inexpensive reagents, requires little chromatographic purification and leads to the strategically protected azidomannose **5** on a multigram scale. In addition, compound **5** is a versatile building block, as it can be elongated from either its reducing or nonreducing end (Scheme 2).

1,2,3,4,6-Penta-*O*-acetyl- α -D-glucopyranose **1** was obtained by conventional *O*-acetylation of D-glucose. Compound **1** was transformed into the known orthoester **2** (**10**) (33% HBr in acetic acid, then *sym*-collidine and MeOH in dichloromethane) in 92% overall yield. After Zemplén¹¹ deacetylation, reaction with benzyl bromide and sodium hydride yielded tribenzyl intermediate **3** (77% from **2**). The orthoester was then opened in allyl alcohol with trimethylsilyl triflate as a catalyst; subsequent 2-*O* deprotection with sodium methylate in methanol afforded allyl glycoside **4**¹² in 85% over two steps. The 2-OH was then activated as a triflate with triflic anhydride in dichloromethane, followed by nucleophilic displacement with tetrabutylammonium azide, providing target azidomannose **5** in 65% yield from **4** (39% o.y. from **1**). Compound **5** is a flexible monosaccharidic building block for oligosaccharide synthesis. In fact, it was easily deprotected at the anomeric position with PdCl₂ affording 2-azido-3,4,6-tri-*O*-benzyl-2-deoxy-D-mannopyranose (**6**, 76%), which can be employed as a glycosyl donor upon proper activation of the anomeric hydroxy group. On the other hand, chemoselective acetylation of **5** with acetic acid and ZnCl₂ provided 6-*O*-acetyl azidomannose **7** (84%), allowing the elongation from the nonreducing end of the sugar.

CONCLUSIONS

In conclusion, we reported herein an easy, quick and straightforward synthetic route giving access to a strategically protected azidomannose in a multigram scale and in high overall yield. This compound is a versatile, useful tool for oligosaccharide synthesis, as demonstrated by selective deblocking of either the anomeric or the 6-*O*-position.



EXPERIMENTAL SECTION

General Methods. ^1H NMR spectra were recorded on Bruker AC 300 and Varian Gemini 200 spectrometers. Solvents were dried by standard procedures; allyl alcohol was dried over 4 Å m.s. Melting points were determined with a Büchi apparatus and are not corrected. Optical rotations were measured at room temperature (23°C) with a Perkin-Elmer 241 polarimeter. TLC was carried out on Merck Silica-gel 60 F₂₅₄ plates (0.25 mm thickness), and spots were visualized by spraying with a solution containing H₂SO₄ (31 mL), ammonium molybdate (21 g) and Ce(SO₄)₂ (1 g) in 500 mL water, followed by heating at 110°C for 5 min. Column chromatography was performed by the flash procedure using Merck Silica-gel 60 (230–400 mesh). Elemental analyses were performed using the Carlo Erba elemental analyzer 1108. Allyl alcohol was dried over m.s. 4 Å.

3,4,6-Tri-*O*-acetyl-[1,2-*O*-(1-methoxyethylidene)]- α -D-glucopyranose (2). A solution of 1,2,3,4,6-penta-*O*-acetyl-D-glucopyranose **1** (50.5 g, 129.4 mmol) in 50 mL of dichloromethane was cooled to 0°C and hydrobromic acid 33% in acetic acid (50 mL, 252 mmol) was added dropwise. The reaction was monitored by TLC (eluent hexane/EtOAc = 7:3 v/v), and after 1.5 h the mixture was neutralised with ice-cold satd NaHCO₃, the organic layer was washed with water (2 × 200 mL) and dried (Na₂SO₄). Removal of the solvent afforded the bromide intermediate as a white glassy solid (53 g). Crude bromide (53 g, 129 mmol) was dissolved under a nitrogen atmosphere in 70 mL of *sym*-collidine and warmed to 45°C. TBAB (20.7 g, 64.5 mmol) and dry methanol (26 mL, 645 mmol) were added. The reaction was monitored by TLC (eluent hexane/EtOAc = 6:4 v/v). After 16 h the solution was diluted with dichloromethane, the organic layer was washed with water (2 × 200 mL), dried over Na₂SO₄ and concentrated under reduced pressure. The residue was suspended in diethyl ether (60 mL), the excess of TBAB was filtered off, and the ethereal solution was concentrated. Chromatographic purification (eluent hexane/EtOAc = 8:2 v/v containing 1% of TEA) afforded pure compound **2** as a white syrup (43.15 g, 92%), whose spectroscopic data were in agreement with those reported in literature.¹⁰

3,4,6-Tri-*O*-benzyl-[1,2-*O*-(1-methoxyethylidene)]- α -D-glucopyranose (3). Compound **2** (43.15 g, 119 mmol) was dissolved under a nitrogen atmosphere in dry methanol (60 mL) and a 1M solution of CH₃ONa in dry methanol (23.8 mL) was added dropwise. The reaction was monitored by TLC (eluent hexane/EtOAc = 7:3 v/v and CH₂Cl₂/MeOH = 85:15 v/v), and after 1.5 h the mixture was concentrated affording 28 g of the orthoester intermediate as a yellow oil. The crude compound (28 g, 119 mmol) was dissolved in dry DMF (100 mL) under a nitrogen atmosphere. After the dropwise addition of BnBr (63.6 mL, 535 mmol), NaH (25.70 g, 1.071 mol) was added portionwise. The reaction was stirred at room temperature (TLC: hexane/EtOAc 7:3 v/v), and after 20 h the excess of NaH was quenched by slow addition of methanol under vigorous stirring. The reaction mixture was diluted with diethyl ether, the organic layer was washed with water



(2 × 100 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. A quick filtration on silica gel (eluent hexane/EtOAc = 9:1 v/v containing 1% of TEA) afforded compound **3** as a yellow syrup (46.53 g, 77%), whose spectroscopic data were in agreement with those reported in literature.¹¹

Allyl (3,4,6-Tri-*O*-benzyl)-β-D-glucopyranoside (4). Compound **3** (46.53 g, 92 mmol) was dissolved in 100 mL of dry allyl alcohol under a nitrogen atmosphere. The solution was cooled to 0°C, and TMSOTf (3.2 mL, 20.2 mmol) was added dropwise. The reaction was monitored by TLC (eluent hexane/EtOAc = 7:3 v/v). After 20 min the reaction gave two products, the 2-*O*-acetyl and the 2-hydroxy derivatives respectively, in a ratio of about 1:1. The solution was neutralised with TEA and diluted with dichloromethane, the organic layer was washed with water (2 × 100 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The residue was dissolved in 80 mL of dry methanol and a 1M solution of CH₃ONa in methanol (18.4 mL, 18.4 mmol) was added. The reaction was stirred at room temperature (TLC: hexane/EtOAc 7:3 v/v). After 20 h the mixture was neutralised with Amberlite IR-120 (H⁺ form), filtered and concentrated. The residue was purified on silica gel (eluent hexane/EtOAc = 85:15 v/v) providing compound **4** (38.3 g, 85%), whose spectroscopic data were in agreement with those reported in literature.¹²

Allyl (2-Azido-3,4,6-tri-*O*-benzyl-2-deoxy)-β-D-mannopyranoside (5). To a solution of compound **4** (38.3 g, 78.2 mmol) in dry dichloromethane (150 mL) under a nitrogen atmosphere, pyridine (25.3 mL, 313 mmol) was added. The mixture was cooled to 0°C, and after 15 min triflic anhydride (38.7 mL, 234.6 mmol) was added dropwise. After stirring 1.5 h the reaction was diluted with dichloromethane, the organic layer was washed with HCl 5% until acidic pH, then with satd NaHCO₃ until neutralisation, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude 2-*O*-triflate intermediate was dissolved under a nitrogen atmosphere in dry toluene (200 mL), Bu₄N⁺N₃⁻ (55.46 mL, 195 mmol) was added quickly, and the solution was stirred at 55°C (TLC: hexane/EtOAc 7:3 v/v). After 24 h the reaction mixture was concentrated and chromatographic purification (eluent hexane/EtOAc = 9:1 v/v) afforded compound **5** as a yellow syrup (26.17 g, 65%). [α]_D²⁰ = -36.8° (c 1, chloroform); ¹H NMR (300 MHz, CDCl₃) δ 7.43–7.04 (m, 15H, H_{Ar}), 5.89–5.76 (m, 1H, CH=CH₂), 5.28–5.22 (m, 1H, CH=CHH), 5.04–5.01 (m, 1H, CH=CHH), 4.89 (d, 1H, J = 11.1 Hz, CHHPh), 4.56–4.40 (m, 4H, 3/2 CH₂Ph, H-1), 4.38–4.14 (m, 4H, CH₂Ph, CH_{all}, H-2), 4.01 (t, 1H, J_{3,4} = J_{4,5} = 9.3 Hz, H-4), 3.93–3.85 (m, 1H, CH_{all}), 3.71 (d, 2H, H-6, H-6'), 3.48 (dd, 1H, J_{3,2} = J_{3,4} = 8.8 Hz, H-3), 3.32 (dt, 1H, J_{5,4} = 9.6 Hz, J_{5,6} = J_{5,6'} = 3.3 Hz, H-5).

Anal. Calcd for C₃₀H₃₃O₅N₃ (515.61): C, 69.88; H, 6.45; N, 8.15. Found: C, 69.73; H, 6.48; N, 8.10.

2-Azido-3,4,6-tri-*O*-benzyl-2-deoxy-α,β-D-mannopyranose (6). Compound **5** (26.17 g, 50.83 mmol) was dissolved in a HOAc/H₂O = 20:1 v/v mixture



(100 mL), then NaOAc (16.6 g, 122 mmol) and PdCl₂ (6.56 g, 56 mmol) were added. The reaction was monitored by TLC (eluent hexane/EtOAc = 7:3 v/v). After 24 h the reaction mixture was diluted with EtOAc and filtered over celite, washed with satd NaHCO₃ until neutralisation, dried (Na₂SO₄), filtered and concentrated. Purification by flash chromatography (eluent hexane/EtOAc 9:1 v/v) afforded pure compound **6** as a yellow syrup (18.37g, 76%) as a mixture of α/β anomers. The deprotection of the anomeric hydroxyl was ascertained by the disappearance of the allyl signals detected in ¹H NMR spectrum (multiplets at 5.8, 5.2 and 5.0 ppm).

Anal. Calcd for C₂₇H₂₉O₅N₃ (475.54): C, 68.19; H, 6.14; N, 8.83. Found: C, 68.31; H, 6.12; N, 8.81.

Allyl 6-O-Acetyl-2-azido-3,4-di-O-benzyl-2-deoxy- β -D-mannopyranoside (7). Compound **5** (768 mg, 1.49 mmol) was dissolved in a mixture of Ac₂O/HOAc 2:1 v/v containing freshly melted ZnCl₂ (610 mg, 4.47 mmol). After 6 h the reaction was diluted with water and EtOAc, washed with satd NaHCO₃ until neutralisation, dried (Na₂SO₄), filtered and concentrated. Flash chromatography of the crude compound (eluent hexane/EtOAc = 9:1 v/v) afforded mannopyranoside **7** as a yellow syrup (585 mg, 84%). [α]_D²⁰ = -13.8° (c 1, chloroform); ¹H NMR (200 MHz, CDCl₃) δ 7.50–7.10 (m, 10H, H_{Ar}), 6.01–5.81 (m, 1H, CH=CH₂), 5.34–5.20 (m, 2H, CH=CH₂), 4.91 (d, 1H, J = 10.8 Hz, CHHPh), 4.80–4.66 (2H, AB system, CH₂Ph), 4.59 (d, 1H, J = 10.8 Hz, CHHPh), 4.52 (s, 1H, H-1), 4.43–4.32 (m, 2H, H-6, H-6'), 4.28–4.19 (m, 1H, CH_{all}), 4.12–4.02 (m, 1H, CH_{all}), 3.95 (d, 1H, J_{2,3} = 3.2 Hz, H-2), 3.77 (t, 1H, J_{3,4} = J_{4,5} = 9.2 Hz, H-4), 3.67 (dd, 1H, J_{3,4} = 9.2 Hz, J_{3,2} = 3.2 Hz, H-3), 3.41 (ddd, 1H, J_{5,4} = 9.2 Hz, J_{5,6} = 2.6 Hz, J_{5,6'} = 5.2 Hz, H-5), 2.09 (s, 3H, OAc).

Anal. Calcd for C₂₅H₂₉O₆N₃ (467.52): C, 64.23; H, 6.25; N, 8.99. Found: C, 64.14; H, 6.26; N, 9.00.

ACKNOWLEDGMENTS

We gratefully acknowledge financial support from EU (VACNET project, CT 96-0158), CNR (Centro di Studio sulle Sostanze Organiche Naturali) and MURST (COFIN 2000, prot. MM03155477).

REFERENCES

- Rodriguez-Aparicio, L.B.; Ferrero, M.A.; Reglero, A. *N*-Acetyl-D-neuraminic acid synthesis in *Escherichia coli* K1 occurs through condensation of *N*-acetyl-D-mannosamine and pyruvate. *Biochem. J.* **1995**, *308*, 501–505. Rivila-Nuin, B.; Rodriguez-Aparicio, L.B.; Ferrero, M.A.; Reglero, A. Regulation of capsular polysialic acid biosynthesis by *N*-acetyl-D-mannosamine, an intermediate of sialic acid metabolism. *FEBS Lett.* **1998**, *426*(2), 191–195.



2. Ralton, J.E.; Milne, K.G.; Güther, M.L.; Field, R.A.; Ferguson, M.A. The mechanism of inhibition glycosylphosphatidylinositol anchor biosynthesis in *Trypanosoma brucei* by mannosamine. *J. Biol. Chem.* **1993**, *268*(32), 24183–24189.
3. Kinlough-Rathbone, R.L.; Packham, R.A.; Mustard, J.F. Effect of amino sugars on platelet aggregation and on fibrinogen binding. *Thromb. Haemost.* **1984**, *52*(1) 75–80.
4. Yoneyama, T.; Koike, J.; Arakawa, H.; Yokoyama, K.; Sasaki, J.; Kawamura, T.; Ito, E.; Takao, S. Distribution of mannosamine and mannosaminuronic acid among cell walls of *Bacillus* species. *J. Bacteriol.* **1982**, *149*(1), 15–21. Lee, C.J.; Koizumi, K.; Henrichsen, J.; Perch, B.; Lin, C.S.; Egan, W. Capsular polysaccharide of non-groupable streptococci that cross-react with pneumococcal group 19. *J. Immunol.* **1984**, *133*(5), 2706–2711. Zareba, T.W.; Pascu, C.; Hryniewicz, W.; Wadström, T. Binding of extracellular matrix protein by enterococci. *Curr. Microbiol.* **1997**, *34*(1), 6–11.
5. a) Kinzy, W.; Schmidt, R.R. Synthese des Trisaccharids aus der “Repeating Unit” des Kapselpolysaccharids von *Neisseria meningitidis* (Serogruppe L). *Liebigs Ann. Chem.* **1985**, 1537–45. b) Paulsen, H.; Lorentzen, J.P.; Kutschker, W. Erprobte Synthese von 2-Azido-2-desoxy-D-Mannose und 2-Azido-2-desoxy-D-Mannuronsäure als Baustein zum Aufbau von Bakterien-Polysaccharid-Sequenzen. *Carbohydr. Res.* **1985**, *136*, 153–176.
6. Augé, C.; David, S.; Malleron, A. An inexpensive route to 2-azido-2-deoxy-D-mannose and its conversion into an azido analog of *N*-acetylneuraminic acid. *Carbohydr. Res.* **1989**, *188*, 201–205.
7. Santoyo-González, F.; Calvo-Flores, F.G.; García-Mendoza, P.; Hernández-Mateo, F.; Isac-García, J.; Robles-Díaz, R. Synthesis of phenyl 2-azido-2-deoxy-1-selenoglycosides from glycals. *J. Org. Chem.* **1993**, *58*, 6112–25. Czernecki, S.; Ayadi, E.; Randriamandimby, D. Selenoglycosides. 3. Synthesis of phenyl 2-(*N*-acetylamino)- and 2-azido-2-deoxy-1-seleno- α -D-glycopyranosides via azido-phenylselenylation of diversely protected glycals. *J. Org. Chem.* **1994**, *59*, 8256–8260.
8. Dupradeau, F.-Y.; Hakomori, S.; Toyokuni, T. Electrophilic azidation of 2-deoxy-aldono-1,5-lactones: an alternative route to 2-azido-2-deoxy-aldopyranoses. *J. Chem. Soc., Chem. Commun.* **1995**, 221–22.
9. Ishido, Y.; Sakairi, N. Regioselective *O*-deacetylation of fully acylated glycosides and 1,2-*O*-isopropylidenealdofuranose derivatives with hydrazine hydrate. *Carbohydr. Res.* **1981**, *97*, 151–155.
10. Alfredsson, G. Borèn, H.B.; Garegg, P.J. 3-*O*- α -D-glucopyranosyl-D-mannose. *Acta Chem. Scand.* **1972**, *26*, 2531.
11. Ogawa, T.; Nozachi, M.; Matsui, M. Total synthesis of stevioside. *Tetrahedron* **1980**, *36*, 2641–2648.
12. Ogawa, T.; Takanashi, Y. Synthesis of β -D-(1 \rightarrow 2)-linked D-glucopentaose, a part of the structure of the exocellular β -D-glucan of *Agrobacterium tumefaciens*. *Carbohydr. Res.* **1983**, *123*, C16–C18.

Received April 9, 2001

Accepted October 11, 2001



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081CAR100108658>