

# Borinic Acid Catalyzed, Regioselective Chloroacylations and Chlorosulfonylations of 2,3-Epoxy Alcohols

Kashif Tanveer, Kareem Jarrah, and Mark S. Taylor\*

Department of Chemistry, University of Toronto, 80 St. George Street, Toronto, Ontario M5S 3H6, Canada

## **(5)** Supporting Information

**ABSTRACT:** In the presence of a borinic acid derived catalyst, 2,3epoxy alcohols undergo couplings with acyl and sulfonyl chlorides. This transformation directly generates *O*-acylated or *O*-sulfonylated chlorohydrin diols, with significant levels of regioselectivity for both the ring-opening and *O*-functionalization steps.



2,3-Epoxy alcohols are versatile building blocks that are readily accessible from allylic alcohols using directed epoxidation methods, including enantioselective variants.<sup>1</sup> While presenting diverse opportunities for synthetic manipulations, the 2,3-epoxy alcohol motif presents challenges in regiocontrol. The problem of regioselective epoxy alcohol ring opening (at C-2 versus C-3) has been studied in depth, and protocols based on the use of Lewis acid promoters and/or electronically biased substrates have been developed.<sup>2</sup> New developments continue to emerge, including the recently reported catalytic methods for ring opening of 2,3-epoxy alcohols.<sup>3</sup> After epoxide opening, a second regioselective transformation is often required to differentiate OH groups in the 1,2- or 1,3-diol product. Chemists have generally tackled these problems in regiocontrol independently, using two-step reaction sequences. Herein, we describe organoboron-catalyzed couplings of acyl or sulfonyl chlorides with 2,3epoxy alcohols. These transformations directly generate Oacylated or O-sulfonylated chlorohydrin diols, with significant levels of regioselectivity for both the ring-opening and Ofunctionalization steps.

The present results build on our group's finding that borinic acids (R<sub>2</sub>BOH) serve as catalysts for regioselective activation of OH groups in 1,2- and 1,3-diol motifs (Scheme 1A).<sup>4</sup> This mode of catalysis can be used to achieve monoacylations, sulfonylations, alkylations, and glycosylations of di- and triols, including carbohydrate derivatives. Mechanistic studies suggest that the regioselectivity-determining step is the reaction of the electrophile with a tetracoordinate borinic ester generated by reversible covalent binding of a substrate-derived diol group. We speculated that a borinic ester of this type could be generated in an alternative way by borinic acid promoted opening of a 2,3-epoxy alcohol. Precedent for this idea includes the work of Miyashita and co-workers, who employed boronic acids in stoichiometric amounts as promoters of C2-selective ring openings of 2,3-epoxy alcohols (Scheme 1B).<sup>5</sup> If a borinic acid were to accelerate the ring-opening step in a similar way, then the reaction of the resulting borinic ester with an electrophile could close the catalytic cycle, delivering a bis-functionalized product (Scheme 1C).

Preliminary experiments revealed that treatment of *cis*-epoxy alcohol **2a** with benzoyl chloride (BzCl) in the presence of

Scheme 1. Proposed Strategy for Regioselective Ring Opening and Monofunctionalization of 2,3-Epoxy Alcohols





precatalyst **1a** and diisopropylethylamine, conditions that we have employed previously for monoacylations of diols, led to the formation of monobenzoylated chlorohydrin **3a** in 71% yield along with epoxy ester **5a** (Table 1).<sup>3c,6</sup> The *syn* configuration of **3a** was established by X-ray crystallography. In comparison to methods for addition of "silyl–X" reagents across epoxides (e.g., halosilylations,<sup>7</sup> azidosilylations,<sup>8</sup> and cyanosilylations<sup>9</sup>), additions of "acyl–X" are under-developed. The closest precedent for this type of reactivity is the organotin dichloride catalyzed chloroacylation of epoxides reported by Shibata and coworkers.<sup>10</sup>

The effects of varying the conditions for the reaction of **2a** with benzoyl chloride are summarized in Table 1. In addition to epoxy ester **5a**, the C2-chlorinated regioisomer **4a** was observed as a side product under certain conditions. A survey of bases revealed that 1,8-bis(dimethylamino)naphthalene (Proton-sponge) provided superior results to *i*-Pr<sub>2</sub>NEt, whereas significantly lower yields of **3a** were obtained using Et<sub>3</sub>N, pyridine, and K<sub>2</sub>CO<sub>3</sub>. In the absence of catalyst, chloroacylation of the epoxy alcohol was

 Received:
 May 26, 2015

 Published:
 June 26, 2015

# Table 1. Optimization of Conditions for Chloroacylation of Epoxy Alcohol 2a



<sup>*a*</sup>Yields were determined by <sup>1</sup>H NMR spectroscopy analysis of unpurified reaction mixtures using 1,3,5-trimethoxybenzene as an internal standard. <sup>*b*</sup>Bu<sub>4</sub>N<sup>+</sup>Cl<sup>-</sup> (1.0 equiv) was added. <sup>*c*</sup>0.2 equiv of Proton-sponge (1,8-bis(dimethylamino)naphthalene) were used.

not observed, with the product of direct acylation being generated in 24% yield. Other Lewis acid catalysts  $(PhB(OH)_2)$  $CeCl_3 \cdot 7H_2O_1$ ,  $MgI_{21}$ , and  $Cu(OAc)_2$ ) were ineffective in promoting the chloroacylation reaction. The fact that free diphenylborinic acid also catalyzed the formation of 3a is consistent with the hypothesis that ethanolamine ester 1a acts a precatalyst. Our previous studies of diol acylation suggest that N,O-bis-benzovlation of the ethanolamine ligand triggers entry of 1a into the catalytic cycle. The amine hydrochloride salt generated upon ligand benzovlation could serve as the initial source of Cl<sup>-</sup> for the epoxide ring opening. It may be that the lower yield of 3a obtained using Ph2BOH instead of borinic ester 1a is related to the different Cl<sup>-</sup> concentrations generated using these two precatalysts. We have previously reported that heteraborinine-derived borinic acid 1b displays enhanced activity for certain diol activation reactions, likely due to the high nucleophilicity of its derived borinic esters. This catalyst was inactive under the standard reaction conditions and displayed only modest activity in the presence of Bu<sub>4</sub>N<sup>+</sup>Cl<sup>-</sup>. We speculate that the lower Lewis acidity of 1b relative to Ph<sub>2</sub>BOH results in a reduced rate of epoxide ring opening. However, electrondeficient catalyst 1c did not provide improved results relative to 1a. The optimized reaction conditions thus involved the use of 1a as the catalyst and Proton-sponge as the base: we found that the loading of the latter could be reduced from 1.0 to 0.2 equiv without a deleterious effect on the yield.

The chloroacylation protocol was applied to a range of substituted epoxy alcohols (Scheme 2). Depending on the substrate used, either Proton-sponge (0.2 equiv) or *i*-Pr<sub>2</sub>NEt (1.5 equiv) as the base provided the best results. The regioselectivity data in Scheme 2 are based on analysis of unpurified reaction

Scheme 2. Chloroacylations of Epoxy Alcohols Catalyzed by  $1a^a$ 



<sup>*a*</sup>Regioselectivities were determined by <sup>1</sup>H NMR spectral analysis prior to purification. Isolated yields after purification are listed. Conditions *a*: **1a** (10 mol %), RCOCl (1.5 equiv), Proton-sponge (0.2 equiv) CH<sub>2</sub>Cl<sub>2</sub>, 23 °C. Conditions *b*: **1a** (10 mol %), BzCl (1.5–3.0 equiv), *i*-Pr<sub>3</sub>NEt (1.5 equiv), CH<sub>2</sub>Cl<sub>2</sub>, 23 °C.

mixtures by <sup>1</sup>H NMR spectroscopy. Isolated yields of the major product after purification by silica gel chromatography are listed. *cis*-2,3-Epoxycyclohexanol derivatives 2b-2d underwent C3selective epoxide ring opening and benzoylation at O-1. The *trans* relationship between Cl and OH groups was confirmed for product 3c by X-ray crystallography. Whereas 1-O-benzoylation was anticipated on steric grounds for product 3a (benzoylation of the primary over the secondary OH group being favored), rationalizing the site of acylation in products 3b-3d is less straightforward. Catalyst 1a is known to activate the equatorial OH groups of *cis*-1,2-diol motifs in pyranosides.<sup>4</sup> It is unclear

#### **Organic Letters**

whether this effect is at play in the synthesis of 3b-3d, or whether the preference for 1-O-benzoylation results from deactivation of O-2 by the inductively electron-withdrawing chloro substituent. Tertiary alcohol 2e was unreactive with BzCl, but the use of a less hindered reagent (acetyl chloride) enabled preparation of the 2-O-acylated chlorohydrin 3e. Steric effects are likely responsible for the anomalous regiochemical outcome of this reaction (acylation at O-2 rather than O-1). A *cis*-2,3epoxycyclopentanol derivative (2f) also underwent efficient chloroacylation in the presence of 1a.

The reactions of substrates 2g-2j, derived from acyclic allylic alcohols, illustrate effects of the identity and position of epoxide substituents on the borinic acid catalyzed chloroacylation reaction. 2-Substituted 2g and 2,3-disubstituted 2h were converted to the products of 1,3-chloroacylation with high levels of regiocontrol. However, the ring opening of benzyloxysubstituted cis-epoxy alcohol 2i was less selective than that of unfunctionalized 2a. Sharpless and co-workers reported that 2,3epoxy alcohols, ethers, and acetals undergo C3-selective ring openings due to the inductive electron-withdrawing effect of the oxygenated substituents.<sup>2a</sup> In the case of **2i**, the CH<sub>2</sub>OH and CH<sub>2</sub>OBn substituents exert opposing effects, thus leading to a mixture of C2- and C3-chloro products. It should be noted that the magnitude of the effect observed by Sharpless and co-workers was rather modest, with regioisomer ratios generally not exceeding 4:1. Thus, the borinic acid catalyst acts to enhance the level of selectivity for attack of chloride at C-3. trans-Epoxy alcohol 2j displayed diminished regioselectivity in the ringopening step relative to its cis isomer 2a. Given that Lewis acid promoted, C3-selective ring openings of trans-epoxy alcohols have been reported, a rationale for this difference in selectivities is not apparent. Although the chloroacylation of 2j was not highyielding, this transformation demonstrated that the antidiastereomer 3j was not formed in detectable amounts from the reaction of substrate 2a (and likewise, that syn-configured 3a was not generated from 2j). Indeed, diastereomeric side products were not evident in any of the reactions shown in Scheme 2, suggesting an S<sub>N</sub>2-type inversion as has been documented for other Lewis acid mediated protocols for preparation of halohydrins from epoxides.

The protocol was applied successfully to 3,4-epoxy alcohol 2k. The C3-ring-opened product was obtained, as verified by analysis of the chlorine-induced isotope shift in the <sup>13</sup>C NMR spectrum.<sup>11</sup> Variation of the acylating agent was also tolerated: coupling with 4-nitrobenzoyl chloride yielded the substituted ester 3l, and bisbenzoate 3m was generated using benzoic anhydride in place of BzCl. However, some limitations were encountered in this regard: alkanoyl chloride, and benzyl chloroformates (e.g., butyryl chloride, isobutyryl chloride, and benzyl chloroformate) did not give rise to synthetically useful yields of chloroacylation using test substrate 2b. Attempted fluoroacylation of 2b using benzoyl fluoride resulted only in the formation of the corresponding epoxy ester.

The use of sulfonyl chlorides under the conditions described above resulted in the formation of monosulfonylated chlorohydrins 6a-6e (Scheme 3). Both *para*-toluenesulfonyl chloride (TsCl) and 4-nitrobenzenesulfonyl chloride (NsCl) were successfully employed in this transformation. The products possess differentiated electrophilic sites in a 1,3-relationship and present opportunities for further selective manipulations.<sup>12</sup>

To obtain preliminary information regarding the mechanism of borinic acid catalyzed chloroacylation, the experiments shown in Scheme 4 were conducted. Epoxy ester **5b** was recovered Scheme 3. Chlorosulfonylations of Epoxy Alcohols Catalyzed by  $1a^a$ 



<sup>a</sup>Regioselectivities were determined by <sup>1</sup>H NMR spectral analysis prior to purification. Isolated yields after purification are listed. Conditions *a*: **1a** (10 mol %), RSO<sub>2</sub>Cl (1.5 equiv), Proton-sponge (0.2 equiv) CH<sub>2</sub>Cl<sub>2</sub>, 23 °C. Conditions *b*: **1a** (10 mol %), RSO<sub>2</sub>Cl (1.5–3.0 equiv), *i*-Pr<sub>2</sub>NEt (1.5 equiv), CH<sub>2</sub>Cl<sub>2</sub>, 23 °C.

# Scheme 4. Chloracylation of Epoxy Alcohols: Control Experiments



unchanged when subjected to BzCl and *i*-Pr<sub>2</sub>NEt, in the presence of either catalytic or stoichiometric quantities of **1a** (eq 1). This result suggests that ring opening precedes alcohol acylation in the catalytic cycle and indicates that the hydroxyl group is necessary for epoxide activation by the borinic acid. Further support for the former point was obtained by subjecting chlorohydrin diol 7 to the reaction conditions, which resulted in the formation of monobenzoate **3b** (eq 2). In the absence of catalyst, the benzoylation of 7 proceeded at a lower rate and with a low level of selectivity for the opposite regioisomer (1.5:1 2-OBz/3-OBz). The borinic acid thus influences the selectivity of the benzoylation step. Finally, *trans*-2,3-epoxycyclohexanol **2l** was inert to the conditions of chloroacylation, presumably due to its inability to interact with the borinic acid through two-point binding (eq 3).

The experiments described above, as well as our mechanistic studies of borinic acid catalyzed reactions of diols, are consistent with the proposed catalytic cycle shown in Scheme 5. Benzoylation of the ethanolamine ligand, which is required for

# Scheme 5. Proposed Catalytic Cycle



precatalyst activation, serves to generate 2 equiv of *i*-Pr<sub>2</sub>NEt·HCl relative to **1a**. Reversible covalent interaction of the organoboron catalyst with substrate activates the epoxide toward ring opening by the released Cl<sup>-</sup>. The observed C3-selectivity is consistent with results obtained for other Lewis acid promoted reactions of epoxy alcohols, including those involving halide nucleophiles.<sup>3c,6</sup> Acylation of the formed borinic ester, and displacement of the product by epoxy alcohol, results in catalyst turnover and regenerates an equivalent of chloride.

In conclusion, an efficient, atom-economical process for regioselective chloroacylation and chlorosulfonylation of 2,3epoxy alcohols has been developed. In situ acylation or sulfonylation of the chlorohydrin diol not only results in differentiation of the two OH groups but also enables what might otherwise be a challenging catalyst turnover step. The proposed mechanism, involving sequential enhancements of the electroand nucleophilicity of the bound substrate, highlights the versatile reactivity of borinic acid complexes. Efforts to gain insight into the mechanism of this process, and to extend this mode of catalysis to other sets of reaction partners, are underway in our laboratory.

# ASSOCIATED CONTENT

#### Supporting Information

Experimental procedures, characterization data, and NMR spectral data for all new compounds. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.Sb01541.

# AUTHOR INFORMATION

## **Corresponding Author**

\*E-mail: mtaylor@chem.utoronto.ca.

#### Notes

The authors declare no competing financial interest.

# ACKNOWLEDGMENTS

Funding was provided by NSERC (Discovery Grant and Canada Research Chairs Programs, Graduate Scholarship to K.T., Undergraduate Scholarship to K.J.), Boehringer Ingelheim Ltd., and the Canada Foundation for Innovation (Projects Nos. 17545 and 19119). We are grateful to Prof. Rob Batey (Department of Chemistry, University of Toronto) for helpful suggestions.

# REFERENCES

(1) Katsuki, T.; Martin, V. Org. React. 2004, 48, 1–299.

(2) (a) Behrens, C. H.; Sharpless, K. B. J. Org. Chem. **1985**, 50, 5696– 5704. (b) Guivisdalsky, P.; Bittman, R. J. Am. Chem. Soc. **1989**, 111, 3077–3079. (c) Chini, M.; Crotti, P.; Flippin, L. A.; Gardelli, C.; Giovani, E.; Macchia, F.; Pineschi, M. J. Org. Chem. **1993**, 58, 1221– 1227. (d) Hanson, R. M. Chem. Rev. **1991**, 91, 437–475.

(3) (a) Wang, C.; Yamamoto, H. J. Am. Chem. Soc. 2014, 136, 6888–6891. (b) Uesugi, S.; Watanabe, T.; Imaizumi, T.; Shibuya, M.; Kanoh, N.; Iwabuchi, Y. Org. Lett. 2014, 16, 4408–4411. (c) Wang, C.; Yamamoto, H. Org. Lett. 2014, 16, 5937–5939.

(4) (a) Lee, D.; Taylor, M. S. J. Am. Chem. Soc. 2011, 133, 3724–3727.
(b) Lee, D.; Williamson, C. L.; Chan, L.; Taylor, M. S. J. Am. Chem. Soc. 2012, 134, 8260–8267. (c) Dimitrijević, E.; Taylor, M. S. Chem. Sci. 2013, 4, 3298–3303. (d) Taylor, M. S. Acc. Chem. Res. 2015, 48, 295–305.

(5) (a) Sasaki, M.; Tanino, K.; Hirai, A.; Miyashita, M. *Org. Lett.* **2003**, *5*, 1789–1791. (b) Tomata, Y.; Sasaki, M.; Tanino, K.; Miyashita, M. *Tetrahedron Lett.* **2003**, *44*, 8975–8977.

(6) For a review of methods for selective opening of 2,3-epoxy alcohols with halides: Bonini, C.; Righi, G. *Synthesis* **1994**, *16*, 225–238.

(7) (a) Kricheldorf, H. R.; Mörber, G.; Regel, W. Synthesis 1981, 383–384. (b) Andrews, G. C.; Crawford, T. C.; Contillo, L. G., Jr. Tetrahedron Lett. 1981, 22, 3803–3806. (c) Oriyama, T.; Ishiwata, A.; Hori, Y.; Yatabe, T.; Hasumi, N.; Koga, G. Synlett 1995, 1004–1006. (d) Stamatov, S. D.; Stawinski, J. Eur. J. Org. Chem. 2008, 2635–2643. (8) (a) Blandy, C.; Choukroun, R.; Gervais, D. Tetrahedron Lett. 1983, 24, 4189–4192. (b) Sinou, D.; Emziane, M. Tetrahedron Lett. 1986, 27, 4423–4426. (c) Martinez, L. E.; Leighton, J. L.; Carsten, D. H.; Jacobsen, E. N. J. Am. Chem. Soc. 1995, 117, 5897–5898.

(9) (a) Mullis, J. C.; Weber, W. P. J. Org. Chem. 1982, 47, 2873–2875.
(b) Imi, K.; Yanagihara, N.; Utimoto, K. J. Org. Chem. 1987, 52, 1013–1016. (c) Cole, B. M.; Shimizu, K. D.; Krueger, C. A.; Harrity, J. P. A.; Snapper, M. L.; Hoveyda, A. H. Angew. Chem., Int. Ed. Engl. 1996, 35, 1668–1671. (d) Schaus, S. E.; Jacobsen, E. N. Org. Lett. 2000, 2, 1001–1004.

(10) (a) Shibata, I.; Baba, A.; Matsuda, H. *Tetrahedron Lett.* **1986**, *27*, 3021–3024. For other examples of epoxide chloroacylation: (b) Iqbal, J.; Khan, M. A.; Srivastava, R. R. *Tetrahedron Lett.* **1988**, *29*, 4985–4986.
(c) Qian, C.; Zhu, D. *Synth. Commun.* **1994**, *24*, 2203–2214.
(d) Taniguchi, Y.; Tanaka, S.; Kitamura, T.; Fujiwara, Y. *Tetrahedron Lett.* **1998**, *39*, 4559–4560. (e) Luzzio, F. A.; Bobb, R. A. *Tetrahedron* **1999**, *55*, 1851–1858. (f) Maghaddam, R. M.; Saeidian, H.; Mirjafaray, Z.; Javan, M. J.; Farimani, M. M.; Seirafi, M. *Heteroatom Chem.* **2009**, *20*, 157–163. (g) Umeda, R.; Nishimura, T.; Kaiba, K.; Tanaka, T.; Takahashi, Y.; Nishiyama, Y. *Tetrahedron* **2011**, *67*, 7217–7221.

(11) (a) Sergeyev, N. M.; Sergeyeva, N. D.; Raynes, W. T. J. Magn. Reson., Ser. A **1995**, 115, 174–192. (b) Braddock, D. C.; Bhuva, R.; Millan, D. S.; Pérez-Fuertes, Y.; Roberts, C. A.; Sheppard, R. N.; Solanki, S.; Stokes, E. S. E.; White, A. J. P. Org. Lett. **2007**, *9*, 445–448.

(12) Synthetic applications of chlorohydrin diol monosulfonates:
(a) Ennis, M. D.; Baze, M. E. *Tetrahedron Lett.* 1986, 27, 6031–6034.
(b) Overman, L. E.; Thompson, A. S. J. Am. Chem. Soc. 1988, 110, 2248–2256. (c) Gao, L.-X.; Murai, A. *Tetrahedron Lett.* 1992, 33, 4349–4352. (d) Bedke, D. K.; Shibuya, G. M.; Pereira, A.; Gerwick, W. H.; Haines, T. H.; Vanderwal, C. D. J. Am. Chem. Soc. 2009, 131, 7570–7572.